

Clarence

WE VALUE YOUR OPINION!

WE WANT TO KNOW ABOUT YOUR SHOPPING EXPERIENCE TODAY AT WAL-MART.

Please complete a survey about today's store visit at:

<http://www.survey.walmart.com>

You will need to enter the following online:

ID #: 7FN7DZXW9H4

IN RETURN FOR YOUR TIME YOU COULD RECEIVE ONE OF FIVE \$1000 WALMART SHOPPING CARDS

Must be 18 or older and a legal resident of the 50 US or DC to enter. No purchase necessary to enter or win. To enter without purchase and for complete official rules visit www.entry.survey.walmart.com. Sweepstakes period ends on the date shown in the official rules. Survey must be taken within TWO weeks

Esta oferta también se encuentra en español en la página del Internet

THANK YOU

Walmart 
Save money. Live better.

(214) 319 - 2616
MANAGER GLEN JOHNSON
7401 SAMUELL BLVD
DALLAS TX 75228
ST# 266, OP# 00005269 TE# 79 TR# 02063
RX# 7196414 D38 QTY TH 4.00 N
SUBTOTAL 4.00
TOTAL 4.00
CASH TEND 4.00
CHANGE DUE 0.00

ITEMS SOLD 1

TC# 6198 7089 9811 6909 5840


Low prices. Every day. On everything.
Backed by our Ad Match Guarantee.
01/27/13 13.1

ABNER, CLARENCE

Rx #: 7196414

If you have any questions, please feel free to contact your pharmacist at (214)319-0857 or GREER, SHARON S at (800)893-9698

Date: 01/27/2013

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088

Directions : TAKE ONE TABLET BY MOUTH 4 TIMES DAILY AS NEEDED FOR PAIN

IBUPROFEN 600MG TAB

1644

COMMON USES: This medicine is a nonsteroidal anti-inflammatory drug (NSAID) used to treat mild to moderate pain, osteoarthritis, and rheumatoid arthritis. It may also be used to treat other conditions as determined by your doctor.

HOW TO USE THIS MEDICINE: Follow the directions for using this medicine provided by your doctor. This medicine may come with a medication guide. Read it carefully. Ask your doctor, nurse, or pharmacist any questions that you may have about this medicine. TAKE THIS MEDICINE with a full glass (8 oz./240 ml) of water. DO NOT lie down for 30 minutes after taking this medicine. The dosage is based on your medical condition and response to therapy. If repeat doses are needed, they are usually given 6 or 8 hours apart, or as directed by your doctor. When used in children, the dose is based on your child's weight. Read the product instructions to find the appropriate dose for your child's weight. Consult the pharmacist or doctor if you have questions or if you need help in choosing the appropriate dosage form. THIS MEDICINE MAY BE TAKEN WITH FOOD if it upsets your stomach. Taking it with food may not decrease the risk of stomach or bowel problems (such as bleeding or ulcers) that may occur while taking this medicine. Talk with your doctor or pharmacist if you experience persistent stomach upset. STORE THIS MEDICINE at room temperature, away from heat and light. Do not store in the bathroom. IF YOU MISS A DOSE OF THIS MEDICINE, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.

Database Edition: 131. Information Expires 04/18/2013

Walmart
Pharmacy (214)319-0857
7401 SAMUELL BLVD
DALLAS, TX 75228 -0000

ABNER, CLARENCE 01/27/2013 NEW
5721 GREEN HOLLOW THE COLONY, TX 75056
Rx #: 7196414 Ref # 0 QTY: 20 DAW: 0 DS: 5
NDC: 68645-0221-59 IBUPROFEN 600MG TAB LEG
GREER, SHARON S NABP: 4501854

Patient Pay \$ 4.00

Walmart
Pharmacy (214)319-0857
7401 SAMUELL BLVD
DALLAS, TX 75228 -0000

ABNER, CLARENCE 01/27/2013 NEW
5721 GREEN HOLLOW THE COLONY, TX 75056
Rx #: 7196414 Ref # 0 QTY: 20 DAW: 0 DS: 5
NDC: 68645-0221-59 IBUPROFEN 600MG TAB LEG
GREER, SHARON S NABP: 4501854

Patient Pay \$ 4.00

ABNER
CLARENCE

1644

Signature Required N

01/27/2013 10:40:05 AM ***

Page No : 1 of 4

Drop Off: 1/26/2013 3:42:24 PM
Ready By: 1/26/2013 5:42:00 PM

Priority: Will Pick Up

ABNER
CLARENCE
5721 GREEN HOLLOW
THE COLONY, TX 75056

01/27/2013 (214)319-0857
RX: 7196414 REF = 0



OC# 465 923 861 076 592 384 107 659 238

TOTAL: \$4.00

ABNER, CLARENCE

Rx #: 7196414

If you have any questions, please feel free to contact your pharmacist at (214)319-0857 or GREER, SHARON S at (800)893-9698


Date: 01/27/2013

Directions : TAKE ONE TABLET BY MOUTH 4 TIMES DAILY AS NEEDED FOR PAIN

IBUPROFEN 600MG TAB

1644

CAUTIONS: THIS MEDICINE INCREASES YOUR RISK OF SERIOUS STOMACH OR BOWEL PROBLEMS (such as ulcers and bleeding). This risk is increased if you are elderly or are in poor health, if you have a history of smoking or drinking alcohol, if you take corticosteroid medicines (such as prednisone) or "blood thinners" (such as warfarin), or if you take this medicine for a long period of time. THIS MEDICINE MAY ALSO INCREASE YOUR RISK for certain serious heart and blood vessel problems (such as heart attack and stroke). TAKE THIS MEDICINE EXACTLY AS PRESCRIBED BY YOUR DOCTOR, at the lowest possible dose for the shortest time needed. Talk with your doctor or pharmacist for further information. DO NOT TAKE THIS MEDICINE IF YOU HAVE HAD A SEVERE ALLERGIC REACTION to aspirin or any medicine containing aspirin or to a nonsteroidal anti-inflammatory drug (such as Feldene, Motrin, Naprosyn, Clinoril). A severe reaction includes a severe rash, hives, breathing difficulties, or dizziness. If you have a question about whether you are allergic to this medicine or if a certain medicine is a nonsteroidal anti-inflammatory drug, contact your doctor or pharmacist. DO NOT EXCEED THE RECOMMENDED DOSE or take this medicine for longer than 10 days for pain or 3 days for fever, unless directed by your doctor. Laboratory and/or medical tests, including blood counts, liver function tests, and kidney function tests, may be performed to monitor your progress or to check for side effects, especially if you are taking this medicine for a long period of time. KEEP ALL DOCTOR AND LABORATORY APPOINTMENTS while you are taking this medicine. DO NOT DRIVE, OPERATE MACHINERY, OR DO ANYTHING ELSE THAT COULD BE DANGEROUS until you know how you react to this medicine. ALCOHOL WARNING: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take this medicine or other pain relievers/fever reducers. BEFORE YOU BEGIN TAKING ANY NEW MEDICINE, either prescription or over-the-counter, check with your doctor or pharmacist. If you are also taking aspirin, as prescribed by your doctor for reasons such as heart attack or stroke prevention (usually these dosages are 81-325 mg per day), continue to take the aspirin and consult your doctor or pharmacist before using this medicine. CAUTION IS ADVISED WHEN USING THIS MEDICINE IN THE ELDERLY because they may be more sensitive to the effects of this medicine, especially the risk of stomach or bowel effects (such as bleeding or ulcers), or kidney effects. FOR WOMEN: USE OF THIS MEDICINE DURING PREGNANCY has resulted in fetal and newborn death. If you think you may be pregnant, contact your doctor immediately. THIS MEDICINE IS EXCRETED IN BREAST MILK. IF YOU ARE OR WILL BE BREAST-FEEDING while you are using this medicine, check with your doctor or pharmacist to discuss the risks to your baby.

 Information Expires 04/18/2013




ABNER, CLARENCE

Rx #: 7196414 If you have any questions, please feel free to contact your pharmacist at (214)319-0857
or GREER, SHARON S at (800)893-9698
Date: 01/27/2013

Directions : TAKE ONE TABLET BY MOUTH 4 TIMES DAILY AS NEEDED FOR PAIN

IBUPROFEN 600MG TAB 1644

POSSIBLE SIDE EFFECTS: SIDE EFFECTS, that may go away during treatment, include nausea, vomiting, diarrhea, gas, constipation, indigestion, dizziness, lightheadedness, drowsiness, or headache. If they continue or are bothersome, check with your doctor. CHECK WITH YOUR DOCTOR AS SOON AS POSSIBLE if you experience ringing in ears. CONTACT YOUR DOCTOR IMMEDIATELY if you experience rapid or pounding heartbeat; easy bruising or bleeding; very stiff neck; or mental/mood changes. CONTACT YOUR DOCTOR IMMEDIATELY if you experience sharp or crushing chest pain; sudden shortness of breath; sudden leg pain; sudden severe headache, vomiting, dizziness, or fainting; changes in vision; numbness of an arm or leg; slurred speech; one-sided weakness; sudden unexplained weight gain; change in amount of urine produced; severe or persistent stomach pain; vomit that looks like coffee grounds; black tarry stools; itching, reddened, swollen, blistered, painful, or peeling skin; yellowing of the skin or eyes; dark urine; right-sided tenderness; severe or persistent tiredness; fever, chills, or sore throat; severe or persistent nausea; swelling of hands, ankles, feet, face, lips, eyes, throat, or tongue; difficulty swallowing or breathing; or hoarseness. AN ALLERGIC REACTION TO THIS MEDICINE is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash, itching, swelling, severe dizziness, or trouble breathing. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.



ABNER, CLARENCE

Rx #: 7196414

If you have any questions, please feel free to contact your pharmacist at (214)319-0857 or GREER, SHARON S at (800)893-9698

Date: 01/27/2013

Directions : TAKE ONE TABLET BY MOUTH 4 TIMES DAILY AS NEEDED FOR PAIN

IBUPROFEN 600MG TAB


1644

BEFORE USING THIS MEDICINE: WARNING: THE RISK OF SERIOUS AND SOMETIMES FATAL HEART PROBLEMS, HEART ATTACK, AND STROKE may be increased with the use of this medicine. This risk may be increased the longer you use this medicine. Risk may also be higher in patients who have heart problems or who are at risk for heart problems. **THIS MEDICINE SHOULD NOT BE USED** to treat pain before or after coronary artery heart bypass (CABG) surgery. **THE RISK OF SERIOUS AND SOMETIMES FATAL STOMACH AND BOWEL PROBLEMS**, including bleeding, ulcers, and holes in the stomach and bowel, is increased while using this medicine. These problems may occur at any time during therapy, with or without symptoms. Elderly patients are at higher risk for serious stomach problems. Ask your doctor or pharmacist for more information about this medicine and its side effects. Some medicines or medical conditions may interact with this medicine. **INFORM YOUR DOCTOR OR PHARMACIST** of all prescription and over-the-counter medicine that you are taking. **DO NOT TAKE THIS MEDICINE** if you are also taking heparins or tacrolimus. **ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION** may be needed if you are taking serotonin reuptake blocker medicines such as fluoxetine or citalopram, "blood thinners" such as warfarin, bisphosphonates such as alendronate or risedronate, cyclosporine, corticosteroids such as prednisone, high blood pressure medicines (including ACE inhibitors such as captopril, angiotensin II receptor antagonists such as losartan, and beta-blockers such as metoprolol), "water pills" (diuretics such as furosemide, hydrochlorothiazide, triamterene), lithium, methotrexate, or aspirin. **DO NOT START OR STOP ANY MEDICINE** without doctor or pharmacist approval. Inform your doctor of any other medical conditions including poorly controlled diabetes, dehydration, heart problems (such as heart failure or history of heart attack), swelling of the hands, feet, or ankles (edema), high blood pressure, history of stroke, blood clotting problems, stomach or bowel problems (such as bleeding or ulcers), history of tobacco use or alcohol use, kidney problems, liver problems, blood or bleeding problems (such as anemia), asthma, growths in the nose (nasal polyps), any allergies (especially history of angioedema with symptoms of lip, tongue, throat swelling), pregnancy, or breast-feeding. **USE OF THIS MEDICINE IS NOT RECOMMENDED** if you have a history of allergy to aspirin or other NSAIDs (e.g., naproxen, celecoxib). **USE OF THIS MEDICINE IS NOT RECOMMENDED** if you have history of severe kidney disease or if you are going to have or have recently had coronary artery heart bypass (CABG) surgery. Contact your doctor or pharmacist if you have any questions or concerns about taking this medicine.

OVERDOSE: If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms of overdose may include severe stomach pain, coffee ground-like vomit, unusually fast or slow heartbeat, trouble breathing, extreme drowsiness, loss of consciousness, and seizures.

PATIENT-ORIENTED DISCLAIMER: This information should not be used to decide whether or not to take this medicine or any other medicine. Only your health care provider has the knowledge and training to decide which medicines are right for you. This information does not endorse any medicine as safe, effective, or approved for treating any patient or health condition. This is only a brief summary of general information about this medicine. It does NOT include all information about the possible uses, directions, warnings, precautions, interactions, adverse effects, or risks that may apply to this medicine. This information is not specific medical advice and does not replace information you receive from your health care provider. You must talk with your healthcare provider for complete information about the risks and benefits of using this medicine.

Do not flush unused medications or pour down a sink or drain.

 Information Expires 04/18/2013



4917 Harry Hines Blvd. Dallas, TX 75235
(214) 590-6120

Patient: ABNER, CLARENCE FRANKLIN
6827 LATTA PKWY

DALLAS, TX 75227

MRN 4258376



AB

Phone: (214) 799-1296

DOB: 7/10/1962

- i. Please use this form to reorder your prescriptions!
- ii. Please verify that the above address is correct!

*****MAIL ORDER*** ***MAIL ORDER*** ***MAIL ORDER*****

Counsel	Rx Number	Medication	Qty	Refills Remain	Days	Date Filled	Health Plan	Amt Due	Reorde
<input type="checkbox"/> Mail	072754848/00	CYCLOBENZAPRINE 10 MG TAB Mfr: MYLAN NDC: 00378-0751-10	30.00	0	10	02/07/2013	105701 Parkland Health Plus Level 2	\$3.00	<input type="checkbox"/>
		Take 1 tablet by mouth 3 times a day as needed for muscle spasms					Dr Mir, Tasaduq H Proxy: Candelario, David W DAW: 0		<input type="checkbox"/>

Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-332-1088.
Do Not Flush unused medications or pour down a sink or drain.

A pharmacist is available during normal business hours to cover questions concerning your prescriptions.
Un farmaceutico esta disponible durante las horas regulares para contestar preguntas relacionadas a su receta.

Save for Insurance & Tax Records. This is Not a Cash Receipt.

Read this medicine information sheet carefully each time you get this medicine filled.

Cyclobenzaprine Tablets

Pronunciation: SYE-kloe-BEN-za-preen

Brand Name: Flexeril

This medicine is used for:

Treating muscle spasms caused by painful muscle conditions. It should be used along with rest and physical therapy. It may also be used for other conditions as determined by your doctor.

This medicine is a muscle relaxant. It works in parts of the brain and nervous system to help reduce muscle spasms.

Do NOT use this medicine if:

- you are allergic to any ingredient in this medicine
- you have an overactive thyroid, moderate to severe liver problems, or certain heart problems (eg, irregular heartbeat, congestive heart failure, heart block, conduction problems), or if you have recently had a heart attack
- you are taking or have taken a monoamine oxidase inhibitor (MAOI) (eg, phenelzine, rasagiline) in the last 14 days

Contact your doctor or health care provider right away if any of these apply to you.

Before using this medicine:

Some medical conditions may interact with this medicine. Tell your doctor or pharmacist if you have any medical conditions, especially if any of the following apply to you:

- if you are pregnant, planning to become pregnant, or are breast-feeding
- if you are taking any prescription or nonprescription medicine, herbal preparation, or dietary supplement
- if you have allergies to medicines, foods, or other substances
- if you have a history of liver problems, heart problems, cerebral palsy, brain or spinal cord disease, or stroke
- if you have a history of glaucoma, increased pressure in the eye, or trouble urinating

Some MEDICINES MAY INTERACT with this medicine. Tell your health care provider if you are taking any other medicines, especially any of the following:

- MAOIs (eg, phenelzine, rasagiline) because serious, sometimes fatal reactions, including high fever and severe seizures, may occur
- Tramadol because the risk of seizures may be increased
- Droperidol or fluoxetine because severe heart problems, including irregular heartbeat, may occur
- Anticholinergics (eg, methscopolamine, benztropine), barbiturates (eg, phenobarbital), cimetidine, fluconazole, fluvoxamine, mibefradil, naproxen, or phenothiazines (eg, chlorpromazine) because they may increase the risk of this medicine's side effects
- Carbamazepine because it may decrease this medicine's effectiveness or increase the risk of this medicine's side effects
- Guanethidine or guanfacine because their effectiveness may be decreased by this medicine
- Sympathomimetics (eg, albuterol, epinephrine, phenylephrine) because their effectiveness may be decreased or the risk of their side effects may be increased by this medicine

This may not be a complete list of all interactions that may occur. Ask your health care provider if this medicine may interact with other medicines that you take. Check with your health care provider before you start, stop, or change the dose of any medicine.

How to use this medicine:

Use this medicine as directed by your doctor. Check the label on the medicine for exact dosing instructions.

- Take this medicine by mouth with or without food. If stomach upset occurs, take with food to reduce stomach irritation.
- Do not suddenly stop taking this medicine without checking with your doctor.
- If you miss a dose of this medicine, take it as soon as possible. If it is almost time for your next dose, skip the missed dose. Go back to your regular dosing schedule. Do not take 2 doses at once.

Ask your health care provider any questions you may have about how to use this medicine.

Important safety information about this medicine:

- This medicine may cause drowsiness, dizziness, or blurred vision. These effects may be worse if you take it with alcohol or certain medicines. Use this medicine with caution. Do not drive or perform other possibly unsafe tasks until you know how you react to it.
- Do not drink alcohol or use medicines that may cause drowsiness (eg, sleep aids, muscle relaxers) while you are using this medicine: it may add to their effects. Ask your pharmacist if you have questions about which medicines may cause drowsiness.
- Do NOT take more than the recommended dose or use for longer than 2 to 3 weeks without checking with your doctor.
- If your symptoms (eg, pain, tenderness, decreased range of motion) do not get better within 2 to 3 weeks or if they get worse, contact your doctor.
- Do not become overheated in hot weather or while you are being active; heatstroke may occur.

- If you experience dry mouth, use sugarless candy or gum, or melt bits of ice in your mouth. If dry mouth continues for more than 2 weeks, contact your dentist or doctor.
- Use this medicine with extreme caution in the ELDERLY; they may be more sensitive to its effects, especially confusion, hallucinations, and fast or irregular heartbeat.
- This medicine should be used with extreme caution in CHILDREN younger than 15 years old; safety and effectiveness in these children have not been confirmed.
- PREGNANCY and BREAST-FEEDING: If you become pregnant, contact your doctor. You will need to discuss the benefits and risks of using this medicine while you are pregnant. It is not known if this medicine is found in breast milk. If you are or will be breast-feeding while you use this medicine, check with your doctor. Discuss any possible risks to your baby.

Possible side effects of this medicine:

All medicines may cause side effects, but many people have no, or minor, side effects. Check with your doctor if any of these most COMMON side effects persist or become bothersome:

Constipation; diarrhea; dizziness; drowsiness; dry mouth; fatigue; nausea; nervousness; stomach pain or upset.

Seek medical attention right away if any of these SEVERE side effects occur:

Severe allergic reactions (rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, or tongue); chest pain; confusion; fainting; fast or irregular heartbeat; mental or mood changes; numbness of an arm or a leg; one-sided weakness; seizures; sudden severe stomach pain; severe dizziness or vomiting; speech or vision problems; trouble urinating; yellowing of the skin or eyes.

This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your health care provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects at <http://www.fda.gov/medwatch>.

If OVERDOSE is suspected:

Contact 1-800-222-1222 (the American Association of Poison Control Centers), your local poison control center (<http://www.aapcc.org>), or emergency room immediately. Symptoms may include agitation; chest pain; coma; confusion; fast or irregular heartbeat; loss of coordination; hallucinations; seizures; severe drowsiness, dizziness, or headache; slurred speech; tremor; vomiting.

Proper storage of this medicine:

Store this medicine at room temperature, between 68 and 77 degrees F (20 and 25 degrees C). Store away from heat, moisture, and light. Do not store in the bathroom. Keep this medicine out of the reach of children and away from pets.

General information:

- If you have any questions about this medicine, please talk with your doctor, pharmacist, or other health care provider.
- This medicine is to be used only by the patient for whom it is prescribed. Do not share it with other people.
- If your symptoms do not improve or if they become worse, check with your doctor.
- Check with your pharmacist about how to dispose of unused medicine.

This information is a summary only. It does not contain all information about this medicine. If you have questions about the medicine you are taking or would like more information, check with your doctor, pharmacist, or other health care provider.

Issue Date: April 6, 2011
 Database Edition 11.2.1.001
 Copyright © 2011 Wolters Kluwer Health, Inc.

**Medication
List**Parkland Health & Hospital System 5201 Harry Hines Blvd Dallas, Texas 75235 214-
590-8000**Discharge Medication Reconciliation Report**Patient Name: **Abner, Clarence Franklin**
Allergies: **Hydrocodone**MRN: **4258376**Admission Date: **Patient not admitted****Important Information!**

This is a list of your medications based on available information. Take these medications as directed. If you are taking medications not on this list and have questions about them, ask your doctor or call your pharmacist for more information.

Please bring this list and your medications to every visit or appointment.

Medications to Take At Home

Medication Name	Dose Instructions	Comments/Reason for Prescription
albuterol 90 mcg/actuation HFA inhaler	Inhale 2 Puffs by mouth every 4 hours as needed for shortness of breath.	> >
ASCORBATE CALCIUM (VITAMIN C ORAL)	Take by mouth.	> >
cyclobenzaprine 10 mg tablet	Take 1 Tab by mouth 3 times a day as needed for muscle spasms.	> >
DOCOSAHEXANOIC ACID/EPA (FISH OIL ORAL)	Take by mouth.	> >
etodolac 400 mg tablet	Take 1 Tab by mouth twice a day as needed for pain.	> >
fluticasone 50 mcg/actuation nasal spray	Use 2 Sprays in each nostril one time a day.	> >
MULTIVITAMINS, THERAPEUTIC (THERAPEUTIC MULTIVITAMIN ORAL)	Take by mouth.	> >
pantoprazole EC 40 mg tablet	Take 1 Tab by mouth one time a day.	> >

Medication List Comments:

10/6/11: Patient also taking a pain medication but can't recall the name of the medication. R.Daniels, RN

Discharge medications last reviewed by:User
CANDELARIO, DAVID WAYNE, DO [DWCAND]Date and Time
1/31/2013 12:03 PM



Parkland

Parkland Health & Hospital System
(214) 590-4900

Receipt

Your Reference Number:

2013057207-10

02/26/2013 12:19 PM

TRANSACTIONS

Patient Payment 2013057207-10-1	\$10.00
------------------------------------	---------

MRN: 4000029

MEMBER NAME: RETAILPHARMACY,PHP-INDIGENT

ACCOUNT NBR: 1

Hospital Account Payment	\$10.00
--------------------------	---------

TOTAL AMOUNT: \$10.00


PAYMENT

Master Card Credit Sale	\$10.00
-------------------------	---------

4917 Harry Hines Blvd. Dallas, TX 75235
(214) 590-6120

Patient: **ABNER, CLARENCE FRANKLIN**
6827 LATTA PKWY

MRN 4258376



AB

DALLAS, TX 75227

Phone: (214) 799-1296

DOB: 7/10/1962

- i. Please use this form to reorder your prescriptions!**
- ii. Please verify that the above address is correct!**

Counsel	Rx Number	Medication	Qty	Refills Remain	Days	Date Filled	Health Plan	Amt Due	Reorde
<input type="checkbox"/>	072761480/00	CYCLOBENZAPRINE 10 MG TAB Mfr: MYLAN NDC: 00378-0751-10	30.00	0	10	02/14/2013	105701 Parkland Health Plus Level 2	\$5.00	<input type="checkbox"/>
		Take 1 tablet by mouth 3 times a day as needed for muscle spasms					Dr Osayande, Amimi S Proxy: Nguyen, Jessica N DAW: 0		

Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-332-1088.
Do Not Flush unused medications or pour down a sink or drain.

A pharmacist is available during normal business hours to cover questions concerning your prescriptions.
Un farmaceutico esta disponible durante las horas regulares para contestar preguntas relacionadas a su receta.

Save for Insurance & Tax Records. This is Not a Cash Receipt.

PARKLAND MAIL ORDER PHARMACY

4917 Harry Hines Blvd.
 Dallas, Texas 75235-7718
 Phone: (214) 590-1400
 Fax: (214) 590-2879
 Hours 8:00AM - 4:00PM
 Monday - Friday

PLEASE COMPLETE ALL AREAS POR FAVOR COMPLETE TODAS LAS AREAS		
NAME / NOMBRE		M R N
ADDRESS / DIRECCION		APT #
CITY / CIUDAD	STATE / ESTADO	ZIP / CODIGO POSTAL
EMPLOYEE ID #	PHONE / TELEFONO	

PRESCRIPTION NUMBER NUMERO DE RECETA	DRUG MEDICAMENTO	PRICE PRECIO
1		
2		
3		
4		
5		
6		
7		
8		
9		
10		
11		
12		
13		
TOTAL		

ENCLOSE YOUR CHECK, MONEY ORDER OR CREDIT CARD INFORMATION AS INDICATED BELOW. **(Please DO NOT send cash.)**
 INCLUYA SU CHEQUE, GIRO POSTAL O LA INFORMACION DE SU TARJETA DE CREDITO COMO SE LE INDICA ABAJO. **(Por favor no envíe dinero en efectivo)**
 Check Money Order VISA (13 or 16 Digits) American Express (15 Digits) MasterCard (16 Digits) Discover Card Parkland Payroll Deduct

CARD
ACCOUNT
NUMBER

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	

EXPIRATION
DATE of
CARD

		-		
--	--	---	--	--

Signature of Authorized Buyer: X

Firma de la persona autorizada

Parkland Prescription Center

02/26/2013 11:37 AM

4917 Harry Hines Blvd. Dallas, TX 75235
(214) 590-6120

Patient: ABNER, CLARENCE FRANKLIN
6827 LATTA PKWY

DALLAS, TX 75227

MRN 4258376



AB

Phone: (214) 799-1296

DOB: 7/10/1962

- i. Please use this form to reorder your prescriptions!
- ii. Please verify that the above address is correct!

Counsel	Rx Number	Medication	Qty	Refills Remain	Days	Date Filled	Health Plan	Amt Due	Reorde
<input type="checkbox"/>	072771761/00	DOXAZOSIN 1 MG TAB Mfr: TEVA PHARMACEUTICALS US NDC: 00093-8120-01 Take 1 tablet by mouth daily at bedtime	30.00	3	30	02/26/2013	105701 Parkland Health Plus Level 2	\$5.00	<input type="checkbox"/>
<input type="checkbox"/>	072771762/00	CYCLOBENZAPRINE 10 MG TAB Mfr: MYLAN NDC: 00378-0751-10 Take 1 tablet by mouth 3 times a day as needed for muscle spasms	60.00	0	20	02/26/2013	105701 Parkland Health Plus Level 2	\$5.00	<input type="checkbox"/>

Dr. Osayanda, Animi S
Proxy: Newton, Haley C
DAW: 0

Dr. Osayanda, Animi S
Proxy: Newton, Haley C
DAW: 0

Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-332-1088.
Do Not Flush unused medications or pour down a sink or drain.

A pharmacist is available during normal business hours to cover questions concerning your prescriptions.
Un farmaceutico esta disponible durante las horas regulares para contestar preguntas relacionadas a su receta.

Save for Insurance & Tax Records. This is Not a Cash Receipt.