

Call your doctor for medical advice about side effects.  
You may report side effects to FDA at 1-800-FDA-1088.

**INGREDIENT NAME:** IBUPROFEN (eye-byoo-PROE-fen)

**COMMON USES:** This medicine is a nonsteroidal anti-inflammatory drug (NSAID) used to treat mild to moderate pain, osteoarthritis, and rheumatoid arthritis. It may also be used to treat other conditions as determined by your doctor.

**BEFORE USING THIS MEDICINE:** **WARNING: THE RISK OF SERIOUS AND SOMETIMES FATAL HEART PROBLEMS, HEART ATTACK, AND STROKE** may be increased with the use of this medicine. This risk may be increased the longer you use this medicine. Risk may also be higher in patients who have heart problems or who are at risk for heart problems. **THIS MEDICINE SHOULD NOT BE USED** to treat pain before or after coronary artery heart bypass (CABG) surgery. **THE RISK OF SERIOUS AND SOMETIMES FATAL STOMACH AND BOWEL PROBLEMS**, including bleeding, ulcers, and holes in the stomach and bowel, is increased while using this medicine. These problems may occur at any time during therapy, with or without symptoms. Elderly patients are at higher risk for serious stomach problems. Ask your doctor or pharmacist for more information about this medicine and its side effects. Some medicines or medical conditions may interact with this medicine. **INFORM YOUR DOCTOR OR PHARMACIST** of all prescription and over-the-counter medicine that you are taking. **DO NOT TAKE THIS MEDICINE** if you are also taking heparins or tacrolimus. **ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION** may be needed if you are taking serotonin reuptake blocker medicines such as fluoxetine or citalopram, "blood thinners" such as warfarin, bisphosphonates such as alendronate or risedronate, cyclosporine, corticosteroids such as prednisone, high blood pressure medicines (including ACE inhibitors such as captopril, angiotensin II receptor antagonists such as losartan, and beta-blockers such as metoprolol), "water pills" (diuretics such as furosemide, hydrochlorothiazide, triamterene), lithium, methotrexate, or aspirin. **DO NOT START OR STOP ANY MEDICINE** without doctor or pharmacist approval. Inform your doctor of any other medical conditions including poorly controlled diabetes, dehydration, heart problems (such as heart failure or history of heart attack), swelling of the hands, feet, or ankles (edema), high blood pressure, history of stroke, blood clotting problems, stomach or bowel problems (such as bleeding or ulcers), history of tobacco use or alcohol use, kidney problems, liver problems, blood or bleeding problems (such as anemia), asthma, growths in the nose (nasal polyps), any allergies (especially history of angioedema with symptoms of lip, tongue, throat swelling), pregnancy, or breast-feeding. **USE OF THIS MEDICINE IS NOT RECOMMENDED** if you have a history of allergy to aspirin or other NSAIDs (e.g., naproxen, celecoxib). **USE OF THIS MEDICINE IS NOT RECOMMENDED** if you have history of severe kidney disease or if you are going to have or have recently had coronary artery heart bypass (CABG) surgery. Contact your doctor or pharmacist if you have any questions or concerns about taking this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for using this medicine provided by your doctor. This medicine may come with a medication guide. Read it carefully. Ask your doctor, nurse, or pharmacist any questions that you may have about this medicine. **TAKE THIS MEDICINE** with a full glass (8 oz./240 ml) of water. **DO NOT** lie down for 30 minutes after taking this medicine. The dosage is based on your medical condition and response to therapy. If repeat doses are needed, they are usually given 6 or 8 hours apart, or as directed by your doctor. When used in children, the dose is based on your child's weight. Read the product instructions to find the appropriate dose for your child's weight. Consult the pharmacist or doctor if you have questions or if you need help in choosing the appropriate dosage form. **THIS MEDICINE MAY BE TAKEN WITH FOOD** if it upsets your stomach. Taking it with food may not decrease the risk of stomach or bowel problems (such as bleeding or ulcers) that may occur while taking this medicine. Talk with your doctor or pharmacist if you experience persistent stomach upset. **STORE THIS MEDICINE** at room temperature, away from heat and light. **Do not store in the bathroom.** **IF YOU MISS A DOSE OF THIS MEDICINE**, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. **Do not take 2 doses at once.**

**CAUTIONS:** **THIS MEDICINE INCREASES YOUR RISK OF SERIOUS STOMACH OR BOWEL PROBLEMS** (such as ulcers and bleeding). This risk is increased if you are elderly or are in poor health, if you have a history of smoking or drinking alcohol, if you take corticosteroid medicines (such as prednisone) or "blood thinners" (such as warfarin), or if you take this medicine for a long period of time. **THIS MEDICINE MAY ALSO INCREASE YOUR RISK** for certain serious heart and blood vessel problems (such as heart attack and stroke). **TAKE THIS MEDICINE EXACTLY AS PRESCRIBED BY YOUR DOCTOR**, at the lowest possible dose for the

shortest time needed. Talk with your doctor or pharmacist for further information. **DO NOT TAKE THIS MEDICINE IF YOU HAVE HAD A SEVERE ALLERGIC REACTION** to aspirin or any medicine containing aspirin or to a nonsteroidal anti-inflammatory drug (such as Feldene, Motrin, Naprosyn, Clinoril). A severe reaction includes a severe rash, hives, breathing difficulties, or dizziness. If you have a question about whether you are allergic to this medicine or if a certain medicine is a nonsteroidal anti-inflammatory drug, contact your doctor or pharmacist. **DO NOT EXCEED THE RECOMMENDED DOSE** or take this medicine for longer than 10 days for pain or 3 days for fever, unless directed by your doctor. Laboratory and/or medical tests, including blood counts, liver function tests, and kidney function tests, may be performed to monitor your progress or to check for side effects, especially if you are taking this medicine for a long period of time. **KEEP ALL DOCTOR AND LABORATORY APPOINTMENTS** while you are taking this medicine. **DO NOT DRIVE, OPERATE MACHINERY, OR DO ANYTHING ELSE THAT COULD BE DANGEROUS** until you know how you react to this medicine. **ALCOHOL WARNING:** If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take this medicine or other pain relievers/fever reducers. **BEFORE YOU BEGIN TAKING ANY NEW MEDICINE**, either prescription or over-the-counter, check with your doctor or pharmacist. If you are also taking aspirin, as prescribed by your doctor for reasons such as heart attack or stroke prevention (usually these dosages are 81-325 mg per day), continue to take the aspirin and consult your doctor or pharmacist before using this medicine. **CAUTION IS ADVISED WHEN USING THIS MEDICINE IN THE ELDERLY** because they may be more sensitive to the effects of this medicine, especially the risk of stomach or bowel effects (such as bleeding or ulcers) or kidney effects. **FOR WOMEN:** **USE OF THIS MEDICINE DURING PREGNANCY** has resulted in fetal and newborn death. If you think you may be pregnant, contact your doctor immediately. **THIS MEDICINE IS EXCRETED IN BREAST MILK.** **IF YOU ARE OR WILL BE BREAST-FEEDING** while you are using this medicine, check with your doctor or pharmacist to discuss the risks to your baby.

**POSSIBLE SIDE EFFECTS:** **SIDE EFFECTS**, that may go away during treatment, include nausea, vomiting, diarrhea, gas, constipation, indigestion, dizziness, lightheadedness, drowsiness, or headache. If they continue or are bothersome, check with your doctor. **CHECK WITH YOUR DOCTOR AS SOON AS POSSIBLE** if you experience ringing in ears. **CONTACT YOUR DOCTOR IMMEDIATELY** if you experience rapid or pounding heartbeat; easy bruising or bleeding; very stiff neck; or mental/mood changes. **CONTACT YOUR DOCTOR IMMEDIATELY** if you experience sharp or crushing chest pain; sudden shortness of breath; sudden leg pain; sudden severe headache, vomiting, dizziness, or fainting; changes in vision; numbness of an arm or leg; slurred speech; one-sided weakness; sudden unexplained weight gain; change in amount of urine produced; severe or persistent stomach pain; vomit that looks like coffee grounds; black tarry stools; itching, reddened, swollen, blistered, painful, or peeling skin; yellowing of the skin or eyes; dark urine; right-sided tenderness; severe or persistent tiredness; fever, chills, or sore throat; severe or persistent nausea; swelling of hands, ankles, feet, face, lips, eyes, throat, or tongue; difficulty swallowing or breathing; or hoarseness. **AN ALLERGIC REACTION TO THIS MEDICINE** is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash, itching, swelling, severe dizziness, or trouble breathing. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms of overdose may include severe stomach pain, coffee ground-like vomit, unusually fast or slow heartbeat, trouble breathing, extreme drowsiness, loss of consciousness, and seizures.

**ADDITIONAL INFORMATION:** **DO NOT SHARE THIS MEDICINE** with others for whom it was not prescribed. **DO NOT USE THIS MEDICINE** for other health conditions. **KEEP THIS MEDICINE** out of the reach of children and pets. **IF USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME**, obtain refills before your supply runs out.

**KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.**

Do not flush unused medications or pour down a sink or drain.

WIC# 216382

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

**RX # 0529364-09015**

DATE: 07/09/13

**IBUPROFEN 600MG TABLETS**

QTY: 60 NO REFILLS - DR. AUTH REQUIRED

Copy NDC: 53746-0465-05

Retail Price: \$16.33 Your Insurance Saved You: \$10.57

\$ 5.76

W. CHRISTENSEN, MD  
MFG: AMNEAL  
NJT/NJT/NJT /ADO

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 131906513113095999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Customer  
Receipt

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

**RX # 0529364-09015**

DATE: 07/09/13

**IBUPROFEN 600MG TABLETS**

QTY: 60 NO REFILLS - DR. AUTH REQUIRED

Copy NDC: 53746-0465-05

Retail Price: \$16.33 Your Insurance Saved You: \$10.57

\$ 5.76

W. CHRISTENSEN, MD  
MFG: AMNEAL  
NJT/NJT/NJT /ADO

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 131906513113095999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Duplicate  
Receipt

Pharmacy use only

TUE 9:35PM  
Copy

IBUPROFEN 600MG TABLETS  
53746-0465-05  
FAST RACK

QTY 60  
20 DRAM



WHITE  
FRONT: IP 465

NJT/NJT/NJT /ADO

Med Guide



**INGREDIENT NAME:** HYDROCODONE (hye-droe-KO-done) and ACETAMINOPHEN (a-seat-a-MIN-oh-fen)

**COMMON USES:** This medicine is a narcotic and analgesic combination used to relieve moderate to moderately severe pain. It may also be used to treat other conditions as determined by your doctor.

**BEFORE USING THIS MEDICINE:** **WARNING:** THIS MEDICINE CONTAINS ACETAMINOPHEN. Severe and sometimes fatal liver problems, including the need for liver transplant, have been reported with the use of acetaminophen. Most cases of these liver problems occurred in patients taking excessive doses of acetaminophen (more than 4,000 mg per day). Also, patients who developed these liver problems were often using more than 1 medicine that contained acetaminophen. Discuss any questions or concerns with your doctor. Some medicines or medical conditions may interact with this medicine. **INFORM YOUR DOCTOR OR PHARMACIST** of all prescription and over-the-counter medicine that you are taking. **DO NOT TAKE THIS MEDICINE** if you are also taking sodium oxybate (GHB). **ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION** may be needed if you are taking anticoagulants (eg, warfarin), buprenorphine, isoniazid, linezolid, monoamine oxidase inhibitors (MAOIs) (eg, phenelzine), naltrexone, other narcotic pain medicines (eg, morphine), perampanel, sleep medicines (eg, zolpidem), tricyclic antidepressants (eg, amitriptyline), or medicines that may harm the liver (eg, methotrexate, ketoconazole, certain medicines for HIV infection). Ask your doctor if you are unsure if any of your medicines might harm the liver. **DO NOT START OR STOP** any medicine without doctor or pharmacist approval. Inform your doctor of any other medical conditions, including a history of liver problems (eg, hepatitis), kidney problems, lung or breathing problems (eg, asthma, chronic obstructive pulmonary disease / COPD, drug or alcohol abuse), dependence, heart problems (eg, cor pulmonale, slow or irregular heartbeat), low blood pressure, stomach or bowel problems (eg, blockage, inflammation, paralysis), gallbladder problems, prostate problems (eg, an enlarged prostate), difficulty urinating, seizures, thyroid problems, adrenal gland problems (eg, Addison disease), mental or mood problems, inflammation of the pancreas (pancreatitis), curvature of the spine (scoliosis), suicidal thoughts or actions, allergies, pregnancy, or breast-feeding. Tell your doctor if you have severe drowsiness, or have or recently have had any head injury, increased pressure in the brain, growths in the brain (eg, tumors), or infection of the brain or nervous system. Tell your doctor if you have recently had surgery (eg, stomach or bowel surgery) or are scheduled to have surgery. Tell your doctor if you are in very poor health; are very overweight or are dehydrated; have low blood volume, stomach pain, or constipation; drink alcohol; or are going through withdrawal from alcohol or other substances. **USE OF THIS MEDICINE IS NOT RECOMMENDED** if you have very slow or difficult breathing, or you are having an asthma attack. Contact your doctor or pharmacist if you have any questions or concerns about taking this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for using this medicine provided by your doctor. TAKE THIS MEDICINE by mouth with or without food. If stomach upset occurs, take with food to reduce stomach irritation. IF YOU HAVE BEEN TAKING THIS MEDICINE REGULARLY or for longer than a few weeks, do not suddenly stop taking it without checking with your doctor. Your doctor may need to gradually lower your dose. **SOME OF THESE PRODUCTS CONTAIN SULFITES.** Sulfites may cause an allergic reaction in some patients (eg, asthma patients). If you have ever had an allergic reaction to sulfites, ask your pharmacist if your product has sulfites in it. **STORE THIS MEDICINE** at room temperature, between 68 and 77 degrees F (20 and 25 degrees C), away from heat, moisture, and light. Do not store in the bathroom. **KEEP THIS MEDICINE** out of the reach of children and away from pets. IF YOU MISS A DOSE OF THIS MEDICINE and you are taking it regularly, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.

**CAUTIONS:** **DO NOT TAKE THIS MEDICINE** if you are allergic to any ingredient in this medicine. **BEFORE USING THIS MEDICINE,** check with your doctor if you have had an allergic reaction to other narcotic medicines (eg, codeine, morphine, oxycodone). **DO NOT EXCEED THE RECOMMENDED DOSE,** take more often than prescribed, or use for longer than prescribed without checking with your doctor. Exceeding the recommended dose or taking this medicine for longer than prescribed may be habit-forming. If using this medicine for an extended period of time, **DO NOT SUDDENLY STOP** taking this medicine without your doctor's approval. When using for an extended period, this medicine may not work as well and may require different dosing. Talk with your doctor if this medicine stops working well. **THIS MEDICINE MAY AFFECT CERTAIN LAB TEST RESULTS.** Make sure that your doctor and laboratory personnel know you are taking this medicine. **BEFORE YOU HAVE ANY MEDICAL OR DENTAL TREATMENTS,** emergency care, or surgery, tell the doctor or dentist that you are using this medicine. **THIS MEDICINE MAY CAUSE DIZZINESS,** lightheadedness, or fainting. Alcohol, hot weather, exercise, and fever can increase these effects. To prevent them, sit up or stand slowly, especially in the morning. Also, sit or lie down at the first sign of dizziness, lightheadedness, or weakness. **THIS MEDICINE MAY ALSO CAUSE DROWSINESS OR BLURRED VISION.** Do not drive, operate machinery, or do anything else that could be dangerous until you know how you react to this medicine. Using this medicine alone, or with other medicines, or with alcohol may lessen your ability to drive or to perform other potentially dangerous tasks. **DO NOT DRINK ALCOHOL** while you are using this medicine. **CHECK WITH YOUR DOCTOR** before you use medicines that may cause drowsiness (eg, sleep aids, muscle relaxers) while you are using this medicine; it may add to their effects. Ask your pharmacist if you have questions about which medicines may cause drowsiness. **THIS MEDICINE CONTAINS ACETAMINOPHEN.** Do not take additional acetaminophen for pain or fever without checking with your doctor or pharmacist. Ask your pharmacist if you have questions about which medicines contain acetaminophen. **THIS MEDICINE MAY HARM YOUR LIVER.** Your risk may be greater if you drink alcohol while you are using this medicine. Talk to your doctor before you take this medicine or other fever reducers if you drink alcohol. **CONTACT YOUR DOCTOR RIGHT AWAY** if you take more than 4,000 mg of acetaminophen per day, even if you feel well. **BEFORE YOU BEGIN TAKING ANY NEW MEDICINE,** either prescription or over-the-counter, check with your doctor or pharmacist. **CAUTION IS ADVISED WHEN USING THIS MEDICINE IN THE ELDERLY** because they may be more sensitive to the effects of the medicine. **FOR WOMEN:** IF YOU BECOME PREGNANT, discuss with your doctor the benefits and risks of using this medicine during pregnancy. Long-term use of this medicine during pregnancy may cause dependence in the fetus or newborn. Discuss any questions or concerns with your doctor. **THIS MEDICINE IS EXCRETED** in breast milk. **DO NOT BREAST-FEED** while taking this medicine.

**POSSIBLE SIDE EFFECTS:** SIDE EFFECTS that may occur while taking this medicine include dizziness, drowsiness, lightheadedness, nausea, or vomiting. If

they continue or are bothersome, check with your doctor. **THIS MEDICINE MAY CAUSE CONSTIPATION.** Talk with your doctor or pharmacist about using a stool softener or laxative to prevent constipation. It is also important to maintain a diet adequate in fiber, drink plenty of water, and exercise to prevent constipation. If you become constipated while taking this medicine, talk with your doctor or pharmacist. **CONTACT YOUR DOCTOR IMMEDIATELY** if you experience a change in the amount of urine produced; change or loss in hearing; chest pain; confusion; fainting; fever, chills, or persistent sore throat; mental or mood changes (eg, agitation, anxiety, depression); red, swollen, blistered, or peeling skin; severe or persistent constipation; severe or persistent dizziness, drowsiness, headache, or lightheadedness; shortness of breath; slow, shallow, or difficult breathing; symptoms of liver problems (eg, yellowing of the skin or eyes, pale stools, dark urine, persistent loss of appetite, unusual or severe stomach pain); trouble urinating; unusual bruising or bleeding; unusual tiredness or weakness; or vision changes. **AN ALLERGIC REACTION** to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, throat, or tongue; or unusual hoarseness. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** IF OVERDOSE IS SUSPECTED, contact your local poison control center or emergency room immediately. Symptoms of overdose may include bluish skin or nails; cold and clammy skin; coma; decreased urination; difficult, shallow, or slow breathing; fainting; limp muscles; loss of consciousness; nausea or vomiting; pinpoint pupils; seizures; severe dizziness, drowsiness, or light-headedness; slow heartbeat; symptoms of liver problems (eg, yellowing of the skin or eyes, pale stools, dark urine, loss of appetite, unusual or severe stomach pain); unusual bruising or bleeding; or unusual sweating or tiredness.

**ADDITIONAL INFORMATION:** If your symptoms do not improve or if they become worse, contact your doctor. **DO NOT SHARE THIS MEDICINE** with others for whom it was not prescribed. **DO NOT USE THIS MEDICINE** for other health conditions. **CHECK WITH YOUR PHARMACIST** about how to dispose of unused medicine.

**KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.**

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

**RX # 0517526-09015**

DATE: 05/08/13

**HYDROCODONE /ACETAMINOPHEN 5-325 TB**

QTY: 90 1 REFILL BEFORE 11/06/13

New NDC: 00591-3202-01

Retail Price: \$65.89 Your Insurance Saved You: \$55.89

**\$ 10.00**

A. PARK, MD  
MFG: WATSON  
JAC/ADO/ADO / NJT

PLAN: WPB  
GROUP#: 01960917  
CLAIM REF#: 131284531485091999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Customer Receipt

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

**RX # 0517526-09015**

DATE: 05/08/13

**HYDROCODONE /ACETAMINOPHEN 5-325 TB**

QTY: 90 1 REFILL BEFORE 11/06/13

New NDC: 00591-3202-01

Retail Price: \$65.89 Your Insurance Saved You: \$55.89

**\$ 10.00**

A. PARK, MD  
MFG: WATSON  
JAC/ADO/ADO / NJT

PLAN: WPB  
GROUP#: 01960917  
CLAIM REF#: 131284531485091999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Duplicate Receipt

Pharmacy use only

WED 4:03PM


New

HYDROCODONE /ACETAMINOPHEN 5-325 TB

00591-3202-01

FAST RACK

QTY 90  
20 DRAM



WHITE AND MULTICOLOR  
FRONT: 3202  
BACK: WATSON

JAC/ADO/ADO / NJT

Call your doctor for medical advice about side effects.  
You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 216382



Call your doctor for medical advice about side effects.  
You may report side effects to FDA at 1-800-FDA-1088.

**INGREDIENT NAME:** IBUPROFEN (eye-byoo-PROE-fen)

**COMMON USES:** This medicine is a nonsteroidal anti-inflammatory drug (NSAID) used to treat mild to moderate pain, osteoarthritis, and rheumatoid arthritis. It may also be used to treat other conditions as determined by your doctor.

**BEFORE USING THIS MEDICINE:** WARNING: THE RISK OF SERIOUS AND SOMETIMES FATAL HEART PROBLEMS, HEART ATTACK, AND STROKE may be increased with the use of this medicine. This risk may be increased the longer you use this medicine. Risk may also be higher in patients who have heart problems or who are at risk for heart problems. THIS MEDICINE SHOULD NOT BE USED to treat pain before or after coronary artery heart bypass (CABG) surgery. THE RISK OF SERIOUS AND SOMETIMES FATAL STOMACH AND BOWEL PROBLEMS, including bleeding, ulcers, and holes in the stomach and bowel, is increased while using this medicine. These problems may occur at any time during therapy, with or without symptoms. Elderly patients are at higher risk for serious stomach problems. Ask your doctor or pharmacist for more information about this medicine and its side effects. Some medicines or medical conditions may interact with this medicine. INFORM YOUR DOCTOR OR PHARMACIST of all prescription and over-the-counter medicine that you are taking. DO NOT TAKE THIS MEDICINE if you are also taking heparin or tacrolimus. ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION may be needed if you are taking serotonin reuptake blocker medicines such as fluoxetine or citalopram, "blood thinners" such as warfarin, bisphosphonates such as alendronate or risedronate, cyclosporine, corticosteroids such as prednisone, high blood pressure medicines (including ACE inhibitors such as captopril, angiotensin II receptor antagonists such as losartan, and beta-blockers such as metoprolol), "water pills" (diuretics such as furosemide, hydrochlorothiazide, triamterene), lithium, methotrexate, or aspirin. DO NOT START OR STOP ANY MEDICINE without doctor or pharmacist approval. Inform your doctor of any other medical conditions including poorly controlled diabetes, dehydration, heart problems (such as heart failure or history of heart attack), swelling of the hands, feet, or ankles (edema), high blood pressure, history of stroke, blood clotting problems, stomach or bowel problems (such as bleeding or ulcers), history of tobacco use or alcohol use, kidney problems, liver problems, blood or bleeding problems (such as anemia), asthma, growths in the nose (nasal polyps), any allergies (especially history of angioedema with symptoms of lip, tongue, throat swelling), pregnancy, or breast-feeding. USE OF THIS MEDICINE IS NOT RECOMMENDED if you have a history of allergy to aspirin or other NSAIDs (e.g., naproxen, celecoxib). USE OF THIS MEDICINE IS NOT RECOMMENDED if you have history of severe kidney disease or if you are going to have or have recently had coronary artery heart bypass (CABG) surgery. Contact your doctor or pharmacist if you have any questions or concerns about taking this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for using this medicine provided by your doctor. This medicine may come with a medication guide. Read it carefully. Ask your doctor, nurse, or pharmacist any questions that you may have about this medicine. TAKE THIS MEDICINE with a full glass (8 oz./240 ml) of water. DO NOT lie down for 30 minutes after taking this medicine. The dosage is based on your medical condition and response to therapy. If repeat doses are needed, they are usually given 6 or 8 hours apart, or as directed by your doctor. When used in children, the dose is based on your child's weight. Read the product instructions to find the appropriate dose for your child's weight. Consult the pharmacist or doctor if you have questions or if you need help in choosing the appropriate dosage form. THIS MEDICINE MAY BE TAKEN WITH FOOD if it upsets your stomach. Taking it with food may not decrease the risk of stomach or bowel problems (such as bleeding or ulcers) that may occur while taking this medicine. Talk with your doctor or pharmacist if you experience persistent stomach upset. STORE THIS MEDICINE at room temperature, away from heat and light. Do not store in the bathroom. IF YOU MISS A DOSE OF THIS MEDICINE, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.

**CAUTIONS:** THIS MEDICINE INCREASES YOUR RISK OF SERIOUS STOMACH OR BOWEL PROBLEMS (such as ulcers and bleeding). This risk is increased if you are elderly or are in poor health, if you have a history of smoking or drinking alcohol, if you take corticosteroid medicines (such as prednisone) or "blood thinners" (such as warfarin), or if you take this medicine for a long period of time. THIS MEDICINE MAY ALSO INCREASE YOUR RISK for certain serious heart and blood vessel problems (such as heart attack and stroke). TAKE THIS MEDICINE EXACTLY AS PRESCRIBED BY YOUR DOCTOR, at the lowest possible dose for the

shortest time needed. Talk with your doctor or pharmacist for further information. DO NOT TAKE THIS MEDICINE IF YOU HAVE HAD A SEVERE ALLERGIC REACTION to aspirin or any medicine containing aspirin or to a nonsteroidal anti-inflammatory drug (such as Feldene, Motrin, Naprosyn, Clinoril). A severe reaction includes a severe rash, hives, breathing difficulties, or dizziness. If you have a question about whether you are allergic to this medicine or if a certain medicine is a nonsteroidal anti-inflammatory drug, contact your doctor or pharmacist. DO NOT EXCEED THE RECOMMENDED DOSE or take this medicine for longer than 10 days for pain or 3 days for fever, unless directed by your doctor. Laboratory and/or medical tests, including blood counts, liver function tests, and kidney function tests, may be performed to monitor your progress or to check for side effects, especially if you are taking this medicine for a long period of time. KEEP ALL DOCTOR AND LABORATORY APPOINTMENTS while you are taking this medicine. DO NOT DRIVE, OPERATE MACHINERY, OR DO ANYTHING ELSE THAT COULD BE DANGEROUS until you know how you react to this medicine. ALCOHOL WARNING: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take this medicine or other pain relievers/fever reducers. BEFORE YOU BEGIN TAKING ANY NEW MEDICINE, either prescription or over-the-counter, check with your doctor or pharmacist. If you are also taking aspirin, as prescribed by your doctor for reasons such as heart attack or stroke prevention (usually these dosages are 81-325 mg per day), continue to take the aspirin and consult your doctor or pharmacist before using this medicine. CAUTION IS ADVISED WHEN USING THIS MEDICINE IN THE ELDERLY because they may be more sensitive to the effects of this medicine, especially the risk of stomach or bowel effects (such as bleeding or ulcers), or kidney effects. FOR WOMEN: USE OF THIS MEDICINE DURING PREGNANCY has resulted in fetal and newborn death. If you think you may be pregnant, contact your doctor immediately. THIS MEDICINE IS EXCRETED IN BREAST MILK. IF YOU ARE OR WILL BE BREAST-FEEDING while you are using this medicine, check with your doctor or pharmacist to discuss the risks to your baby.

**POSSIBLE SIDE EFFECTS:** SIDE EFFECTS, that may go away during treatment, include nausea, vomiting, diarrhea, gas, constipation, indigestion, dizziness, lightheadedness, drowsiness, or headache. If they continue or are bothersome, check with your doctor. CHECK WITH YOUR DOCTOR AS SOON AS POSSIBLE if you experience ringing in ears. CONTACT YOUR DOCTOR IMMEDIATELY if you experience rapid or pounding heartbeat; easy bruising or bleeding; very stiff neck; or mental/mood changes. CONTACT YOUR DOCTOR IMMEDIATELY if you experience sharp or crushing chest pain; sudden shortness of breath; sudden leg pain; sudden severe headache, vomiting, dizziness, or fainting; changes in vision; numbness of an arm or leg; slurred speech; one-sided weakness; sudden unexplained weight gain; change in amount of urine produced; severe or persistent stomach pain; vomit that looks like coffee grounds; black tarry stools; itching, reddened, swollen, blistered, painful, or peeling skin; yellowing of the skin or eyes; dark urine; right-sided tenderness; severe or persistent tiredness; fever, chills, or sore throat; severe or persistent nausea; swelling of hands, ankles, feet, face, lips, eyes, throat, or tongue; difficulty swallowing or breathing; or hoarseness. AN ALLERGIC REACTION TO THIS MEDICINE is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash, itching, swelling, severe dizziness, or trouble breathing. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms of overdose may include severe stomach pain, coffee ground-like vomit, unusually fast or slow heartbeat, trouble breathing, extreme drowsiness, loss of consciousness, and seizures.

**ADDITIONAL INFORMATION:** DO NOT SHARE THIS MEDICINE with others for whom it was not prescribed. DO NOT USE THIS MEDICINE for other health conditions. KEEP THIS MEDICINE out of the reach of children and pets. IF USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME, obtain refills before your supply runs out.

**KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.**

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

**RX # 0515796-09015**

DATE: 05/26/13

**IBUPROFEN 600MG TABLETS**

QTY: 90 NO REFILLS - DR. AUTH REQUIRED

Refill NDC: 53746-0465-05

Retail Price: \$21.50 Your Insurance Saved You: \$13.60

**\$ 7.90**

W. CHRISTENSEN, MD  
MFG: AMNEAL  
XXX/ADO/ADO/ ADO

PLAN: WPB  
GROUP#: 01960917  
CLAIM REF# 131461234342014999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Customer  
Receipt

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

**RX # 0515796-09015**

DATE: 05/26/13

**IBUPROFEN 600MG TABLETS**

QTY: 90 NO REFILLS - DR. AUTH REQUIRED

Refill NDC: 53746-0465-05

Retail Price: \$21.50 Your Insurance Saved You: \$13.60

**\$ 7.90**

W. CHRISTENSEN, MD  
MFG: AMNEAL  
XXX/ADO/ADO/ ADO

PLAN: WPB  
GROUP#: 01960917  
CLAIM REF# 131461234342014999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Duplicate  
Receipt

Pharmacy use only

SUN 12:00PM

Refill


IBUPROFEN 600MG TABLETS

53746-0465-05

FAST RACK

QTY 90

30 DRAM



WHITE  
FRONT: IP 465

XXX/ADO/ADO/ ADO

Med Guide

Do not flush unused medications or pour down a sink or drain.

WIC# 216382



Call your doctor for medical advice about side effects.  
You may report side effects to FDA at 1-800-FDA-1088.

**INGREDIENT NAME:** TRAMADOL (TRAM-a-dol)

**COMMON USES:** This medicine is an analgesic used to treat moderate to moderately severe pain.

**BEFORE USING THIS MEDICINE:** Some medicines or medical conditions may interact with this medicine. **INFORM YOUR DOCTOR OR PHARMACIST** of all prescription and over-the-counter medicine that you are taking. **DO NOT TAKE THIS MEDICINE** if you are taking sodium oxybate (GHB), or another product that contains tramadol. **ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION** may be needed if you are taking alpha-2 receptor blockers (eg, yohimbine), anorexiant (eg, phentermine), anticoagulants (eg, warfarin), azole antifungals (eg, ketoconazole), butyrophenones (eg, haloperidol), carbamazepine, cyclobenzaprine, digoxin, furazolidone, linezolid, lithium, loxapine, macrolide antibiotics (eg, erythromycin), certain medicines for mental or mood disorders (eg, olanzapine), molidone, monoamine oxidase inhibitors (MAOIs) (eg, phenelzine, selegiline), nefazodone, opioid pain medicine (eg, codeine, hydrocodone), phenothiazines (eg, promethazine), quinine, rifampin, serotonin-norepinephrine reuptake inhibitors (SNRIs) (eg, duloxetine), selective serotonin reuptake inhibitors (SSRIs) (eg, fluoxetine, paroxetine), sleeping medicines (eg, zolpidem), St. John's wort, thioxanthenes (eg, thiothixene), tiagabine, tricyclic antidepressants (eg, amitriptyline), or "triptans" (eg, sumatriptan). **DO NOT START OR STOP** any medicine without doctor or pharmacist approval. Inform your doctor of any other medical conditions, including liver, kidney, or thyroid problems; blockage in the bladder; diabetes; heart problems (eg, cor pulmonale); high blood pressure; pancreas problems; prostate problems; metabolism problems; allergies; pregnancy; or breast-feeding. Tell your doctor if you have or recently have had any head injury, brain injury or tumor, increased pressure in the brain, or infection of the brain or nervous system. Tell your doctor if you have a history of lung or breathing problems (eg, asthma, chronic obstructive pulmonary disease / COPD); seizures (eg, epilepsy); recent stomach or bowel surgery; other stomach or bowel problems (eg, pain, inflammation, ulcers); alcohol or other substance abuse; mood or mental problems (eg, depression); or suicidal thoughts or actions. Tell your doctor if you drink alcohol or if you are going through withdrawal from alcohol or other substances. **USE OF THIS MEDICINE IS NOT RECOMMENDED** if you have severe or persistent diarrhea caused by antibiotic use; suicidal thoughts or actions; or a history of alcohol or other substance abuse or addiction. **USE OF THIS MEDICINE IS NOT RECOMMENDED** if you are intoxicated with alcohol, opioids or narcotics (eg, codeine, morphine), or sedatives or sleeping medicines (eg, temazepam, zolpidem). This medicine should not be used in **CHILDREN** younger than 16 years; safety and effectiveness in these children have not been confirmed. Contact your doctor or pharmacist if you have any questions or concerns about using this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for taking this medicine provided by your doctor. **TAKE THIS MEDICINE** by mouth with or without food. **STORE THIS MEDICINE** at 77 degrees F (25 degrees C) away from heat, moisture, and light. Do not store in the bathroom. Brief storage at temperatures between 59 and 86 degrees F (15 and 30 degrees C) is permitted. **KEEP THIS MEDICINE** out of the reach of children and away from pets. **IF YOU MISS A DOSE OF THIS MEDICINE** and you are taking it regularly, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. **DO NOT** take 2 doses at once.

**CAUTIONS:** **DO NOT TAKE THIS MEDICINE** if you have had an allergic reaction to it or if you are allergic to any ingredient in this product. **DO NOT TAKE THIS MEDICINE** if you have had a severe allergic reaction to codeine or another opioid (eg, morphine). **DO NOT EXCEED THE RECOMMENDED DOSE** or take this medicine for longer than prescribed without checking with your doctor. Exceeding the recommended dose or taking this medicine for longer than prescribed may be habit-forming or may increase the risk of seizures or breathing problems. If using this medicine for an extended period of time, **DO NOT SUDDENLY STOP** taking this medicine without your doctor's approval. Your dose may need to be slowly lowered to avoid side effects. When used for an extended period of time, this medicine may not work as well and may require different dosing. Talk with your doctor if this medicine stops working well. **DO NOT TAKE MORE THAN PRESCRIBED.** **BEFORE YOU HAVE ANY MEDICAL OR DENTAL TREATMENTS, EMERGENCY CARE, OR SURGERY,** tell the doctor or dentist that you are using this medicine. **DO NOT DRINK ALCOHOL** while you are using this medicine. **CHECK WITH YOUR DOCTOR** before you use medicines that may cause drowsiness (eg, sleep aids, muscle relaxers, narcotic pain medicines) while you are using this medicine; it may add to their effects. Ask your pharmacist if you

have questions about which medicines may cause drowsiness. This medicine may cause drowsiness or dizziness. **DO NOT DRIVE, OPERATE MACHINERY, OR DO ANYTHING ELSE THAT COULD BE DANGEROUS** until you know how you react to this medicine. Using this medicine alone, with other medicines, or with alcohol may lessen your ability to drive or to perform other potentially dangerous tasks. Alcohol, hot weather, exercise, or fever may increase dizziness. **IF DIZZINESS OCCURS,** sit up or stand slowly, especially in the morning. Sit or lie down at the first sign of this effect. **THIS MEDICINE MAY INCREASE YOUR RISK OF SEIZURES.** Your risk may be greater if you also have certain medical conditions, use certain medicines, or if you use a lot of alcohol. Talk to your doctor to see if you may have a greater risk of seizures while taking this medicine. **SEROTONIN SYNDROME** is a possibly fatal syndrome that can be caused by this medicine. Your risk may be greater if you take this medicine with certain other medicines (eg, "triptans," MAOIs, antidepressants). Symptoms may include agitation; confusion; hallucinations; coma; fever; fast or irregular heartbeat; tremor; excessive sweating; and nausea, vomiting, or diarrhea. Contact your doctor at once if you have any of these symptoms. **BEFORE YOU BEGIN TAKING ANY NEW MEDICINES,** either prescription or over-the-counter, check with your doctor or pharmacist. **CAUTION IS ADVISED WHEN USING THIS MEDICINE IN THE ELDERLY** because they may be more sensitive to the effects of this medicine. **FOR WOMEN:** This medicine may cause harm to the fetus. **IF YOU THINK YOU MAY BE PREGNANT,** discuss with your doctor the benefits and risks of using this medicine during pregnancy. **THIS MEDICINE IS EXCRETED** in breast milk. **DO NOT BREAST-FEED** while taking this medicine.

**POSSIBLE SIDE EFFECTS:** SIDE EFFECTS that may occur while taking this medicine include constipation; diarrhea; dizziness; drowsiness; dry mouth; headache; indigestion; mildly increased sweating; mild itching; nausea; trouble sleeping; vomiting; weakness. If they continue or are bothersome, check with your doctor. **CONTACT YOUR DOCTOR IMMEDIATELY** if you experience agitation; chest pain; confusion; difficult or painful urination; excessive sweating; fainting; fast or irregular heartbeat; fever; hallucinations; loss of coordination; mood or mental changes (eg, depression); red, blistered, swollen, or peeling skin; seizures; severe dizziness or lightheadedness; severe nausea, vomiting, or diarrhea; severe or persistent headache; slow or shallow breathing; suicidal thoughts or behaviors; tremor; or vision problems. **AN ALLERGIC REACTION** to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, or tongue; unusual hoarseness. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your health care provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** IF OVERDOSE IS SUSPECTED, contact your local poison control center or emergency room immediately. Symptoms may include bluish skin; cold, clammy skin; difficult, shallow, or slow breathing; drowsiness leading to unresponsiveness or coma; excessive sweating; limp muscles; pinpoint pupils; seizures; slow or irregular heartbeat.

**ADDITIONAL INFORMATION:** **DO NOT SHARE THIS MEDICINE** with others for whom it was not prescribed. **DO NOT USE THIS MEDICINE** for other health conditions. **IF YOU WILL BE USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME,** be sure to obtain necessary refills before your supply runs out. **CHECK WITH YOUR PHARMACIST** about how to dispose of unused medicine.

**KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.**

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

**RX # 0510864-09015**

DATE: 05/14/13

**TRAMADOL 50MG TABLETS**

QTY: 180 2 REFILLS BEFORE 04/07/14

Refill NDC:68382-0319-10

Retail Price: \$77.97 Your Insurance Saved You: \$67.97

**\$ 10.00**

W. CHRISTENSEN, MD  
MFG:ZYDUS  
XXX/HYK/HYK/HYK/HYK

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 131343281747063999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Customer  
Receipt

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

**RX # 0510864-09015**

DATE: 05/14/13

**TRAMADOL 50MG TABLETS**

QTY: 180 2 REFILLS BEFORE 04/07/14

Refill NDC:68382-0319-10

Retail Price: \$77.97 Your Insurance Saved You: \$67.97

**\$ 10.00**

W. CHRISTENSEN, MD  
MFG:ZYDUS  
XXX/HYK/HYK/HYK/HYK

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 131343281747063999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951


Duplicate  
Receipt

Pharmacy use only

TUE 12:06PM  
Refill

TRAMADOL 50MG TABLETS  
68382-0319-10  
FAST RACK

QTY 180  
20 DRAM



WHITE  
FRONT: 319

XXX/HYK/HYK/HYK/HYK

Do not flush unused medications or pour down a sink or drain.

WIC# 216382



Call your doctor for medical advice about side effects.  
You may report side effects to FDA at 1-800-FDA-1088.

**INGREDIENT NAME:** IBUPROFEN (eye-byoo-PROE-fen)

**COMMON USES:** This medicine is a nonsteroidal anti-inflammatory drug (NSAID) used to treat mild to moderate pain, osteoarthritis, and rheumatoid arthritis. It may also be used to treat other conditions as determined by your doctor.

**BEFORE USING THIS MEDICINE:** WARNING: THE RISK OF SERIOUS AND SOMETIMES FATAL HEART PROBLEMS, HEART ATTACK, AND STROKE may be increased with the use of this medicine. This risk may be increased the longer you use this medicine. Risk may also be higher in patients who have heart problems or who are at risk for heart problems. THIS MEDICINE SHOULD NOT BE USED to treat pain before or after coronary artery heart bypass (CABG) surgery. THE RISK OF SERIOUS AND SOMETIMES FATAL STOMACH AND BOWEL PROBLEMS, including bleeding, ulcers, and holes in the stomach and bowel, is increased while using this medicine. These problems may occur at any time during therapy, with or without symptoms. Elderly patients are at higher risk for serious stomach problems. Ask your doctor or pharmacist for more information about this medicine and its side effects. Some medicines or medical conditions may interact with this medicine. INFORM YOUR DOCTOR OR PHARMACIST of all prescription and over-the-counter medicine that you are taking. DO NOT TAKE THIS MEDICINE if you are also taking heparin or tacrolimus. ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION may be needed if you are taking serotonin reuptake blocker medicines such as fluoxetine or citalopram, "blood thinners" such as warfarin, bisphosphonates such as alendronate or risedronate, cyclosporine, corticosteroids such as prednisone, high blood pressure medicines (including ACE inhibitors such as captopril, angiotensin II receptor antagonists such as losartan, and beta-blockers such as metoprolol), "water pills" (diuretics such as furosemide, hydrochlorothiazide, triamterene), lithium, methotrexate, or aspirin. DO NOT START OR STOP ANY MEDICINE without doctor or pharmacist approval. Inform your doctor of any other medical conditions including poorly controlled diabetes, dehydration, heart problems (such as heart failure or history of heart attack), swelling of the hands, feet, or ankles (edema), high blood pressure, history of stroke, blood clotting problems, stomach or bowel problems (such as bleeding or ulcers), history of tobacco use or alcohol use, kidney problems, liver problems, blood or bleeding problems (such as anemia), asthma, growths in the nose (nasal polyps), any allergies (especially history of angioedema with symptoms of lip, tongue, throat swelling), pregnancy, or breast-feeding. USE OF THIS MEDICINE IS NOT RECOMMENDED if you have a history of allergy to aspirin or other NSAIDs (e.g., naproxen, celecoxib). USE OF THIS MEDICINE IS NOT RECOMMENDED if you have history of severe kidney disease or if you are going to have or have recently had coronary artery heart bypass (CABG) surgery. Contact your doctor or pharmacist if you have any questions or concerns about taking this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for using this medicine provided by your doctor. This medicine may come with a medication guide. Read it carefully. Ask your doctor, nurse, or pharmacist any questions that you may have about this medicine. TAKE THIS MEDICINE with a full glass (8 oz./240 ml) of water. DO NOT lie down for 30 minutes after taking this medicine. The dosage is based on your medical condition and response to therapy. If repeat doses are needed, they are usually given 6 or 8 hours apart, or as directed by your doctor. When used in children, the dose is based on your child's weight. Read the product instructions to find the appropriate dose for your child's weight. Consult the pharmacist or doctor if you have questions or if you need help in choosing the appropriate dosage form. THIS MEDICINE MAY BE TAKEN WITH FOOD if it upsets your stomach. Taking it with food may not decrease the risk of stomach or bowel problems (such as bleeding or ulcers) that may occur while taking this medicine. Talk with your doctor or pharmacist if you experience persistent stomach upset. STORE THIS MEDICINE at room temperature, away from heat and light. Do not store in the bathroom. IF YOU MISS A DOSE OF THIS MEDICINE, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.

**CAUTIONS:** THIS MEDICINE INCREASES YOUR RISK OF SERIOUS STOMACH OR BOWEL PROBLEMS (such as ulcers and bleeding). This risk is increased if you are elderly or are in poor health, if you have a history of smoking or drinking alcohol, if you take corticosteroid medicines (such as prednisone) or "blood thinners" (such as warfarin), or if you take this medicine for a long period of time. THIS MEDICINE MAY ALSO INCREASE YOUR RISK for certain serious heart and blood vessel problems (such as heart attack and stroke). TAKE THIS MEDICINE EXACTLY AS PRESCRIBED BY YOUR DOCTOR, at the lowest possible dose for the

shortest time needed. Talk with your doctor or pharmacist for further information. DO NOT TAKE THIS MEDICINE IF YOU HAVE HAD A SEVERE ALLERGIC REACTION to aspirin or any medicine containing aspirin or to a nonsteroidal anti-inflammatory drug (such as Feldene, Motrin, Naprosyn, Clinoril). A severe reaction includes a severe rash, hives, breathing difficulties, or dizziness. If you have a question about whether you are allergic to this medicine or if a certain medicine is a nonsteroidal anti-inflammatory drug, contact your doctor or pharmacist. DO NOT EXCEED THE RECOMMENDED DOSE or take this medicine for longer than 10 days for pain or 3 days for fever, unless directed by your doctor. Laboratory and/or medical tests, including blood counts, liver function tests, and kidney function tests, may be performed to monitor your progress or to check for side effects, especially if you are taking this medicine for a long period of time. KEEP ALL DOCTOR AND LABORATORY APPOINTMENTS while you are taking this medicine. DO NOT DRIVE, OPERATE MACHINERY, OR DO ANYTHING ELSE THAT COULD BE DANGEROUS until you know how you react to this medicine. ALCOHOL WARNING: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take this medicine or other pain relievers/fever reducers. BEFORE YOU BEGIN TAKING ANY NEW MEDICINE, either prescription or over-the-counter, check with your doctor or pharmacist. If you are also taking aspirin, as prescribed by your doctor for reasons such as heart attack or stroke prevention (usually these dosages are 81-325 mg per day), continue to take the aspirin and consult your doctor or pharmacist before using this medicine. CAUTION IS ADVISED WHEN USING THIS MEDICINE IN THE ELDERLY because they may be more sensitive to the effects of this medicine, especially the risk of stomach or bowel effects (such as bleeding or ulcers), or kidney effects. FOR WOMEN: USE OF THIS MEDICINE DURING PREGNANCY has resulted in fetal and newborn death. If you think you may be pregnant, contact your doctor immediately. THIS MEDICINE IS EXCRETED IN BREAST MILK. IF YOU ARE OR WILL BE BREAST-FEEDING while you are using this medicine, check with your doctor or pharmacist to discuss the risks to your baby.

**POSSIBLE SIDE EFFECTS:** SIDE EFFECTS, that may go away during treatment, include nausea, vomiting, diarrhea, gas, constipation, indigestion, dizziness, lightheadedness, drowsiness, or headache. If they continue or are bothersome, check with your doctor. CHECK WITH YOUR DOCTOR AS SOON AS POSSIBLE if you experience ringing in ears. CONTACT YOUR DOCTOR IMMEDIATELY if you experience rapid or pounding heartbeat; easy bruising or bleeding; very stiff neck; or mental/mood changes. CONTACT YOUR DOCTOR IMMEDIATELY if you experience sharp or crushing chest pain; sudden shortness of breath; sudden leg pain; sudden severe headache, vomiting, dizziness, or fainting; changes in vision; numbness of an arm or leg; slurred speech; one-sided weakness; sudden unexplained weight gain; change in amount of urine produced; severe or persistent stomach pain; vomit that looks like coffee grounds; black tarry stools; itching, reddened, swollen, blistered, painful, or peeling skin; yellowing of the skin or eyes; dark urine; right-sided tenderness; severe or persistent tiredness; fever, chills, or sore throat; severe or persistent nausea; swelling of hands, ankles, feet, face, lips, eyes, throat, or tongue; difficulty swallowing or breathing; or hoarseness. AN ALLERGIC REACTION TO THIS MEDICINE is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash, itching, swelling, severe dizziness, or trouble breathing. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms of overdose may include severe stomach pain, coffee ground-like vomit, unusually fast or slow heartbeat, trouble breathing, extreme drowsiness, loss of consciousness, and seizures.

**ADDITIONAL INFORMATION:** DO NOT SHARE THIS MEDICINE with others for whom it was not prescribed. DO NOT USE THIS MEDICINE for other health conditions. KEEP THIS MEDICINE out of the reach of children and pets. IF USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME, obtain refills before your supply runs out.

**KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.**

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)417-7627

**RX # 0515796-09015**

DATE: 04/30/13

**IBUPROFEN 600MG TABLETS**

QTY: 90 1 REFILL BEFORE 04/30/14

New-E NDC: 53746-0465-05

Retail Price: \$21.50 Your Insurance Saved You: \$13.60

**\$ 7.90**

W. CHRISTENSEN, MD  
MFG: AMNEAL  
SDB/HYK/HYK/HYK/HYK

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 131203502028034999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Customer  
Receipt

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)417-7627

**RX # 0515796-09015**

DATE: 04/30/13

**IBUPROFEN 600MG TABLETS**

QTY: 90 1 REFILL BEFORE 04/30/14

New-E NDC: 53746-0465-05

Retail Price: \$21.50 Your Insurance Saved You: \$13.60

**\$ 7.90**

W. CHRISTENSEN, MD  
MFG: AMNEAL  
SDB/HYK/HYK/HYK/HYK

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 131203502028034999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Duplicate  
Receipt

Pharmacy use only

TUE 1:09PM

New-E


IBUPROFEN 600MG TABLETS

53746-0465-05

FAST RACK

QTY 90

30 DRAM



WHITE  
FRONT: IP 465

SDB/HYK/HYK/HYK/HYK

Med Guide

Do not flush unused medications or pour down a sink or drain.

WIC# 216382



Call your doctor for medical advice about side effects.  
You may report side effects to FDA at 1-800-FDA-1088.

**INGREDIENT NAME:** TRAMADOL (TRAM-a-dol)

**COMMON USES:** This medicine is an analgesic used to treat moderate to moderately severe pain.

**BEFORE USING THIS MEDICINE:** Some medicines or medical conditions may interact with this medicine. INFORM YOUR DOCTOR OR PHARMACIST of all prescription and over-the-counter medicine that you are taking. DO NOT TAKE THIS MEDICINE if you are taking sodium oxybate (GHB), or another product that contains tramadol. ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION may be needed if you are taking alpha-2 receptor blockers (eg, yohimbine), anorexiant (eg, phentermine), anticoagulants (eg, warfarin), azole antifungals (eg, ketoconazole), butyrophenones (eg, haloperidol), carbamazepine, cyclobenzaprine, digoxin, furazolidone, linezolid, lithium, loxapine, macrolide antibiotics (eg, erythromycin), certain medicines for mental or mood disorders (eg, olanzapine), molidone, monoamine oxidase inhibitors (MAOIs) (eg, phenelzine, selegiline), nefazodone, opioid pain medicine (eg, codeine, hydrocodone), phenothiazines (eg, promethazine), quinidine, rifampin, serotonin-norepinephrine reuptake inhibitors (SNRIs) (eg, duloxetine), selective serotonin reuptake inhibitors (SSRIs) (eg, fluoxetine, paroxetine), sleeping medicines (eg, zolpidem), St. John's wort, thioxanthenes (eg, thiothixene), tiagabine, tricyclic antidepressants (eg, amitriptyline), or "triptans" (eg, sumatriptan). DO NOT START OR STOP any medicine without doctor or pharmacist approval. Inform your doctor of any other medical conditions, including liver, kidney, or thyroid problems; blockage in the bladder; diabetes; heart problems (eg, cor pulmonale); high blood pressure; pancreas problems; prostate problems; metabolism problems; allergies; pregnancy; or breast-feeding. Tell your doctor if you have or recently have had any head injury, brain injury or tumor, increased pressure in the brain, or infection of the brain or nervous system. Tell your doctor if you have a history of lung or breathing problems (eg, asthma, chronic obstructive pulmonary disease / COPD); seizures (eg, epilepsy); recent stomach or bowel surgery; other stomach or bowel problems (eg, pain, inflammation, ulcers); alcohol or other substance abuse; mood or mental problems (eg, depression); or suicidal thoughts or actions. Tell your doctor if you drink alcohol or if you are going through withdrawal from alcohol or other substances. USE OF THIS MEDICINE IS NOT RECOMMENDED if you have severe or persistent diarrhea caused by antibiotic use; suicidal thoughts or actions; or a history of alcohol or other substance abuse or addiction. USE OF THIS MEDICINE IS NOT RECOMMENDED if you are intoxicated with alcohol, opioids or narcotics (eg, codeine, morphine), or sedatives or sleeping medicines (eg, temazepam, zolpidem). This medicine should not be used in CHILDREN younger than 16 years; safety and effectiveness in these children have not been confirmed. Contact your doctor or pharmacist if you have any questions or concerns about using this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for taking this medicine provided by your doctor. TAKE THIS MEDICINE by mouth with or without food. STORE THIS MEDICINE at 77 degrees F (25 degrees C) away from heat, moisture, and light. Do not store in the bathroom. Brief storage at temperatures between 59 and 86 degrees F (15 and 30 degrees C) is permitted. KEEP THIS MEDICINE out of the reach of children and away from pets. IF YOU MISS A DOSE OF THIS MEDICINE and you are taking it regularly, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. DO NOT take 2 doses at once.

**CAUTIONS:** DO NOT TAKE THIS MEDICINE if you have had an allergic reaction to it or if you are allergic to any ingredient in this product. DO NOT TAKE THIS MEDICINE if you have had a severe allergic reaction to codeine or another opioid (eg, morphine). DO NOT EXCEED THE RECOMMENDED DOSE or take this medicine for longer than prescribed without checking with your doctor. Exceeding the recommended dose or taking this medicine for longer than prescribed may be habit-forming or may increase the risk of seizures or breathing problems. If using this medicine for an extended period of time, DO NOT SUDDENLY STOP taking this medicine without your doctor's approval. Your dose may need to be slowly lowered to avoid side effects. When used for an extended period of time, this medicine may not work as well and may require different dosing. Talk with your doctor if this medicine stops working well. DO NOT TAKE MORE THAN PRESCRIBED. BEFORE YOU HAVE ANY MEDICAL OR DENTAL TREATMENTS, EMERGENCY CARE, OR SURGERY, tell the doctor or dentist that you are using this medicine. DO NOT DRINK ALCOHOL while you are using this medicine. CHECK WITH YOUR DOCTOR BEFORE you use medicines that may cause drowsiness (eg, sleep aids, muscle relaxers, narcotic pain medicines) while you are using this medicine; it may add to their effects. Ask your pharmacist if you

have questions about which medicines may cause drowsiness. This medicine may cause drowsiness or dizziness. DO NOT DRIVE, OPERATE MACHINERY, OR DO ANYTHING ELSE THAT COULD BE DANGEROUS until you know how you react to this medicine. Using this medicine alone, with other medicines, or with alcohol may lessen your ability to drive or to perform other potentially dangerous tasks. Alcohol, hot weather, exercise, or fever may increase dizziness. IF DIZZINESS OCCURS, sit up or stand slowly, especially in the morning. Sit or lie down at the first sign of this effect. THIS MEDICINE MAY INCREASE YOUR RISK OF SEIZURES. Your risk may be greater if you also have certain medical conditions, use certain medicines, or if you use a lot of alcohol. Talk to your doctor to see if you may have a greater risk of seizures while taking this medicine. SEROTONIN SYNDROME is a possibly fatal syndrome that can be caused by this medicine. Your risk may be greater if you take this medicine with certain other medicines (eg, "triptans," MAOIs, antidepressants). Symptoms may include agitation; confusion; hallucinations; coma; fever; fast or irregular heartbeat; tremor; excessive sweating; and nausea, vomiting, or diarrhea. Contact your doctor at once if you have any of these symptoms. BEFORE YOU BEGIN TAKING ANY NEW MEDICINES, either prescription or over-the-counter, check with your doctor or pharmacist. CAUTION IS ADVISED WHEN USING THIS MEDICINE IN THE ELDERLY because they may be more sensitive to the effects of this medicine. FOR WOMEN: This medicine may cause harm to the fetus. IF YOU THINK YOU MAY BE PREGNANT, discuss with your doctor the benefits and risks of using this medicine during pregnancy. THIS MEDICINE IS EXCRETED in breast milk. DO NOT BREAST-FEED while taking this medicine.

**POSSIBLE SIDE EFFECTS:** SIDE EFFECTS that may occur while taking this medicine include constipation; diarrhea; dizziness; drowsiness; dry mouth; headache; indigestion; mildly increased sweating; mild itching; nausea; trouble sleeping; vomiting; weakness. If they continue or are bothersome, check with your doctor. CONTACT YOUR DOCTOR IMMEDIATELY if you experience agitation; chest pain; confusion; difficult or painful urination; excessive sweating; fainting; fast or irregular heartbeat; fever; hallucinations; loss of coordination; mood or mental changes (eg, depression); red, blistered, swollen, or peeling skin; seizures; severe dizziness or lightheadedness; severe nausea, vomiting, or diarrhea; severe or persistent headache; slow or shallow breathing; suicidal thoughts or behaviors; tremor; or vision problems. AN ALLERGIC REACTION to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, or tongue; unusual hoarseness. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your health care provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** IF OVERDOSE IS SUSPECTED, contact your local poison control center or emergency room immediately. Symptoms may include bluish skin; cold, clammy skin; difficult, shallow, or slow breathing; drowsiness leading to unresponsiveness or coma; excessive sweating; limp muscles; pinpoint pupils; seizures; slow or irregular heartbeat.

**ADDITIONAL INFORMATION:** DO NOT SHARE THIS MEDICINE with others for whom it was not prescribed. DO NOT USE THIS MEDICINE for other health conditions. IF YOU WILL BE USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME, be sure to obtain necessary refills before your supply runs out. CHECK WITH YOUR PHARMACIST about how to dispose of unused medicine.

**KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.**

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)417-7627

**RX # 0498533-09015**

DATE: 02/08/13

**TRAMADOL 50MG TABLETS**

QTY: 90 NO REFILLS - DR. AUTH REQUIRED

New-E NDC: 68382-0319-10

Retail Price: \$46.49 Your Insurance Saved You: \$36.49

\$ 10.00

W. CHRISTENSEN, MD  
MFG: ZYDUS  
CAM/ADO/ADO/ /ADO

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 130394454708034998

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Customer  
Receipt

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)417-7627

**RX # 0498533-09015**

DATE: 02/08/13

**TRAMADOL 50MG TABLETS**

QTY: 90 NO REFILLS - DR. AUTH REQUIRED

New-E NDC: 68382-0319-10

Retail Price: \$46.49 Your Insurance Saved You: \$36.49

\$ 10.00

W. CHRISTENSEN, MD  
MFG: ZYDUS  
CAM/ADO/ADO/ /ADO

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 130394454708034998

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Duplicate  
Receipt

Pharmacy use only

FRI 3:14PM  
New-E

TRAMADOL 50MG TABLETS  
68382-0319-10  
FAST RACK

QTY 90  
10 DRAM



WHITE  
FRONT: 319

CAM/ADO/ADO/ /ADO

Do not flush unused medications or pour down a sink or drain.

WIC# 216382



Call your doctor for medical advice about side effects.  
You may report side effects to FDA at 1-800-FDA-1088.

**INGREDIENT NAME:** METHYLPREDNISOLONE  
(meth-ill-pred-NISS-oh-lone)

**COMMON USES:** This medicine is a corticosteroid used to treat severe allergies, arthritis, asthma, and skin conditions. It may also be used to treat other conditions as determined by your doctor.

**BEFORE USING THIS MEDICINE:** Some medicines or medical conditions may interact with this medicine. INFORM YOUR DOCTOR OR PHARMACIST of all prescription and over-the-counter medicine that you are taking. DO NOT TAKE THIS MEDICINE if you are also taking mifepristone. ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION may be needed if you are taking aspirin, barbiturates, cyclosporine, licorice (glycyrrhiza), macrolide antibiotics (e.g., erythromycin), indomethacin, estrogen (e.g., birth control pills), ketoconazole, rifampin, troleandomycin, or medicine for seizures. USE OF THIS MEDICINE IS NOT RECOMMENDED if you have a fungal infection in the body. Inform your doctor of any medical conditions including heart disease, liver disease, kidney disease, HIV infection, high blood pressure, peptic ulcer, osteoporosis, myasthenia gravis, herpes eye infection, low levels of thyroid hormone (hypothyroidism), diabetes, tuberculosis, mental/mood disorders, seizures, recent infection, allergies, pregnancy, or breast-feeding. Contact your doctor or pharmacist if you have any questions or concerns about taking this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for using this medicine provided by your doctor. The directions for taking this medicine properly are also on the package. This medicine may be taken on an empty stomach or with food. Taking it with food may help to decrease the chance of stomach upset. Ask your doctor or pharmacist any questions you may have about how you should take this medicine. STORE THIS MEDICINE at room temperature, away from heat and light. IF YOU MISS A DOSE OF THIS MEDICINE and you are taking 1 dose daily, take it as soon as possible. If you do not remember until the next day, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once. For all other dosing regimens: if you miss a dose, contact your doctor or pharmacist promptly. Your dosing schedule may need to be adjusted.

**CAUTIONS:** DO NOT STOP TAKING THIS MEDICINE without checking with your doctor. Stopping this medicine suddenly may cause serious side effects. KEEP ALL DOCTOR AND LABORATORY APPOINTMENTS while you are using this medicine. BEFORE YOU HAVE ANY MEDICAL OR DENTAL TREATMENTS, EMERGENCY CARE, OR SURGERY, tell the doctor or dentist that you are using this medicine. CORTICOSTEROIDS MAY LOWER YOUR ABILITY TO FIGHT INFECTION or may cause infections to be more severe. The risk may be greater if you take them for a long time. Avoid contact with people who have infections such as colds, chickenpox, measles, or tuberculosis (TB). Tell your doctor right away if you have contact with anyone who has these infections. Tell your doctor if you develop symptoms such as fever, sore throat, rash, or chills. Check with your doctor if you have questions about your risk of infection. CHECK WITH YOUR DOCTOR BEFORE HAVING IMMUNIZATIONS (VACCINATIONS) while you are using this medicine. BEFORE YOU BEGIN TAKING ANY NEW MEDICINE, either

prescription or over-the-counter, check with your doctor or pharmacist. IF YOU PLAN ON BECOMING PREGNANT, discuss with your doctor the benefits and risks of using this medicine during pregnancy. IF YOU ARE OR WILL BE BREAST-FEEDING while you are using this medicine, check with your doctor or pharmacist to discuss the risks to your baby. IF YOU HAVE DIABETES, this medicine may affect your blood sugar. Check blood sugar levels closely and ask your doctor before adjusting the dose of your diabetes medicine.

**POSSIBLE SIDE EFFECTS:** SIDE EFFECTS that may occur while you are taking this medicine include dizziness; facial flushing; feeling of whirling motion; headache; or increased sweating. If they continue or are bothersome, check with your doctor. CONTACT YOUR DOCTOR IMMEDIATELY if you experience bloody, black, or tarry stools; changes in body fat; changes in menstrual periods; chest pain; fainting; fever, chills, or sore throat; increased hunger, thirst, or urination; mental or mood changes; muscle pain, weakness, or wasting; seizures; severe nausea or vomiting; slow wound healing; stomach pain; sudden severe dizziness or headache; swelling of feet or legs; tendon, bone, or joint pain; thinning of the skin; unusual bruising or bleeding; unusual skin sensations; unusual weight gain; vision changes or other eye problems; or vomit that looks like coffee grounds. AN ALLERGIC REACTION to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; or swelling of the mouth, face, lips, or tongue. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** If overdose is suspected, contact your local poison control center or emergency room immediately.

**ADDITIONAL INFORMATION:** Patients taking this medicine for long-term therapy should carry a warning card or medical ID bracelet that identifies the possible need of supplemental steroids during periods of stress. DO NOT SHARE THIS MEDICINE with others for whom it was not prescribed. DO NOT USE THIS MEDICINE for other health conditions. KEEP THIS MEDICINE out of the reach of children.

**KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.**

**ROBERT A PLOCK**  
6827 Latta Pkwy, Dallas, TX 75227  
(214)417-7627

**RX # 0498536-09015**

DATE: 02/08/13

**METHYLPREDNISOLONE 4MG DOSPAK 21'S**

QTY: 21 NO REFILLS - DR. AUTH REQUIRED

New-E NDC:59746-0001-03

Retail Price: \$27.99 Your Insurance Saved You: \$17.99

**\$ 10.00**

W. CHRISTENSEN, MD  
MFG:CADISTA  
CAM/ADO/ADO/ ADO

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 130394459256047999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Customer  
Receipt

**ROBERT A PLOCK**  
6827 Latta Pkwy, Dallas, TX 75227  
(214)417-7627

**RX # 0498536-09015**

DATE: 02/08/13

**METHYLPREDNISOLONE 4MG DOSPAK 21'S**

QTY: 21 NO REFILLS - DR. AUTH REQUIRED

New-E NDC:59746-0001-03

Retail Price: \$27.99 Your Insurance Saved You: \$17.99

**\$ 10.00**

W. CHRISTENSEN, MD  
MFG:CADISTA  
CAM/ADO/ADO/ ADO

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 130394459256047999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951


Duplicate  
Receipt

Pharmacy use only

FRI 3:14PM  
New-E

METHYLPREDNISOLONE 4MG DOSPAK 21'S  
59746-0001-03  
UNIT OF USE

QTY 21



WHITE  
FRONT: TL 001

CAM/ADO/ADO/ ADO

Do not flush unused medications or pour down a sink or drain.

WIC# 216382



Call your doctor for medical advice about side effects.  
You may report side effects to FDA at 1-800-FDA-1088.

**INGREDIENT NAME:** IBUPROFEN (eye-byoo-PROE-fen)

**COMMON USES:** This medicine is a nonsteroidal anti-inflammatory drug (NSAID) used to treat mild to moderate pain, osteoarthritis, and rheumatoid arthritis. It may also be used to treat other conditions as determined by your doctor.

**BEFORE USING THIS MEDICINE:** WARNING: THE RISK OF SERIOUS AND SOMETIMES FATAL HEART PROBLEMS, HEART ATTACK, AND STROKE may be increased with the use of this medicine. This risk may be increased the longer you use this medicine. Risk may also be higher in patients who have heart problems or who are at risk for heart problems. THIS MEDICINE SHOULD NOT BE USED to treat pain before or after coronary artery heart bypass (CABG) surgery. THE RISK OF SERIOUS AND SOMETIMES FATAL STOMACH AND BOWEL PROBLEMS, including bleeding, ulcers, and holes in the stomach and bowel, is increased while using this medicine. These problems may occur at any time during therapy, with or without symptoms. Elderly patients are at higher risk for serious stomach problems. Ask your doctor or pharmacist for more information about this medicine and its side effects. Some medicines or medical conditions may interact with this medicine. INFORM YOUR DOCTOR OR PHARMACIST of all prescription and over-the-counter medicine that you are taking. DO NOT TAKE THIS MEDICINE if you are also taking heparins or tacrolimus. ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION may be needed if you are taking serotonin reuptake blocker medicines such as fluoxetine or citalopram, "blood thinners" such as warfarin, bisphosphonates such as alendronate or risedronate, cyclosporine, corticosteroids such as prednisone, high blood pressure medicines (including ACE inhibitors such as captopril, angiotensin II receptor antagonists such as losartan, and beta-blockers such as metoprolol), "water pills" (diuretics such as furosemide, hydrochlorothiazide, triamterene), lithium, methotrexate, or aspirin. DO NOT START OR STOP ANY MEDICINE without doctor or pharmacist approval. Inform your doctor of any other medical conditions including poorly controlled diabetes, dehydration, heart problems (such as heart failure or history of heart attack), swelling of the hands, feet, or ankles (edema), high blood pressure, history of stroke, blood clotting problems, stomach or bowel problems (such as bleeding or ulcers), history of tobacco use or alcohol use, kidney problems, liver problems, blood or bleeding problems (such as anemia), asthma, growths in the nose (nasal polyps), any allergies (especially history of angioedema with symptoms of lip, tongue, throat swelling), pregnancy, or breast-feeding. USE OF THIS MEDICINE IS NOT RECOMMENDED if you have a history of allergy to aspirin or other NSAIDs (e.g., naproxen, celecoxib). USE OF THIS MEDICINE IS NOT RECOMMENDED if you have history of severe kidney disease or if you are going to have or have recently had coronary artery heart bypass (CABG) surgery. Contact your doctor or pharmacist if you have any questions or concerns about taking this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for using this medicine provided by your doctor. This medicine may come with a medication guide. Read it carefully. Ask your doctor, nurse, or pharmacist any questions that you may have about this medicine. TAKE THIS MEDICINE with a full glass (8 oz./240 ml) of water. DO NOT lie down for 30 minutes after taking this medicine. The dosage is based on your medical condition and response to therapy. If repeat doses are needed, they are usually given 6 or 8 hours apart, or as directed by your doctor. When used in children, the dose is based on your child's weight. Read the product instructions to find the appropriate dose for your child's weight. Consult the pharmacist or doctor if you have questions or if you need help in choosing the appropriate dosage form. THIS MEDICINE MAY BE TAKEN WITH FOOD if it upsets your stomach. Taking it with food may not decrease the risk of stomach or bowel problems (such as bleeding or ulcers) that may occur while taking this medicine. Talk with your doctor or pharmacist if you experience persistent stomach upset. STORE THIS MEDICINE at room temperature, away from heat and light. Do not store in the bathroom. IF YOU MISS A DOSE OF THIS MEDICINE, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.

**CAUTIONS:** THIS MEDICINE INCREASES YOUR RISK OF SERIOUS STOMACH OR BOWEL PROBLEMS (such as ulcers and bleeding). This risk is increased if you are elderly or are in poor health, if you have a history of smoking or drinking alcohol, if you take corticosteroid medicines (such as prednisone) or "blood thinners" (such as warfarin), or if you take this medicine for a long period of time. THIS MEDICINE MAY ALSO INCREASE YOUR RISK for certain serious heart and blood vessel problems (such as heart attack and stroke). TAKE THIS MEDICINE EXACTLY AS PRESCRIBED BY YOUR DOCTOR, at the lowest possible dose for the

shortest time needed. Talk with your doctor or pharmacist for further information. DO NOT TAKE THIS MEDICINE IF YOU HAVE HAD A SEVERE ALLERGIC REACTION to aspirin or any medicine containing aspirin or to a nonsteroidal anti-inflammatory drug (such as Feldene, Motrin, Naprosyn, Clinoril). A severe reaction includes a severe rash, hives, breathing difficulties, or dizziness. If you have a question about whether you are allergic to this medicine or if a certain medicine is a nonsteroidal anti-inflammatory drug, contact your doctor or pharmacist. DO NOT EXCEED THE RECOMMENDED DOSE or take this medicine for longer than 10 days for pain or 3 days for fever, unless directed by your doctor. Laboratory and/or medical tests, including blood counts, liver function tests, and kidney function tests, may be performed to monitor your progress or to check for side effects, especially if you are taking this medicine for a long period of time. KEEP ALL DOCTOR AND LABORATORY APPOINTMENTS while you are taking this medicine. DO NOT DRIVE, OPERATE MACHINERY, OR DO ANYTHING ELSE THAT COULD BE DANGEROUS until you know how you react to this medicine. ALCOHOL WARNING: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take this medicine or other pain relievers/fever reducers. BEFORE YOU BEGIN TAKING ANY NEW MEDICINE, either prescription or over-the-counter, check with your doctor or pharmacist. If you are also taking aspirin, as prescribed by your doctor for reasons such as heart attack or stroke prevention (usually these dosages are 81-325 mg per day), continue to take the aspirin and consult your doctor or pharmacist before using this medicine. CAUTION IS ADVISED WHEN USING THIS MEDICINE IN THE ELDERLY because they may be more sensitive to the effects of this medicine, especially the risk of stomach or bowel effects (such as bleeding or ulcers), or kidney effects. FOR WOMEN: USE OF THIS MEDICINE DURING PREGNANCY has resulted in fetal and newborn death. If you think you may be pregnant, contact your doctor immediately. THIS MEDICINE IS EXCRETED IN BREAST MILK. IF YOU ARE OR WILL BE BREAST-FEEDING while you are using this medicine, check with your doctor or pharmacist to discuss the risks to your baby.

**POSSIBLE SIDE EFFECTS:** SIDE EFFECTS, that may go away during treatment, include nausea, vomiting, diarrhea, gas, constipation, indigestion, dizziness, lightheadedness, drowsiness, or headache. If they continue or are bothersome, check with your doctor. CHECK WITH YOUR DOCTOR AS SOON AS POSSIBLE if you experience ringing in ears. CONTACT YOUR DOCTOR IMMEDIATELY if you experience rapid or pounding heartbeat; easy bruising or bleeding; very stiff neck; or mental/mood changes. CONTACT YOUR DOCTOR IMMEDIATELY if you experience sharp or crushing chest pain; sudden shortness of breath; sudden leg pain; sudden severe headache, vomiting, dizziness, or fainting; changes in vision; numbness of an arm or leg; slurred speech; one-sided weakness; sudden unexplained weight gain; change in amount of urine produced; severe or persistent stomach pain; vomit that looks like coffee grounds; black tarry stools; itching, reddened, swollen, blistered, painful, or peeling skin; yellowing of the skin or eyes; dark urine; right-sided tenderness; severe or persistent tiredness; fever, chills, or sore throat; severe or persistent nausea; swelling of hands, ankles, feet, face, lips, eyes, throat, or tongue; difficulty swallowing or breathing; or hoarseness. AN ALLERGIC REACTION TO THIS MEDICINE is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash, itching, swelling, severe dizziness, or trouble breathing. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms of overdose may include severe stomach pain, coffee ground-like vomit, unusually fast or slow heartbeat, trouble breathing, extreme drowsiness, loss of consciousness, and seizures.

**ADDITIONAL INFORMATION:** DO NOT SHARE THIS MEDICINE with others for whom it was not prescribed. DO NOT USE THIS MEDICINE for other health conditions. KEEP THIS MEDICINE out of the reach of children and pets. IF USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME, obtain refills before your supply runs out.

**KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.**

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)417-7627

**RX # 0503783-09015**

DATE: 03/04/13

**IBUPROFEN 600MG TABLETS**

QTY: 30 1 REFILL BEFORE 03/04/14

New NDC: 53746-0465-05

Retail Price: \$11.99 Your Insurance Saved You: \$ 8.57

**\$ 3.42**

W. CHRISTENSEN, MD  
MFG: AMNEAL  
PLB/HYK/HYK/ /HYK

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 130635701275028999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Customer  
Receipt

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)417-7627

**RX # 0503783-09015**

DATE: 03/04/13

**IBUPROFEN 600MG TABLETS**

QTY: 30 1 REFILL BEFORE 03/04/14

New NDC: 53746-0465-05

Retail Price: \$11.99 Your Insurance Saved You: \$ 8.57

**\$ 3.42**

W. CHRISTENSEN, MD  
MFG: AMNEAL  
PLB/HYK/HYK/ /HYK

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 130635701275028999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951


Duplicate  
Receipt

Pharmacy use only

MON 6:33PM  
New

IBUPROFEN 600MG TABLETS  
53746-0465-05  
FAST RACK

QTY 30  
10 DRAM



WHITE  
FRONT: IP 465

PLB/HYK/HYK/ /HYK

Med Guide

Do not flush unused medications or pour down a sink or drain.

WIC# 216382



Call your doctor for medical advice about side effects.  
You may report side effects to FDA at 1-800-FDA-1088.

#### INGREDIENT NAME: TRAMADOL (TRAM-a-dol)

**COMMON USES:** This medicine is an analgesic used to treat moderate to moderately severe pain.

**BEFORE USING THIS MEDICINE:** Some medicines or medical conditions may interact with this medicine. **INFORM YOUR DOCTOR OR PHARMACIST** of all prescription and over-the-counter medicine that you are taking. **DO NOT TAKE THIS MEDICINE** if you are taking sodium oxybate (GHB), or another product that contains tramadol. **ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION** may be needed if you are taking alpha-2 receptor blockers (eg, yohimbine), anorexiant (eg, phentermine), anticoagulants (eg, warfarin), azole antifungals (eg, ketoconazole), butyrophenones (eg, haloperidol), carbamazepine, cyclobenzaprine, digoxin, furazolidone, linezolid, lithium, loxapine, macrolide antibiotics (eg, erythromycin), certain medicines for mental or mood disorders (eg, olanzapine), molidone, monoamine oxidase inhibitors (MAOIs) (eg, phenelzine, selegiline), nefazodone, opioid pain medicine (eg, codeine, hydrocodone), phenothiazines (eg, promethazine), quinidine, rifampin, serotonin-norepinephrine reuptake inhibitors (SNRIs) (eg, duloxetine), selective serotonin reuptake inhibitors (SSRIs) (eg, fluoxetine, paroxetine), sleeping medicines (eg, zolpidem), St. John's wort, thioxanthenes (eg, thiothixene), tiagabine, tricyclic antidepressants (eg, amitriptyline), or "triptans" (eg, sumatriptan). **DO NOT START OR STOP** any medicine without doctor or pharmacist approval. Inform your doctor of any other medical conditions, including liver, kidney, or thyroid problems; blockage in the bladder; diabetes; heart problems (eg, cor pulmonale); high blood pressure; pancreas problems; prostate problems; metabolism problems; allergies; pregnancy; or breast-feeding. Tell your doctor if you have or recently have had any head injury, brain injury or tumor, increased pressure in the brain, or infection of the brain or nervous system. Tell your doctor if you have a history of lung or breathing problems (eg, asthma, chronic obstructive pulmonary disease / COPD); seizures (eg, epilepsy); recent stomach or bowel surgery; other stomach or bowel problems (eg, pain, inflammation, ulcers); alcohol or other substance abuse; mood or mental problems (eg, depression); or suicidal thoughts or actions. Tell your doctor if you drink alcohol or if you are going through withdrawal from alcohol or other substances. **USE OF THIS MEDICINE IS NOT RECOMMENDED** if you have severe or persistent diarrhea caused by antibiotic use; suicidal thoughts or actions; or a history of alcohol or other substance abuse or addiction. **USE OF THIS MEDICINE IS NOT RECOMMENDED** if you are intoxicated with alcohol, opioids or narcotics (eg, codeine, morphine), or sedatives or sleeping medicines (eg, temazepam, zolpidem). This medicine should not be used in CHILDREN younger than 16 years; safety and effectiveness in these children have not been confirmed. Contact your doctor or pharmacist if you have any questions or concerns about using this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for taking this medicine provided by your doctor. **TAKE THIS MEDICINE** by mouth with or without food. **STORE THIS MEDICINE** at 77 degrees F (25 degrees C) away from heat, moisture, and light. Do not store in the bathroom. Brief storage at temperatures between 59 and 86 degrees F (15 and 30 degrees C) is permitted. **KEEP THIS MEDICINE** out of the reach of children and away from pets. **IF YOU MISS A DOSE OF THIS MEDICINE** and you are taking it regularly, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. **DO NOT** take 2 doses at once.

**CAUTIONS:** **DO NOT TAKE THIS MEDICINE** if you have had an allergic reaction to it or if you are allergic to any ingredient in this product. **DO NOT TAKE THIS MEDICINE** if you have had a severe allergic reaction to codeine or another opioid (eg, morphine). **DO NOT EXCEED THE RECOMMENDED DOSE** or take this medicine for longer than prescribed without checking with your doctor. Exceeding the recommended dose or taking this medicine for longer than prescribed may be habit-forming or may increase the risk of seizures or breathing problems. If using this medicine for an extended period of time, **DO NOT SUDDENLY STOP** taking this medicine without your doctor's approval. Your dose may need to be slowly lowered to avoid side effects. When used for an extended period of time, this medicine may not work as well and may require different dosing. Talk with your doctor if this medicine stops working well. **DO NOT TAKE MORE THAN PRESCRIBED.** **BEFORE YOU HAVE ANY MEDICAL OR DENTAL TREATMENTS, EMERGENCY CARE, OR SURGERY,** tell the doctor or dentist that you are using this medicine. **DO NOT DRINK ALCOHOL** while you are using this medicine. **CHECK WITH YOUR DOCTOR** before you use medicines that may cause drowsiness (eg, sleep aids, muscle relaxers, narcotic pain medicines) while you are using this medicine; it may add to their effects. Ask your pharmacist if you

have questions about which medicines may cause drowsiness. This medicine may cause drowsiness or dizziness. **DO NOT DRIVE, OPERATE MACHINERY, OR DO ANYTHING ELSE THAT COULD BE DANGEROUS** until you know how you react to this medicine. Using this medicine alone, with other medicines, or with alcohol may lessen your ability to drive or to perform other potentially dangerous tasks. Alcohol, hot weather, exercise, or fever may increase dizziness. **IF DIZZINESS OCCURS,** sit up or stand slowly, especially in the morning. Sit or lie down at the first sign of this effect. **THIS MEDICINE MAY INCREASE YOUR RISK OF SEIZURES.** Your risk may be greater if you also have certain medical conditions, use certain medicines, or if you use a lot of alcohol. Talk to your doctor to see if you may have a greater risk of seizures while taking this medicine. **SEROTONIN SYNDROME** is a possibly fatal syndrome that can be caused by this medicine. Your risk may be greater if you take this medicine with certain other medicines (eg, "triptans," MAOIs, antidepressants). Symptoms may include agitation; confusion; hallucinations; coma; fever; fast or irregular heartbeat; tremor; excessive sweating; and nausea, vomiting, or diarrhea. Contact your doctor at once if you have any of these symptoms. **BEFORE YOU BEGIN TAKING ANY NEW MEDICINES,** either prescription or over-the-counter, check with your doctor or pharmacist. **CAUTION IS ADVISED WHEN USING THIS MEDICINE IN THE ELDERLY** because they may be more sensitive to the effects of this medicine. **FOR WOMEN:** This medicine may cause harm to the fetus. **IF YOU THINK YOU MAY BE PREGNANT,** discuss with your doctor the benefits and risks of using this medicine during pregnancy. **THIS MEDICINE IS EXCRETED** in breast milk. **DO NOT BREAST-FEED** while taking this medicine.

**POSSIBLE SIDE EFFECTS:** SIDE EFFECTS that may occur while taking this medicine include constipation; diarrhea; dizziness; drowsiness; dry mouth; headache; indigestion; mildly increased sweating; mild itching; nausea; trouble sleeping; vomiting; weakness. If they continue or are bothersome, check with your doctor. **CONTACT YOUR DOCTOR IMMEDIATELY** if you experience agitation; chest pain; confusion; difficult or painful urination; excessive sweating; fainting; fast or irregular heartbeat; fever; hallucinations; loss of coordination; mood or mental changes (eg, depression); red, blistered, swollen, or peeling skin; seizures; severe dizziness or lightheadedness; severe nausea, vomiting, or diarrhea; severe or persistent headache; slow or shallow breathing; suicidal thoughts or behaviors; tremor; or vision problems. **AN ALLERGIC REACTION** to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, or tongue; unusual hoarseness. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your health care provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** **IF OVERDOSE IS SUSPECTED,** contact your local poison control center or emergency room immediately. Symptoms may include bluish skin; cold, clammy skin; difficult, shallow, or slow breathing; drowsiness leading to unresponsiveness or coma; excessive sweating; limp muscles; pinpoint pupils; seizures; slow or irregular heartbeat.

**ADDITIONAL INFORMATION:** **DO NOT SHARE THIS MEDICINE** with others for whom it was not prescribed. **DO NOT USE THIS MEDICINE** for other health conditions. **IF YOU WILL BE USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME,** be sure to obtain necessary refills before your supply runs out. **CHECK WITH YOUR PHARMACIST** about how to dispose of unused medicine.

**KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.**

#### ROBERT A PLOCK

6827 Latta Pkwy, Dallas, TX 75227  
(214)417-7627

**RX # 0503784-09015**

DATE: 03/04/13

#### TRAMADOL 50MG TABLETS

QTY: 90 NO REFILLS - DR. AUTH REQUIRED

New NDC: 68382-0319-10

Retail Price: \$46.49 Your Insurance Saved You: \$36.49

\$ 10.00

W. CHRISTENSEN, MD  
MFG: ZYDUS  
PLB/HYK/HYK /HYK

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 130635705595019999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Customer  
Receipt

#### ROBERT A PLOCK

6827 Latta Pkwy, Dallas, TX 75227  
(214)417-7627

**RX # 0503784-09015**

DATE: 03/04/13

#### TRAMADOL 50MG TABLETS

QTY: 90 NO REFILLS - DR. AUTH REQUIRED

New NDC: 68382-0319-10

Retail Price: \$46.49 Your Insurance Saved You: \$36.49

\$ 10.00

W. CHRISTENSEN, MD  
MFG: ZYDUS  
PLB/HYK/HYK /HYK

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 130635705595019999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Duplicate  
Receipt

Pharmacy use only

MON 6:33PM  
New

TRAMADOL 50MG TABLETS  
68382-0319-10  
FAST RACK

QTY 90  
10 DRAM



WHITE  
FRONT: 319

PLB/HYK/HYK /HYK

Do not flush unused medications or pour down a sink or drain.

WIC# 216382



Call your doctor for medical advice about side effects.  
You may report side effects to FDA at 1-800-FDA-1088.

**INGREDIENT NAME:** IBUPROFEN (eye-byoo-PROE-fen)

**COMMON USES:** This medicine is a nonsteroidal anti-inflammatory drug (NSAID) used to treat mild to moderate pain, osteoarthritis, and rheumatoid arthritis. It may also be used to treat other conditions as determined by your doctor.

**BEFORE USING THIS MEDICINE:** WARNING: THE RISK OF SERIOUS AND SOMETIMES FATAL HEART PROBLEMS, HEART ATTACK, AND STROKE may be increased with the use of this medicine. This risk may be increased the longer you use this medicine. Risk may also be higher in patients who have heart problems or who are at risk for heart problems. THIS MEDICINE SHOULD NOT BE USED to treat pain before or after coronary artery heart bypass (CABG) surgery. THE RISK OF SERIOUS AND SOMETIMES FATAL STOMACH AND BOWEL PROBLEMS, including bleeding, ulcers, and holes in the stomach and bowel, is increased while using this medicine. These problems may occur at any time during therapy, with or without symptoms. Elderly patients are at higher risk for serious stomach problems. Ask your doctor or pharmacist for more information about this medicine and its side effects. Some medicines or medical conditions may interact with this medicine. INFORM YOUR DOCTOR OR PHARMACIST of all prescription and over-the-counter medicine that you are taking. DO NOT TAKE THIS MEDICINE if you are also taking heparin or trolimus. ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION may be needed if you are taking serotonin reuptake blocker medicines such as fluoxetine or citalopram, "blood thinners" such as warfarin, bisphosphonates such as alendronate or risedronate, cyclosporine, corticosteroids such as prednisone, high blood pressure medicines (including ACE inhibitors such as captopril, angiotensin II receptor antagonists such as losartan, and beta-blockers such as metoprolol), "water pills" (diuretics such as furosemide, hydrochlorothiazide, triamterene), lithium, methotrexate, or aspirin. DO NOT START OR STOP ANY MEDICINE without doctor or pharmacist approval. Inform your doctor of any other medical conditions including poorly controlled diabetes, dehydration, heart problems (such as heart failure or history of heart attack), swelling of the hands, feet, or ankles (edema), high blood pressure, history of stroke, blood clotting problems, stomach or bowel problems (such as bleeding or ulcers), history of tobacco use or alcohol use, kidney problems, liver problems, blood or bleeding problems (such as anemia), asthma, growths in the nose (nasal polyps), any allergies (especially history of angioedema with symptoms of lip, tongue, throat swelling), pregnancy, or breast-feeding. USE OF THIS MEDICINE IS NOT RECOMMENDED if you have a history of allergy to aspirin or other NSAIDs (e.g., naproxen, celecoxib). USE OF THIS MEDICINE IS NOT RECOMMENDED if you have history of severe kidney disease or if you are going to have or have recently had coronary artery heart bypass (CABG) surgery. Contact your doctor or pharmacist if you have any questions or concerns about taking this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for using this medicine provided by your doctor. This medicine may come with a medication guide. Read it carefully. Ask your doctor, nurse, or pharmacist any questions that you may have about this medicine. TAKE THIS MEDICINE with a full glass (8 oz./240 ml) of water. DO NOT lie down for 30 minutes after taking this medicine. The dosage is based on your medical condition and response to therapy. If repeat doses are needed, they are usually given 6 or 8 hours apart, or as directed by your doctor. When used in children, the dose is based on your child's weight. Read the product instructions to find the appropriate dose for your child's weight. Consult the pharmacist or doctor if you have questions or if you need help in choosing the appropriate dosage form. THIS MEDICINE MAY BE TAKEN WITH FOOD if it upsets your stomach. Taking it with food may not decrease the risk of stomach or bowel problems (such as bleeding or ulcers) that may occur while taking this medicine. Talk with your doctor or pharmacist if you experience persistent stomach upset. STORE THIS MEDICINE at room temperature, away from heat and light. Do not store in the bathroom. IF YOU MISS A DOSE OF THIS MEDICINE, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.

**CAUTIONS:** THIS MEDICINE INCREASES YOUR RISK OF SERIOUS STOMACH OR BOWEL PROBLEMS (such as ulcers and bleeding). This risk is increased if you are elderly or are in poor health, if you have a history of smoking or drinking alcohol, if you take corticosteroid medicines (such as prednisone) or "blood thinners" (such as warfarin), or if you take this medicine for a long period of time. THIS MEDICINE MAY ALSO INCREASE YOUR RISK for certain serious heart and blood vessel problems (such as heart attack and stroke). TAKE THIS MEDICINE EXACTLY AS PRESCRIBED BY YOUR DOCTOR, at the lowest possible dose for the

shortest time needed. Talk with your doctor or pharmacist for further information. DO NOT TAKE THIS MEDICINE IF YOU HAVE HAD A SEVERE ALLERGIC REACTION to aspirin or any medicine containing aspirin or to a nonsteroidal anti-inflammatory drug (such as Feldene, Motrin, Naprosyn, Clinoril). A severe reaction includes a severe rash, hives, breathing difficulties, or dizziness. If you have a question about whether you are allergic to this medicine or if a certain medicine is a nonsteroidal anti-inflammatory drug, contact your doctor or pharmacist. DO NOT EXCEED THE RECOMMENDED DOSE or take this medicine for longer than 10 days for pain or 3 days for fever, unless directed by your doctor. Laboratory and/or medical tests, including blood counts, liver function tests, and kidney function tests, may be performed to monitor your progress or to check for side effects, especially if you are taking this medicine for a long period of time. KEEP ALL DOCTOR AND LABORATORY APPOINTMENTS while you are taking this medicine. DO NOT DRIVE, OPERATE MACHINERY, OR DO ANYTHING ELSE THAT COULD BE DANGEROUS until you know how you react to this medicine. ALCOHOL WARNING: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take this medicine or other pain relievers/fever reducers. BEFORE YOU BEGIN TAKING ANY NEW MEDICINE, either prescription or over-the-counter, check with your doctor or pharmacist. If you are also taking aspirin, as prescribed by your doctor for reasons such as heart attack or stroke prevention (usually these dosages are 81-325 mg per day), continue to take the aspirin and consult your doctor or pharmacist before using this medicine. CAUTION IS ADVISED WHEN USING THIS MEDICINE IN THE ELDERLY because they may be more sensitive to the effects of this medicine, especially the risk of stomach or bowel effects (such as bleeding or ulcers), or kidney effects. FOR WOMEN: USE OF THIS MEDICINE DURING PREGNANCY has resulted in fetal and newborn death. If you think you may be pregnant, contact your doctor immediately. THIS MEDICINE IS EXCRETED IN BREAST MILK. IF YOU ARE OR WILL BE BREAST-FEEDING while you are using this medicine, check with your doctor or pharmacist to discuss the risks to your baby.

**POSSIBLE SIDE EFFECTS:** SIDE EFFECTS, that may go away during treatment, include nausea, vomiting, diarrhea, gas, constipation, indigestion, dizziness, lightheadedness, drowsiness, or headache. If they continue or are bothersome, check with your doctor. CHECK WITH YOUR DOCTOR AS SOON AS POSSIBLE if you experience ringing in ears. CONTACT YOUR DOCTOR IMMEDIATELY if you experience rapid or pounding heartbeat; easy bruising or bleeding; very stiff neck; or mental/mood changes. CONTACT YOUR DOCTOR IMMEDIATELY if you experience sharp or crushing chest pain; sudden shortness of breath; sudden leg pain; sudden severe headache, vomiting, dizziness, or fainting; changes in vision; numbness of an arm or leg; slurred speech; one-sided weakness; sudden unexplained weight gain; change in amount of urine produced; severe or persistent stomach pain; vomit that looks like coffee grounds; black tarry stools; itching, reddened, swollen, blistered, painful, or peeling skin; yellowing of the skin or eyes; dark urine; right-sided tenderness; severe or persistent tiredness; fever, chills, or sore throat; severe or persistent nausea; swelling of hands, ankles, feet, face, lips, eyes, throat, or tongue; difficulty swallowing or breathing; or hoarseness. AN ALLERGIC REACTION TO THIS MEDICINE is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash, itching, swelling, severe dizziness, or trouble breathing. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms of overdose may include severe stomach pain, coffee ground-like vomit, unusually fast or slow heartbeat, trouble breathing, extreme drowsiness, loss of consciousness, and seizures.

**ADDITIONAL INFORMATION:** DO NOT SHARE THIS MEDICINE with others for whom it was not prescribed. DO NOT USE THIS MEDICINE for other health conditions. KEEP THIS MEDICINE out of the reach of children and pets. IF USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME, obtain refills before your supply runs out.

**KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.**

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)417-7627

**RX # 0503783-09015**

DATE: 03/26/13

**IBUPROFEN 600MG TABLETS**

QTY: 30 NO REFILLS - DR. AUTH REQUIRED

Refill NDC: 53746-0465-05

Retail Price: \$11.99 Your Insurance Saved You: \$ 8.35

**\$ 3.64**

W. CHRISTENSEN, MD  
MFG: AMNEAL  
SDB/ADO/ADO/ /ADO

PLAN: WPB  
GROUP#: 01960917  
CLAIM REF# 130854405472081999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Customer  
Receipt

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)417-7627

**RX # 0503783-09015**

DATE: 03/26/13

**IBUPROFEN 600MG TABLETS**

QTY: 30 NO REFILLS - DR. AUTH REQUIRED

Refill NDC: 53746-0465-05

Retail Price: \$11.99 Your Insurance Saved You: \$ 8.35

**\$ 3.64**

W. CHRISTENSEN, MD  
MFG: AMNEAL  
SDB/ADO/ADO/ /ADO

PLAN: WPB  
GROUP#: 01960917  
CLAIM REF# 130854405472081999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Duplicate  
Receipt

Pharmacy use only

TUE 3:44PM

Refill


IBUPROFEN 600MG TABLETS

53746-0465-05

FAST RACK

QTY 30

10 DRAM



WHITE  
FRONT: IP 465

SDB/ADO/ADO/ /ADO

Med Guide

Do not flush unused medications or pour down a sink or drain.

WIC# 216382



**INGREDIENT NAME:** IBUPROFEN (eye-byoo-PROE-fen)

**COMMON USES:** This medicine is a nonsteroidal anti-inflammatory drug (NSAID) used to treat mild to moderate pain, osteoarthritis, and rheumatoid arthritis. It may also be used to treat other conditions as determined by your doctor.

**BEFORE USING THIS MEDICINE:** WARNING: THE RISK OF SERIOUS AND SOMETIMES FATAL HEART PROBLEMS, HEART ATTACK, AND STROKE may be increased with the use of this medicine. This risk may be increased the longer you use this medicine. Risk may also be higher in patients who have heart problems or who are at risk for heart problems. THIS MEDICINE SHOULD NOT BE USED to treat pain before or after coronary artery heart bypass (CABG) surgery. THE RISK OF SERIOUS AND SOMETIMES FATAL STOMACH AND BOWEL PROBLEMS, including bleeding, ulcers, and holes in the stomach and bowel, is increased while using this medicine. These problems may occur at any time during therapy, with or without symptoms. Elderly patients are at higher risk for serious stomach problems. Ask your doctor or pharmacist for more information about this medicine and its side effects. Some medicines or medical conditions may interact with this medicine. INFORM YOUR DOCTOR OR PHARMACIST of all prescription and over-the-counter medicine that you are taking. DO NOT TAKE THIS MEDICINE if you are also taking heparins or tacrolimus. ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION may be needed if you are taking serotonin reuptake blocker medicines such as fluoxetine or citalopram, "blood thinners" such as warfarin, bisphosphonates such as alendronate or risedronate, cyclosporine, corticosteroids such as prednisone, high blood pressure medicines (including ACE inhibitors such as captopril, angiotensin II receptor antagonists such as losartan, and beta-blockers such as metoprolol), "water pills" (diuretics such as furosemide, hydrochlorothiazide, triamterene), lithium, methotrexate, or aspirin. DO NOT START OR STOP ANY MEDICINE without doctor or pharmacist approval. Inform your doctor of any other medical conditions including poorly controlled diabetes, dehydration, heart problems (such as heart failure or history of heart attack), swelling of the hands, feet, or ankles (edema), high blood pressure, history of stroke, blood clotting problems, stomach or bowel problems (such as bleeding or ulcers), history of tobacco use or alcohol use, kidney problems, liver problems, blood or bleeding problems (such as anemia), asthma, growths in the nose (nasal polyps), any allergies (especially history of angioedema with symptoms of lip, tongue, throat swelling), pregnancy, or breast-feeding. USE OF THIS MEDICINE IS NOT RECOMMENDED if you have a history of allergy to aspirin or other NSAIDs (e.g., naproxen, celecoxib). USE OF THIS MEDICINE IS NOT RECOMMENDED if you have history of severe kidney disease or if you are going to have or have recently had coronary artery heart bypass (CABG) surgery. Contact your doctor or pharmacist if you have any questions or concerns about taking this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for using this medicine provided by your doctor. This medicine may come with a medication guide. Read it carefully. Ask your doctor, nurse, or pharmacist any questions that you may have about this medicine. TAKE THIS MEDICINE with a full glass (8 oz./240 ml) of water. DO NOT lie down for 30 minutes after taking this medicine. The dosage is based on your medical condition and response to therapy. If repeat doses are needed, they are usually given 6 or 8 hours apart, or as directed by your doctor. When used in children, the dose is based on your child's weight. Read the product instructions to find the appropriate dose for your child's weight. Consult the pharmacist or doctor if you have questions or if you need help in choosing the appropriate dosage form. THIS MEDICINE MAY BE TAKEN WITH FOOD if it upsets your stomach. Taking it with food may not decrease the risk of stomach or bowel problems (such as bleeding or ulcers) that may occur while taking this medicine. Talk with your doctor or pharmacist if you experience persistent stomach upset. STORE THIS MEDICINE at room temperature, away from heat and light. Do not store in the bathroom. IF YOU MISS A DOSE OF THIS MEDICINE, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.

**CAUTIONS:** THIS MEDICINE INCREASES YOUR RISK OF SERIOUS STOMACH OR BOWEL PROBLEMS (such as ulcers and bleeding). This risk is increased if you are elderly or are in poor health, if you have a history of smoking or drinking alcohol, if you take corticosteroid medicines (such as prednisone) or "blood thinners" (such as warfarin), or if you take this medicine for a long period of time. THIS MEDICINE MAY ALSO INCREASE YOUR RISK for certain serious heart and blood vessel problems (such as heart attack and stroke). TAKE THIS MEDICINE EXACTLY AS PRESCRIBED BY YOUR DOCTOR, at the lowest possible dose for the

shortest time needed. Talk with your doctor or pharmacist for further information. DO NOT TAKE THIS MEDICINE IF YOU HAVE HAD A SEVERE ALLERGIC REACTION to aspirin or any medicine containing aspirin or to a nonsteroidal anti-inflammatory drug (such as Feldene, Motrin, Naprosyn, Clinoril). A severe reaction includes a severe rash, hives, breathing difficulties, or dizziness. If you have a question about whether you are allergic to this medicine or if a certain medicine is a nonsteroidal anti-inflammatory drug, contact your doctor or pharmacist. DO NOT EXCEED THE RECOMMENDED DOSE or take this medicine for longer than 10 days for pain or 3 days for fever, unless directed by your doctor. Laboratory and/or medical tests, including blood counts, liver function tests, and kidney function tests, may be performed to monitor your progress or to check for side effects, especially if you are taking this medicine for a long period of time. KEEP ALL DOCTOR AND LABORATORY APPOINTMENTS while you are taking this medicine. DO NOT DRIVE, OPERATE MACHINERY, OR DO ANYTHING ELSE THAT COULD BE DANGEROUS until you know how you react to this medicine. ALCOHOL WARNING: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take this medicine or other pain relievers/fever reducers. BEFORE YOU BEGIN TAKING ANY NEW MEDICINE, either prescription or over-the-counter, check with your doctor or pharmacist. If you are also taking aspirin, as prescribed by your doctor for reasons such as heart attack or stroke prevention (usually these dosages are 81-325 mg per day), continue to take the aspirin and consult your doctor or pharmacist before using this medicine. CAUTION IS ADVISED WHEN USING THIS MEDICINE IN THE ELDERLY because they may be more sensitive to the effects of this medicine, especially the risk of stomach or bowel effects (such as bleeding or ulcers), or kidney effects. FOR WOMEN: USE OF THIS MEDICINE DURING PREGNANCY has resulted in fetal and newborn death. If you think you may be pregnant, contact your doctor immediately. THIS MEDICINE IS EXCRETED IN BREAST MILK. IF YOU ARE OR WILL BE BREAST-FEEDING while you are using this medicine, check with your doctor or pharmacist to discuss the risks to your baby.

**POSSIBLE SIDE EFFECTS:** SIDE EFFECTS, that may go away during treatment, include nausea, vomiting, diarrhea, gas, constipation, indigestion, dizziness, lightheadedness, drowsiness, or headache. If they continue or are bothersome, check with your doctor. CHECK WITH YOUR DOCTOR AS SOON AS POSSIBLE if you experience ringing in ears. CONTACT YOUR DOCTOR IMMEDIATELY if you experience rapid or pounding heartbeat; easy bruising or bleeding; very stiff neck; or mental/mood changes. CONTACT YOUR DOCTOR IMMEDIATELY if you experience sharp or crushing chest pain; sudden shortness of breath; sudden leg pain; sudden severe headache, vomiting, dizziness, or fainting; changes in vision; numbness of an arm or leg; slurred speech; one-sided weakness; sudden unexplained weight gain; change in amount of urine produced; severe or persistent stomach pain; vomit that looks like coffee grounds; black tarry stools; itching, reddened, swollen, blistered, painful, or peeling skin; yellowing of the skin or eyes; dark urine; right-sided tenderness; severe or persistent tiredness; fever, chills, or sore throat; severe or persistent nausea; swelling of hands, ankles, feet, face, lips, eyes, throat, or tongue; difficulty swallowing or breathing; or hoarseness. AN ALLERGIC REACTION TO THIS MEDICINE is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash, itching, swelling, severe dizziness, or trouble breathing. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms of overdose may include severe stomach pain, coffee ground-like vomit, unusually fast or slow heartbeat, trouble breathing, extreme drowsiness, loss of consciousness, and seizures.

**ADDITIONAL INFORMATION:** DO NOT SHARE THIS MEDICINE with others for whom it was not prescribed. DO NOT USE THIS MEDICINE for other health conditions. KEEP THIS MEDICINE out of the reach of children and pets. IF USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME, obtain refills before your supply runs out.

**KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.**

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)417-7627

**RX # 0513884-09015**

DATE: 04/20/13

**IBUPROFEN 600MG TABLETS**

QTY: 30 3+ REFILLS BEFORE 04/07/14

Copy NDC: 53746-0465-05

Retail Price: \$11.99 Your Insurance Saved You: \$ 8.35

**\$ 3.64**

W. CHRISTENSEN, MD  
MFG: AMNEAL  
CAM/HYK/HYK/AMT/AMT

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 131104699921091998

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Customer  
Receipt

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)417-7627

**RX # 0513884-09015**

DATE: 04/20/13

**IBUPROFEN 600MG TABLETS**

QTY: 30 3+ REFILLS BEFORE 04/07/14

Copy NDC: 53746-0465-05

Retail Price: \$11.99 Your Insurance Saved You: \$ 8.35

**\$ 3.64**

W. CHRISTENSEN, MD  
MFG: AMNEAL  
CAM/HYK/HYK/AMT/AMT

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 131104699921091998

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951


Duplicate  
Receipt

Pharmacy use only

SAT 4:33PM  
Copy

IBUPROFEN 600MG TABLETS  
53746-0465-05  
FAST RACK

QTY 30  
10 DRAM



WHITE  
FRONT: IP 465

CAM/HYK/HYK/AMT/AMT

Med Guide

Call your doctor for medical advice about side effects.  
You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 216382



The chance of a person getting an ulcer or bleeding increases with:

- taking medicines called "corticosteroids" and "anticoagulants"
- longer use
- smoking
- drinking alcohol
- older age
- having poor health

NSAID medicines should only be used:

- exactly as prescribed
- at the lowest dose possible for your treatment
- for the shortest time needed

#### What are Non-Steroidal Anti-Inflammatory Drugs (NSAIDs)?

NSAID medicines are used to treat pain and redness, swelling, and heat (inflammation) from medical conditions such as:

- different types of arthritis
- menstrual cramps and other types of short-term pain

Who should not take a Non-Steroidal Anti-Inflammatory Drug (NSAID)?

Do not take an NSAID medicine:

- if you had an asthma attack, hives, or other allergic reaction with aspirin or any other NSAID medicine
- for pain right before or after heart bypass surgery

Tell your healthcare provider:

- about all of your medical conditions.
- about all of the medicines you take. NSAIDs and some other medicines can interact with each other and cause serious side effects. **Keep a list of your medicines to show to your healthcare provider and pharmacist.**
- if you are pregnant. **NSAID medicines should not be used by pregnant women late in their pregnancy.**
- if you are breastfeeding. **Talk to your doctor.**

What are the possible side effects of Non-Steroidal Anti-Inflammatory Drugs (NSAIDs)?

Serious side effects include:	Other side effects include:
<ul style="list-style-type: none"><li>• heart attack</li><li>• stroke</li><li>• high blood pressure</li><li>• heart failure from body swelling (fluid retention)</li><li>• kidney problems including kidney failure</li><li>• bleeding and ulcers in the stomach and intestine</li><li>• low red blood cells (anemia)</li><li>• life-threatening skin reactions</li><li>• life-threatening allergic reactions</li><li>• liver problems including liver failure</li><li>• asthma attacks in people who have asthma</li></ul>	<ul style="list-style-type: none"><li>• stomach pain</li><li>• constipation</li><li>• diarrhea</li><li>• gas</li><li>• heartburn</li><li>• nausea</li><li>• vomiting</li><li>• dizziness</li></ul>

Can your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

#### Other information about Non-Steroidal Anti-Inflammatory Drugs (NSAIDs)

- Aspirin is an NSAID medicine but it does not increase the chance of a heart attack. Aspirin can cause bleeding in the brain, stomach, and intestines. Aspirin can also cause ulcers in the stomach and intestines.
- Some of these NSAID medicines are sold in lower doses without a prescription (over-the-counter). Talk to your healthcare provider before using over-the-counter NSAIDs for more than 10 days.

#### NSAID medicines that need a prescription

Generic Name	Tradename
Celecoxib	Celebrex
Diclofenac	Cataflam, Voltaren, Arthrotec (combined with misoprostol),
Diflunisal	Dolobid
Etodolac	Lodine, Lodine XL
Fenoprofen	Nalfon, Nalfon 200
Flurbiprofen	Ansaid
Ibuprofen	Motrin, Tab-Profen, Vicoprofen* (combined with hydrocodone), Combunox (combined with oxycodone)
Indomethacin	Indocin, Indocin SR, Indo-Lemmon, Indomethagan
Ketoprofen	Oruvail
Ketorolac	Toradol
Mefenamic Acid	Ponstel
Meloxicam	Mobic
Nabumetone	Relafen
Naproxen	Naprosyn, Anaprox, Anaprox DS, EC-Naprosyn, Naprelan, Naprapac (copackaged with lansoprazole)
Oxaprozin	Daypro
Piroxicam	Feldene
Sulindac	Clinoril
Tolmetin	Tolectin, Tolectin DS, Tolectin 600

\* Vicoprofen contains the same dose of ibuprofen as over-the-counter (OTC) NSAIDs, and is usually used for less than 10 days to treat pain. The OTC NSAID label warns that long term continuous use may increase the risk of heart attack or stroke.

*This Medication Guide has been approved by the U.S. Food and Drug Administration.*

Revised April 2012

ROBERT A PLOCK

6827 Latta Pkwy, Dallas, TX 75227  
(214)417-7627

RX # 0510864-09015

DATE: 04/07/13

TRAMADOL 50MG TABLETS

QTY: 180 3 REFILLS BEFORE 04/07/14

New NDC: 68382-0319-10

Retail Price: \$77.97 Your Insurance Saved You: \$67.97

\$ 10.00

W. CHRISTENSEN, MD  
MFG: ZYDUS  
CAM/HYK/HYK/HYK/HYK

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 130982407165089999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Customer  
Receipt

ROBERT A PLOCK

6827 Latta Pkwy, Dallas, TX 75227  
(214)417-7627

RX # 0510864-09015

DATE: 04/07/13

TRAMADOL 50MG TABLETS

QTY: 180 3 REFILLS BEFORE 04/07/14

New NDC: 68382-0319-10

Retail Price: \$77.97 Your Insurance Saved You: \$67.97

\$ 10.00

W. CHRISTENSEN, MD  
MFG: ZYDUS  
CAM/HYK/HYK/HYK/HYK

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 130982407165089999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951


Duplicate  
Receipt

Pharmacy use only

MON 8:00AM  
New

TRAMADOL 50MG TABLETS  
68382-0319-10  
FAST RACK

QTY 180  
20 DRAM



WHITE  
FRONT: 319

CAM/HYK/HYK/HYK/HYK



Call your doctor for medical advice about side effects.  
You may report side effects to FDA at 1-800-FDA-1088.

#### INGREDIENT NAME: TRAMADOL (TRAM-a-dol)

**COMMON USES:** This medicine is an analgesic used to treat moderate to moderately severe pain.

**BEFORE USING THIS MEDICINE:** Some medicines or medical conditions may interact with this medicine. **INFORM YOUR DOCTOR OR PHARMACIST** of all prescription and over-the-counter medicine that you are taking. **DO NOT TAKE THIS MEDICINE** if you are taking sodium oxybate (GHB), or another product that contains tramadol. **ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION** may be needed if you are taking alpha-2 receptor blockers (eg, yohimbine), anorexics (eg, phentermine), anticoagulants (eg, warfarin), azole antifungals (eg, ketoconazole), butyrophenones (eg, haloperidol), carbamazepine, cyclobenzaprine, digoxin, fentanyl, linezolid, lithium, loxapine, macrolide antibiotics (eg, erythromycin), certain medicines for mental or mood disorders (eg, olanzapine), molidone, monoamine oxidase inhibitors (MAOIs) (eg, phenelzine, selegiline), nefazodone, opioid pain medicine (eg, codeine, hydrocodone), phenothiazines (eg, promethazine), quinine, rifampin, serotonin-norepinephrine reuptake inhibitors (SNRIs) (eg, duloxetine), selective serotonin reuptake inhibitors (SSRIs) (eg, fluoxetine, paroxetine), sleeping medicines (eg, zolpidem), St. John's wort, thioxanthenes (eg, thiothixene), tiagabine, tricyclic antidepressants (eg, amitriptyline), or "triptans" (eg, sumatriptan). **DO NOT START OR STOP** any medicine without doctor or pharmacist approval. Inform your doctor of any other medical conditions, including liver, kidney, or thyroid problems; blockage in the bladder; diabetes; heart problems (eg, cor pulmonale); high blood pressure; pancreas problems; prostate problems; metabolism problems; allergies; pregnancy; or breast-feeding. Tell your doctor if you have or recently have had any head injury, brain injury or tumor, increased pressure in the brain, or infection of the brain or nervous system. Tell your doctor if you have a history of lung or breathing problems (eg, asthma, chronic obstructive pulmonary disease / COPD); seizures (eg, epilepsy); recent stomach or bowel surgery; other stomach or bowel problems (eg, pain, inflammation, ulcers); alcohol or other substance abuse; mood or mental problems (eg, depression); or suicidal thoughts or actions. Tell your doctor if you drink alcohol or if you are going through withdrawal from alcohol or other substances. **USE OF THIS MEDICINE IS NOT RECOMMENDED** if you have severe or persistent diarrhea caused by antibiotic use; suicidal thoughts or actions; or a history of alcohol or other substance abuse or addiction. **USE OF THIS MEDICINE IS NOT RECOMMENDED** if you are intoxicated with alcohol, opioids or narcotics (eg, codeine, morphine), or sedatives or sleeping medicines (eg, temazepam, zolpidem). This medicine should not be used in CHILDREN younger than 16 years; safety and effectiveness in these children have not been confirmed. Contact your doctor or pharmacist if you have any questions or concerns about using this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for taking this medicine provided by your doctor. **TAKE THIS MEDICINE** by mouth with or without food. **STORE THIS MEDICINE** at 77 degrees F (25 degrees C) away from heat, moisture, and light. Do not store in the bathroom. Brief storage at temperatures between 59 and 86 degrees F (15 and 30 degrees C) is permitted. **KEEP THIS MEDICINE** out of the reach of children and away from pets. **IF YOU MISS A DOSE OF THIS MEDICINE** and you are taking it regularly, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. **DO NOT** take 2 doses at once.

**CAUTIONS:** **DO NOT TAKE THIS MEDICINE** if you have had an allergic reaction to it or if you are allergic to any ingredient in this product. **DO NOT TAKE THIS MEDICINE** if you have had a severe allergic reaction to codeine or another opioid (eg, morphine). **DO NOT EXCEED THE RECOMMENDED DOSE** or take this medicine for longer than prescribed without checking with your doctor. Exceeding the recommended dose or taking this medicine for longer than prescribed may be habit-forming or may increase the risk of seizures or breathing problems. If using this medicine for an extended period of time, **DO NOT SUDDENLY STOP** taking this medicine without your doctor's approval. Your dose may need to be slowly lowered to avoid side effects. When used for an extended period of time, this medicine may not work as well and may require different dosing. Talk with your doctor if this medicine stops working well. **DO NOT TAKE MORE THAN PRESCRIBED.** **BEFORE YOU HAVE ANY MEDICAL OR DENTAL TREATMENTS, EMERGENCY CARE, OR SURGERY,** tell the doctor or dentist that you are using this medicine. **DO NOT DRINK ALCOHOL** while you are using this medicine. **CHECK WITH YOUR DOCTOR** before you use medicines that may cause drowsiness (eg, sleep aids, muscle relaxers, narcotic pain medicines) while you are using this medicine; it may add to their effects. Ask your pharmacist if you

have questions about which medicines may cause drowsiness. This medicine may cause drowsiness or dizziness. **DO NOT DRIVE, OPERATE MACHINERY, OR DO ANYTHING ELSE THAT COULD BE DANGEROUS** until you know how you react to this medicine. Using this medicine alone, with other medicines, or with alcohol may lessen your ability to drive or to perform other potentially dangerous tasks. Alcohol, hot weather, exercise, or fever may increase dizziness. **IF DIZZINESS OCCURS,** sit up or stand slowly, especially in the morning. Sit or lie down at the first sign of this effect. **THIS MEDICINE MAY INCREASE YOUR RISK OF SEIZURES.** Your risk may be greater if you also have certain medical conditions, use certain medicines, or if you use a lot of alcohol. Talk to your doctor to see if you may have a greater risk of seizures while taking this medicine. **SEROTONIN SYNDROME** is a possibly fatal syndrome that can be caused by this medicine. Your risk may be greater if you take this medicine with certain other medicines (eg, "triptans," MAOIs, antidepressants). Symptoms may include agitation; confusion; hallucinations; coma; fever; fast or irregular heartbeat; tremor; excessive sweating; and nausea, vomiting, or diarrhea. Contact your doctor at once if you have any of these symptoms. **BEFORE YOU BEGIN TAKING ANY NEW MEDICINES,** either prescription or over-the-counter, check with your doctor or pharmacist. **CAUTION IS ADVISED WHEN USING THIS MEDICINE IN THE ELDERLY** because they may be more sensitive to the effects of this medicine. **FOR WOMEN:** This medicine may cause harm to the fetus. **IF YOU THINK YOU MAY BE PREGNANT,** discuss with your doctor the benefits and risks of using this medicine during pregnancy. **THIS MEDICINE IS EXCRETED** in breast milk. **DO NOT BREAST-FEED** while taking this medicine.

**POSSIBLE SIDE EFFECTS:** **SIDE EFFECTS** that may occur while taking this medicine include constipation; diarrhea; dizziness; drowsiness; dry mouth; headache; indigestion; mildly increased sweating; mild itching; nausea; trouble sleeping; vomiting; weakness. If they continue or are bothersome, check with your doctor. **CONTACT YOUR DOCTOR IMMEDIATELY** if you experience agitation; chest pain; confusion; difficult or painful urination; excessive sweating; fainting; fast or irregular heartbeat; fever; hallucinations; loss of coordination; mood or mental changes (eg, depression); red, blistered, swollen, or peeling skin; seizures; severe dizziness or lightheadedness; severe nausea, vomiting, or diarrhea; severe or persistent headache; slow or shallow breathing; suicidal thoughts or behaviors; tremor; or vision problems. **AN ALLERGIC REACTION** to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, or tongue; unusual hoarseness. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your health care provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** **IF OVERDOSE IS SUSPECTED,** contact your local poison control center or emergency room immediately. Symptoms may include bluish skin; cold, clammy skin; difficult, shallow, or slow breathing; drowsiness leading to unresponsiveness or coma; excessive sweating; limp muscles; pinpoint pupils; seizures; slow or irregular heartbeat.

**ADDITIONAL INFORMATION:** **DO NOT SHARE THIS MEDICINE** with others for whom it was not prescribed. **DO NOT USE THIS MEDICINE** for other health conditions. **IF YOU WILL BE USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME,** be sure to obtain necessary refills before your supply runs out. **CHECK WITH YOUR PHARMACIST** about how to dispose of unused medicine.

**KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.**

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)417-7627

**RX # 0508875-09015**

DATE: 03/28/13

#### TRAMADOL 50MG TABLETS

QTY: 90 NO REFILLS - DR. AUTH REQUIRED

Copy NDC: 68382-0319-10

Retail Price: \$46.49 Your Insurance Saved You: \$36.49

**\$ 10.00**

W. CHRISTENSEN, MD  
MFG: ZYDUS  
SDB/HYK/HYK/ /NJT

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 130874892037062999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Customer  
Receipt

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)417-7627

**RX # 0508875-09015**

DATE: 03/28/13

#### TRAMADOL 50MG TABLETS

QTY: 90 NO REFILLS - DR. AUTH REQUIRED

Copy NDC: 68382-0319-10

Retail Price: \$46.49 Your Insurance Saved You: \$36.49

**\$ 10.00**

W. CHRISTENSEN, MD  
MFG: ZYDUS  
SDB/HYK/HYK/ /NJT

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 130874892037062999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Duplicate  
Receipt

Pharmacy use only

THU 5:04PM  
Copy

TRAMADOL 50MG TABLETS  
68382-0319-10  
FAST RACK

QTY 90  
10 DRAM



WHITE  
FRONT: 319

SDB/HYK/HYK/ /NJT

Do not flush unused medications or pour down a sink or drain.

WIC# 216382



Call your doctor for medical advice about side effects.  
You may report side effects to FDA at 1-800-FDA-1088.

**INGREDIENT NAME:** BETAMETHASONE  
(bay-ta-METH-a-sone)

**COMMON USES:** This medicine is a corticosteroid used to reduce itching, redness, and swelling associated with many skin conditions. It may also be used to treat other conditions as determined by your doctor.

**BEFORE USING THIS MEDICINE:** INFORM YOUR DOCTOR OR PHARMACIST of all prescription and over-the-counter medicine that you are taking. **ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION** may be needed if you are taking prednisone (or similar medicines). Inform your doctor of any other medical conditions, allergies, pregnancy, or breast-feeding.

**HOW TO USE THIS MEDICINE:** Follow the directions for using this medicine provided by your doctor. **TO USE THIS MEDICINE:** apply a small amount of medicine to the affected area. Gently rub the medicine in until it is evenly distributed. Wash your hands after applying this medicine, unless your hands are part of treated area. **DO NOT BANDAGE OR WRAP** the affected area unless directed otherwise by your doctor. **STORE THIS MEDICINE** at room temperature, away from heat and light. **IF YOU MISS A DOSE OF THIS MEDICINE**, apply it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule.

**CAUTIONS:** IF YOUR SYMPTOMS DO NOT IMPROVE WITHIN A FEW DAYS, or if they become worse, check with your doctor. **DO NOT USE THIS MEDICINE** longer than the time prescribed by your doctor. **DO NOT USE THIS MEDICINE** for other skin conditions at a later time. **DO NOT GET THIS MEDICINE** in your eyes. **FOR WOMEN: IF YOU PLAN ON BECOMING PREGNANT**, discuss with your doctor the benefits and risks of using this medicine during pregnancy. **IT IS UNKNOWN IF THIS MEDICINE IS EXCRETED** in breast milk. **IF YOU ARE OR WILL BE BREAST-FEEDING** while you are using this medicine, check with your doctor or pharmacist to discuss the risks to your baby.

**POSSIBLE SIDE EFFECTS:** CHECK WITH

**YOUR DOCTOR AS SOON AS POSSIBLE** if you experience skin thinning and discoloration, stinging, rash, dry skin, itching, burning, redness, or swelling not present before using this medicine. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms may include muscle weakness; severe or persistent headache; symptoms of high blood sugar (eg, increased thirst, hunger, or urination; confusion; drowsiness; flushing; rapid breathing; fruit-like breath odor); unusual weight gain, especially in the face; or vision changes.

**ADDITIONAL INFORMATION:** If your symptoms do not improve within 2 weeks or if they become worse, check with your doctor. **DO NOT SHARE THIS MEDICINE** with others for whom it was not prescribed. **DO NOT USE THIS MEDICINE** for other health conditions. **KEEP THIS MEDICINE** out of the reach of children and pets. **CHECK WITH YOUR PHARMACIST** about how to dispose of unused medicine.

**KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.**

Do not flush unused medications or pour down a sink or drain.

WIC# 216382

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)417-7627

**RX # 0510823-09015**

DATE: 04/07/13

**BETAMETHASONE DIP AUG CREAM 50GM**

QTY: 50 1 REFILL BEFORE 04/07/14

New NDC: 68462-0290-52

Retail Price: \$89.19 Your Insurance Saved You: \$79.19

\$ 10.00

W. CHRISTENSEN, MD  
MFG: GLENMARK  
CAM/MMC/MMC/ /HYK

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 130973832570025998

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Customer  
Receipt

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)417-7627

**RX # 0510823-09015**

DATE: 04/07/13

**BETAMETHASONE DIP AUG CREAM 50GM**

QTY: 50 1 REFILL BEFORE 04/07/14

New NDC: 68462-0290-52

Retail Price: \$89.19 Your Insurance Saved You: \$79.19

\$ 10.00

W. CHRISTENSEN, MD  
MFG: GLENMARK  
CAM/MMC/MMC/ /HYK

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 130973832570025998

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Duplicate  
Receipt

Pharmacy use only

MON 8:00AM  
New

BETAMETHASONE DIP AUG CREAM 50GM  
68462-0290-52  
ALPHA

QTY 50

CAM/MMC/MMC/ /HYK



**INGREDIENT NAME:** HYDROCODONE (hye-droe-KO-done) and ACETAMINOPHEN (a-seat-a-MIN-oh-fen)

**COMMON USES:** This medicine is a narcotic and analgesic combination used to relieve moderate to moderately severe pain. It may also be used to treat other conditions as determined by your doctor.

**BEFORE USING THIS MEDICINE:** **WARNING:** THIS MEDICINE CONTAINS ACETAMINOPHEN. Severe and sometimes fatal liver problems, including the need for liver transplant, have been reported with the use of acetaminophen. Most cases of these liver problems occurred in patients taking excessive doses of acetaminophen (more than 4,000 mg per day). Also, patients who developed these liver problems were often using more than 1 medicine that contained acetaminophen. Discuss any questions or concerns with your doctor. Some medicines or medical conditions may interact with this medicine. **INFORM YOUR DOCTOR OR PHARMACIST** of all prescription and over-the-counter medicine that you are taking. **DO NOT TAKE THIS MEDICINE** if you are also taking sodium oxybate (XGB), **ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION may be needed** if you are taking anticonvulsants (eg, valproic acid, phenytoin, carbamazepine, lamotrigine, phenobarbital, ethosuximide, clobazam, zonisamide, levetiracetam, topiramate, gabapentin, pregabalin, tiagabine, vigabatrin, perampanel, sleep medicines (eg, zolpidem), tricyclic antidepressants (eg, amitriptyline), or medicines that may harm the liver (eg, methotrexate, ketoconazole, certain medicines for HIV infection). **Ask your doctor** if you are unsure if any of your medicines might harm the liver. **DO NOT START OR STOP** any medicine without doctor or pharmacist approval. Inform your doctor of any other medical conditions, including a history of liver problems (eg, hepatitis), kidney problems, lung or breathing problems (eg, asthma, chronic obstructive pulmonary disease / COPD), drug or alcohol abuse or dependence, heart problems (eg, coronary artery disease, heart failure), low blood pressure, stomach or bowel problems (eg, blockage, inflammation, paralysis), gallbladder problems, prostate problems (eg, an enlarged prostate), difficulty urinating, seizures, thyroid problems, adrenal gland problems (eg, Addison disease), mental or mood problems, inflammation of the pancreas (pancreatitis), curvature of the spine (scoliosis), suicidal thoughts or actions, allergies, pregnancy, or breast-feeding. Tell your doctor if you have severe drowsiness, or have or recently have had any head injury, increased pressure in the brain, growths in the brain (eg, tumors), or infection of the brain or nervous system. Tell your doctor if you have recently had surgery (eg, stomach or bowel surgery) or are scheduled to have surgery. Tell your doctor if you are in very poor health; are very overweight or are dehydrated; have low blood volume, stomach pain, or constipation; drink alcohol; or are going through withdrawal from alcohol or other substances. **USE OF THIS MEDICINE IS NOT RECOMMENDED** if you have very slow or difficult breathing, or you are having an asthma attack. Contact your doctor or pharmacist if you have any questions or concerns about taking this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for using this medicine provided by your doctor. **TAKE THIS MEDICINE** by mouth with or without food. If stomach upset occurs, take with food to reduce stomach irritation. **IF YOU HAVE BEEN TAKING THIS MEDICINE REGULARLY** for longer than a few weeks, **do not suddenly stop taking it without checking with your doctor.** Your doctor may need to gradually lower your dose. **SOME OF THESE PRODUCTS CONTAIN SULFITES.** Sulfites may cause an allergic reaction in some patients (eg, asthma patients). If you have ever had an allergic reaction to sulfites, ask your pharmacist if your product has sulfites in it. **STORE THIS MEDICINE** at room temperature, between 68 and 77 degrees F (20 and 25 degrees C), away from heat, moisture, and light. Do not store in the bathroom. **KEEP THIS MEDICINE** out of the reach of children and away from pets. **IF YOU MISS A DOSE OF THIS MEDICINE** and you are taking it regularly, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.

**CAUTIONS:** **DO NOT TAKE THIS MEDICINE** if you are allergic to any ingredient in this medicine. **BEFORE USING THIS MEDICINE,** check with your doctor if you have had an allergic reaction to other narcotic medicines (eg, codeine, morphine, oxycodone). **DO NOT EXCEED THE RECOMMENDED DOSE,** take more often than prescribed, or use for longer than prescribed without checking with your doctor. Exceeding the recommended dose or taking this medicine for longer than prescribed may be habit-forming. If using this medicine for an extended period of time, **DO NOT SUDDENLY STOP** taking this medicine without your doctor's approval. When using for an extended period, this medicine may not work as well and may require different dosing. Talk with your doctor if this medicine stops working well. **THIS MEDICINE MAY AFFECT CERTAIN LAB TEST RESULTS.** Make sure that all of your doctors and laboratory personnel know you are taking this medicine. **BEFORE YOU HAVE ANY MEDICAL OR DENTAL TREATMENTS, emergency care, or surgery,** tell the doctor or dentist that you are using this medicine. **THIS MEDICINE MAY CAUSE DIZZINESS, lightheadedness, or fainting.** Alcohol, hot weather, exercise, and fever can increase these effects. To prevent them, sit up or stand slowly, especially in the morning. Also sit or lie down at the first sign of dizziness, lightheadedness, or weakness. **THIS MEDICINE MAY ALSO CAUSE DROWSINESS OR BLURRED VISION.** Do not drive, operate machinery, or do anything else that could be dangerous until you know how you react to this medicine. Using this medicine alone, with other medicines, or with alcohol may lessen your ability to drive or to perform other potentially dangerous tasks. **DO NOT DRINK ALCOHOL** while you are using this medicine. **CHECK WITH YOUR DOCTOR** before you use medicines that may cause drowsiness (eg, sleep aids, muscle relaxers) while you are using this medicine; it may add to their effects. Ask your pharmacist if you have questions about which medicines may cause drowsiness. **THIS MEDICINE CONTAINS ACETAMINOPHEN.** Do not take additional acetaminophen for pain or fever without checking with your doctor or pharmacist. Ask your pharmacist if you have questions about which medicines contain acetaminophen. **THIS MEDICINE MAY HARM YOUR LIVER.** Your risk may be greater if you drink alcohol while you are using this medicine. Talk to your doctor before you take this medicine or other fever reducers if you drink alcohol. **CONTACT YOUR DOCTOR RIGHT AWAY** if you take more than 4,000 mg of acetaminophen per day, even if you feel well. **BEFORE YOU BEGIN TAKING ANY NEW MEDICINE,** either prescription or over-the-counter, check with your doctor or pharmacist. **CAUTION IS ADVISED WHEN USING THIS MEDICINE IN THE ELDERLY** because they may be more sensitive to the effects of the medicine. **FOR WOMEN:** IF YOU BECOME PREGNANT, discuss with your doctor the benefits and risks of using this medicine during pregnancy. Long-term use of this medicine during pregnancy may cause dependence in the fetus or newborn. Discuss any questions or concerns with your doctor. **THIS MEDICINE IS EXCRETED** in breast milk. **DO NOT BREAST-FEED** while taking this medicine.

**POSSIBLE SIDE EFFECTS:** **SIDE EFFECTS** that may occur while taking this medicine include dizziness, drowsiness, lightheadedness, nausea, or vomiting. If

they continue or are bothersome, check with your doctor. **THIS MEDICINE MAY CAUSE CONSTIPATION.** Talk with your doctor or pharmacist about using a stool softener or laxative to prevent constipation. It is also important to maintain a diet adequate in fiber, drink plenty of water, and exercise to prevent constipation. If you become constipated while taking this medicine, talk with your doctor or pharmacist. **CONTACT YOUR DOCTOR IMMEDIATELY** if you experience a change in the amount of urine produced; change or loss in hearing; chest pain; confusion; fainting; fever, chills, or persistent sore throat; mental or mood changes (eg, agitation, anxiety, depression); red, swollen, blistered, or peeling skin; severe or persistent constipation; severe or persistent dizziness, drowsiness, headache, or lightheadedness; shortness of breath; slow, shallow, or difficult breathing; symptoms of liver problems (eg, yellowing of the skin or eyes, pale stools, dark urine, persistent loss of appetite, unusual or severe stomach pain); trouble urinating; unusual bruising or bleeding; unusual tiredness or weakness; or vision changes. **AN ALLERGIC REACTION** to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, throat, or tongue; or unusual hoarseness. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** IF OVERDOSE IS SUSPECTED, contact your local poison control center or emergency room immediately. Symptoms of overdose may include bluish skin or nails; cold and clammy skin; coma; decreased urination; difficult, shallow, or slow breathing; fainting; limp muscles; loss of consciousness; nausea or vomiting; pinpoint pupils; seizures; severe dizziness, drowsiness, or light-headedness; slow heartbeat; symptoms of liver problems (eg, yellowing of the skin or eyes, pale stools, dark urine, loss of appetite, unusual or severe stomach pain); unusual bruising or bleeding; or unusual sweating or tiredness.

**ADDITIONAL INFORMATION:** If your symptoms do not improve or if they become worse, contact your doctor. **DO NOT SHARE THIS MEDICINE** with others for whom it was not prescribed. **DO NOT USE THIS MEDICINE** for other health conditions. **CALL WITH YOUR PHARMACIST** about how to dispose of unused medicine.

**KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.**

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

**RX # 0517526-09015**

DATE: 05/29/13

**HYDROCODONE /ACETAMINOPHEN 5-325 TB**

QTY: 90 NO REFILLS - DR. AUTH REQUIRED

Refill NDC: 00591-3202-01

Retail Price: \$65.89 Your Insurance Saved You: \$55.89

**\$ 10.00**

A. PARK, MD  
MFG: WATSON  
RTS/ADO/ADO/ADO/ADO

PLAN: WPB  
GROUP#: 01960917  
CLAIM REF# 131491548821076999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Customer  
Receipt

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

**RX # 0517526-09015**

DATE: 05/29/13

**HYDROCODONE /ACETAMINOPHEN 5-325 TB**

QTY: 90 NO REFILLS - DR. AUTH REQUIRED

Refill NDC: 00591-3202-01

Retail Price: \$65.89 Your Insurance Saved You: \$55.89

**\$ 10.00**

A. PARK, MD  
MFG: WATSON  
RTS/ADO/ADO/ADO/ADO

PLAN: WPB  
GROUP#: 01960917  
CLAIM REF# 131491548821076999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Duplicate  
Receipt

Pharmacy use only

WED 11:00AM

Refill


HYDROCODONE /ACETAMINOPHEN 5-325 TB

00591-3202-01

FAST RACK

QTY 90

20 DRAM



WHITE AND MULTICOLOR

FRONT: 3202

BACK: WATSON

RTS/ADO/ADO/ADO/ADO

Do not flush unused medications or pour down a sink or drain.

WIC# 216382

Call your doctor for medical advice about side effects.  
You may report side effects to FDA at 1-800-FDA-1088.



Call your doctor for medical advice about side effects.  
You may report side effects to FDA at 1-800-FDA-1088.

**INGREDIENT NAME:** DIAZEPAM (dye-AZ-e-pam)

**COMMON USES:** This medicine is a benzodiazepine used to treat anxiety disorders or for the short-term relief of the symptoms of anxiety. It is also used to relieve certain types of muscle spasms and to treat symptoms of alcohol withdrawal (eg, agitation, tremor, hallucinations). This medicine is also used with other medicines to treat certain types of seizure disorders. It may also be used for other conditions as determined by your doctor.

**BEFORE USING THIS MEDICINE:** Some medicines or medical conditions may interact with this medicine. **INFORM YOUR DOCTOR OR PHARMACIST** of all prescription and over-the-counter medicine that you are taking. **DO NOT TAKE THIS MEDICINE** if you are also taking an HIV protease inhibitor (eg, ritonavir) or sodium oxybate (GHB). **ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION** may be needed if you are taking rifamycins (eg, rifampin), St. John's wort, azole antifungals (eg, ketoconazole), barbiturates (eg, phenobarbital), cimetidine, clozapine, diltiazem, macrolide antibiotics (eg, erythromycin), methadone, nefazodone, omeprazole, selective serotonin reuptake inhibitors (SSRIs) (eg, fluoxetine, fluvoxamine), telithromycin, disulfiram, hormonal contraceptives (eg, birth control pills), monoamine oxidase inhibitors (MAOIs) (eg, phenelzine), narcotic pain medicines (eg, morphine, codeine), hydantoins (eg, phenytoin), or phenothiazines (eg, thioridazine) or other medicines for mood or mental problems (eg, olanzapine). **DO NOT START OR STOP** any medicine without doctor or pharmacist approval. Inform your doctor of any other medical conditions, including glaucoma or increased pressure in the eye, liver or kidney problems, muscle problems or weakness, a blood disorder (eg, porphyria), allergies, pregnancy, or breast-feeding. Tell your doctor if you have a history of lung or breathing problems (eg, asthma, chronic obstructive pulmonary disease / COPD), seizures, mood or mental problems (eg, depression, psychosis), suicidal thoughts or actions, or alcohol or other substance abuse or dependence. **USE OF THIS MEDICINE IS NOT RECOMMENDED** if you have acute narrow-angle glaucoma, untreated open-angle glaucoma, certain muscle problems (eg, myasthenia gravis), severe liver problems, severe breathing problems, or sleep apnea. Use of this medicine in children younger than 6 months old is not recommended. Discuss with your doctor the risks and benefits of giving this medicine to your child. Contact your doctor or pharmacist if you have any questions or concerns about taking this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for taking this medicine provided by your doctor. **TAKE THIS MEDICINE** by mouth with or without food. **CHECK WITH YOUR DOCTOR** before you eat grapefruit or drink grapefruit juice while you use this medicine. **STORE THIS MEDICINE** at room temperature, 59 to 86 degrees F (15 to 30 degrees C), in a tight, light-resistant container, away from heat, moisture, and light. Do not store in the bathroom. **KEEP THIS MEDICINE** out of the reach of children and away from pets. **IF YOU MISS A DOSE OF THIS MEDICINE AND YOU ARE TAKING IT REGULARLY**, take it as soon as possible. If several hours have passed or if it is nearing time for the next dose, do not double the dose to catch up, unless advised by your health care provider. **DO NOT** take 2 doses at once.

**CAUTIONS:** **DO NOT TAKE THIS MEDICINE** if you have had an allergic reaction to it or if you are allergic to any ingredient in this product. **IF YOU TAKE THIS MEDICINE REGULARLY OVER A LONGER PERIOD OF TIME** (eg, several months), do not suddenly stop it without first talking with your doctor. You may have an increased risk of side effects (eg, mood changes, trouble sleeping). **IF YOU NEED TO STOP THIS MEDICINE** or add a new medicine, your doctor may need to gradually lower your dose. **WHEN USED FOR AN EXTENDED PERIOD OF TIME**, this medicine may not work as well and may require different dosing. Talk with your doctor if this medicine stops working well. **DO NOT EXCEED THE RECOMMENDED DOSE** or take this medicine for longer than prescribed without checking with your doctor. **Exceeding the recommended dose or taking this medicine for longer than prescribed may be habit-forming.** Laboratory and/or medical tests, including liver function and blood counts, may be performed to monitor your progress or to check for side effects. **KEEP ALL DOCTOR AND LABORATORY APPOINTMENTS** while you are taking this medicine. **BEFORE YOU HAVE ANY MEDICAL OR DENTAL TREATMENTS, EMERGENCY CARE, OR SURGERY**, tell the doctor or dentist that you are using this medicine. This medicine may cause drowsiness, dizziness, or blurred vision. **DO NOT DRIVE, OPERATE MACHINERY, OR DO ANYTHING ELSE THAT COULD BE DANGEROUS** until you know how you react to this medicine. Using this medicine alone, with other medicines, or with alcohol may lessen your ability to drive or to perform other potentially dangerous tasks. **DO NOT DRINK ALCOHOL OR USE MEDICINES THAT MAY CAUSE DROWSINESS** (eg, sleep aids, muscle

relaxers) while you are using this medicine; it may add to their effects. Ask your pharmacist if you have questions about which medicines may cause drowsiness. **IF YOU ARE TAKING THIS MEDICINE FOR SEIZURES**, carry an ID card at all times that says you have seizure disorder and you take this medicine. **BEFORE YOU BEGIN TAKING ANY NEW MEDICINE**, either prescription or over-the-counter, check with your doctor or pharmacist. **CAUTION IS ADVISED WHEN USING THIS MEDICINE IN THE ELDERLY** because they may be more sensitive to the effects of the medicine, especially drowsiness, incoordination, and mental or mood changes. **CAUTION IS ADVISED WHEN USING THIS MEDICINE IN CHILDREN** because they may be more sensitive to the effects of the medicine, especially mental or mood changes. **FOR WOMEN: THIS MEDICINE MAY CAUSE HARM TO THE HUMAN FETUS.** **IF YOU THINK YOU MAY BE PREGNANT**, discuss with your doctor the benefits and risks of using this medicine during pregnancy. **THIS MEDICINE IS FOUND IN breast milk.** **DO NOT BREAST-FEED** while taking this medicine.

**POSSIBLE SIDE EFFECTS:** **SIDE EFFECTS** that may occur while taking this medicine include drowsiness; incoordination; muscle weakness; or tiredness. If they continue or are bothersome, check with your doctor. **CONTACT YOUR DOCTOR IMMEDIATELY** if you experience confusion; difficulty breathing; fainting; fever, chills, or persistent sore throat; hallucinations; inability to control urination or difficulty urinating; memory problems or memory loss; new or worsening mental or mood changes (eg, agitation, aggressiveness, behavior changes, depression, irritability, rage, restlessness); severe or persistent dizziness or lightheadedness; slurred speech; tremor; trouble sleeping; vision changes (eg, blurred vision, double vision); yellowing of the skin or eyes. **AN ALLERGIC REACTION** to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; or swelling of the mouth, face, lips, or tongue. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** **IF OVERDOSE IS SUSPECTED**, contact your local poison control center or emergency room immediately. Symptoms of overdose may include clumsiness; coma; confusion; deep sleep; fainting; loss of consciousness; severe drowsiness; slow or shallow breathing; slow reflexes.

**ADDITIONAL INFORMATION:** **DO NOT SHARE THIS MEDICINE** with others for whom it was not prescribed. **DO NOT USE THIS MEDICINE** for other health conditions. **CHECK WITH YOUR PHARMACIST** about how to dispose of unused medicine.

**KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.**

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

**RX # 0517527-09015**

DATE: 05/08/13

**DIAZEPAM 5MG TABLETS**

QTY: 2 NO REFILLS - DR. AUTH REQUIRED

New NDC: 00378-0345-05

Retail Price: \$11.99 Your Insurance Saved You: \$10.25

**\$ 1.74**

A. PARK, MD  
MFG: MYLAN  
JAC/ADO/ADO/ /NJT

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 131284539451079998

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Customer  
Receipt

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

**RX # 0517527-09015**

DATE: 05/08/13

**DIAZEPAM 5MG TABLETS**

QTY: 2 NO REFILLS - DR. AUTH REQUIRED

New NDC: 00378-0345-05

Retail Price: \$11.99 Your Insurance Saved You: \$10.25

**\$ 1.74**

A. PARK, MD  
MFG: MYLAN  
JAC/ADO/ADO/ /NJT

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 131284539451079998

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Duplicate  
Receipt

Pharmacy use only

WED 4:03PM

New

DIAZEPAM 5MG TABLETS

00378-0345-05

FAST RACK

QTY 2  
10 DRAM



ORANGE  
FRONT: MYLAN 345

JAC/ADO/ADO/ /NJT

Do not flush unused medications or pour down a sink or drain.

WIC# 216382



Call your doctor for medical advice about side effects.  
You may report side effects to FDA at 1-800-FDA-1088.

**INGREDIENT NAME:** AMOXICILLIN (a-MOX-i-SIL-in) and CLAVULANATE (KLAU-ue-la-nate)

**COMMON USES:** This medicine is a penicillin antibiotic used to treat infections caused by certain bacteria.

**BEFORE USING THIS MEDICINE:** Some medicines or medical conditions may interact with this medicine. **INFORM YOUR DOCTOR OR PHARMACIST** of all prescription and over-the-counter medicine that you are taking. **DO NOT TAKE THIS MEDICINE** if you are also taking a tetracycline antibiotic (eg, doxycycline). **ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION** may be needed if you are taking anticoagulants (eg, warfarin), chloramphenicol, hormonal birth control (eg, birth control pills), macrolide antibiotics (eg, erythromycin), methotrexate, probenecid, or sulfonamides (eg, sulfamethoxazole). **DO NOT START OR STOP** any medicine without doctor or pharmacist approval. Inform your doctor of any other medical conditions, including kidney problems, gonorrhea, allergies, pregnancy, or breast-feeding. Tell your doctor if you have a history of asthma, hay fever, hives, or liver problems or yellowing of the eyes or skin. **USE OF THIS MEDICINE IS NOT RECOMMENDED** if you have infectious mononucleosis (mono). **USE OF THIS MEDICINE IS NOT RECOMMENDED** if you have a history of liver problems or yellowing of the eyes or skin caused by this medicine. Contact your doctor or pharmacist if you have any questions or concerns about taking this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for taking this medicine provided by your doctor. Take this medicine by mouth at the start of a meal to decrease the chance of stomach upset. **STORE THIS MEDICINE** at or below 77 degrees F (25 degrees C). Store away from heat, moisture, and light. Do not store in the bathroom. **KEEP THIS MEDICINE** out of the reach of children and away from pets. **BE SURE TO USE THIS MEDICINE FOR THE FULL COURSE OF TREATMENT.** If you do not, the medicine may not clear up your infection completely. The bacteria could also become less sensitive to this or other medicines. This could make the infection harder to treat in the future. Do not miss any doses. **IF YOU MISS A DOSE OF THIS MEDICINE**, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.

**CAUTIONS:** **DO NOT TAKE THIS MEDICINE** if you have had an allergic reaction to it, to any ingredient in this product, or to another penicillin antibiotic (eg, ampicillin). **BEFORE TAKING THIS MEDICINE**, check with your doctor if you have had an allergic reaction to a cephalosporin antibiotic (eg, cephalexin) or another beta-lactam antibiotic (eg, imipenem). If you have a question about whether you are allergic to this medicine, or if a medicine is a penicillin, cephalosporin, or beta-lactam antibiotic, contact your doctor or pharmacist. **LABORATORY AND/OR MEDICAL TESTS** may be performed to monitor your progress or to check for side effects. Keep all doctor and laboratory appointments, while you are taking this medicine. **THIS MEDICINE MAY CAUSE DIZZINESS.** Do not drive, operate machinery, or do anything else that could be dangerous until you know how you react to this medicine. Using this medicine alone, along with other medicines, or with alcohol may lessen your ability to drive or to perform other potentially dangerous tasks. **MILD DIARRHEA IS COMMON WITH ANTIBIOTIC USE.** However, a more serious form of diarrhea (pseudomembranous colitis) may rarely occur. This may develop while you use the antibiotic or within several months after you stop using it. Contact your doctor right away if stomach pain or cramps, severe diarrhea, or bloody stools occur. Do not treat diarrhea without first checking with your doctor. **BROWN, YELLOW, OR GRAY TOOTH DISCOLORATION** has occurred rarely in some patients taking this medicine. It occurred most often in children. The discoloration was reduced or removed by brushing or dental cleaning in most cases. Contact your doctor if you experience this effect. **DO NOT RECEIVE A LIVE VACCINE** (eg, typhoid) while you are taking this medicine. Talk with your doctor before you receive any vaccine. **THIS MEDICINE ONLY WORKS** against bacteria; it does not treat viral infections (eg, the common cold). Long-term or repeated use of this medicine may cause a second infection. Tell your doctor if signs of a second infection occur. Your medicine may need to be changed to treat this. **BEFORE YOU BEGIN TAKING ANY NEW MEDICINE**, either prescription or over-the-counter, check with your doctor or pharmacist. **FOR WOMEN: HORMONAL BIRTH CONTROL** (eg, birth control pills) may not work as well while you are using this medicine. To prevent pregnancy, use an extra form of birth control (eg, condoms). **IF YOU BECOME PREGNANT**, discuss with your doctor the benefits and risks of using this medicine during pregnancy. **THIS MEDICINE IS EXCRETED IN BREAST MILK.** **IF YOU ARE OR WILL BE BREAST-FEEDING** while you are using this medicine, check with your doctor or pharmacist to discuss the risks to your baby. **DIABETES PATIENTS:** This medicine

may cause false test results with some urine glucose tests. Check with your doctor before you adjust the dose of your diabetes medicine or change your diet.

**POSSIBLE SIDE EFFECTS:** SIDE EFFECTS that may occur while taking this medicine include diarrhea, nausea, or vomiting. If they continue or are bothersome, check with your doctor. **CONTACT YOUR DOCTOR IMMEDIATELY** if you experience behavior changes; bloody stools; confusion; dark urine; fever, chills, or persistent sore throat; red, swollen, blistered, or peeling skin; seizures; severe or persistent diarrhea; severe stomach pain or cramps; unusual bruising or bleeding; unusual tiredness or weakness; vaginal discharge or irritation; white patches in the mouth or on the tongue; or yellowing of the skin or eyes. **AN ALLERGIC REACTION** to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; or swelling of the mouth, face, lips, or tongue. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** IF OVERDOSE IS SUSPECTED, contact your local poison control center or emergency room immediately. Symptoms may include decreased urination; severe nausea, vomiting, or diarrhea; stomach pain; or unusual drowsiness.

**ADDITIONAL INFORMATION:** If your symptoms do not improve or if they become worse, contact your doctor. **DO NOT SHARE THIS MEDICINE** with others for whom it was not prescribed. **DO NOT USE THIS MEDICINE** for other health conditions. **CHECK WITH YOUR PHARMACIST** about how to dispose of unused medicine.

**KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.**

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

**RX # 0523176-09015**

DATE: 06/06/13

**AMOX-CLAV 875MG TABLETS**

QTY: 14 NO REFILLS - DR. AUTH REQUIRED

New-E NDC:00781-1852-20

Retail Price: \$50.79 Your Insurance Saved You: \$40.79

**\$ 10.00**

W. CHRISTENSEN, MD  
MFG: GENEVA  
SDB/NJT/NJT / NJT

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 131573689802064999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Customer  
Receipt

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

**RX # 0523176-09015**

DATE: 06/06/13

**AMOX-CLAV 875MG TABLETS**

QTY: 14 NO REFILLS - DR. AUTH REQUIRED

New-E NDC:00781-1852-20

Retail Price: \$50.79 Your Insurance Saved You: \$40.79

**\$ 10.00**

W. CHRISTENSEN, MD  
MFG: GENEVA  
SDB/NJT/NJT / NJT

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 131573689802064999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Duplicate  
Receipt

Pharmacy use only

THU 1:23PM

New-E


AMOX-CLAV 875MG TABLETS

00781-1852-20

FAST RACK

QTY 14

10 DRAM



WHITE  
FRONT: GG N7

SDB/NJT/NJT / NJT

Do not flush unused medications or pour down a sink or drain.

WIC# 216382



Call your doctor for medical advice about side effects.  
You may report side effects to FDA at 1-800-FDA-1088.

**INGREDIENT NAME:** TRAMADOL (TRAM-a-dol)

**COMMON USES:** This medicine is an analgesic used to treat moderate to moderately severe pain.

**BEFORE USING THIS MEDICINE:** Some medicines or medical conditions may interact with this medicine. INFORM YOUR DOCTOR OR PHARMACIST of all prescription and over-the-counter medicine that you are taking. DO NOT TAKE THIS MEDICINE if you are taking sodium oxybate (GHB), or another product that contains tramadol. ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION may be needed if you are taking alpha-2 receptor blockers (eg, yohimbine), anorexiant (eg, phentermine), anticoagulants (eg, warfarin), azole antifungals (eg, ketoconazole), butyrophenones (eg, haloperidol), carbamazepine, cyclobenzaprine, digoxin, furazolidone, linezolid, lithium, loxapine, macrolide antibiotics (eg, erythromycin), certain medicines for mental or mood disorders (eg, olanzapine), molidone, monoamine oxidase inhibitors (MAOIs) (eg, phenelzine, selegiline), nefazodone, opioid pain medicine (eg, codeine, hydrocodone), phenothiazines (eg, promethazine), quinidine, rifampin, serotonin-norepinephrine reuptake inhibitors (SNRIs) (eg, duloxetine), selective serotonin reuptake inhibitors (SSRIs) (eg, fluoxetine, paroxetine), sleeping medicines (eg, zolpidem), St. John's wort, thioxanthenes (eg, thiothixene), tiagabine, tricyclic antidepressants (eg, amitriptyline), or "triptans" (eg, sumatriptan). DO NOT START OR STOP any medicine without doctor or pharmacist approval. Inform your doctor of any other medical conditions, including liver, kidney, or thyroid problems; blockage in the bladder; diabetes; heart problems (eg, cor pulmonale); high blood pressure; pancreas problems; prostate problems; metabolism problems; allergies; pregnancy; or breast-feeding. Tell your doctor if you have or recently have had any head injury, brain injury or tumor, increased pressure in the brain, or infection of the brain or nervous system. Tell your doctor if you have a history of lung or breathing problems (eg, asthma, chronic obstructive pulmonary disease / COPD); seizures (eg, epilepsy); recent stomach or bowel surgery; other stomach or bowel problems (eg, pain, inflammation, ulcers); alcohol or other substance abuse; mood or mental problems (eg, depression); or suicidal thoughts or actions. Tell your doctor if you drink alcohol or if you are going through withdrawal from alcohol or other substances. USE OF THIS MEDICINE IS NOT RECOMMENDED if you have severe or persistent diarrhea caused by antibiotic use; suicidal thoughts or actions; or a history of alcohol or other substance abuse or addiction. USE OF THIS MEDICINE IS NOT RECOMMENDED if you are intoxicated with alcohol, opioids or narcotics (eg, codeine, morphine), or sedatives or sleeping medicines (eg, temazepam, zolpidem). This medicine should not be used in CHILDREN younger than 16 years; safety and effectiveness in these children have not been confirmed. Contact your doctor or pharmacist if you have any questions or concerns about using this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for taking this medicine provided by your doctor. TAKE THIS MEDICINE by mouth with or without food. STORE THIS MEDICINE at 77 degrees F (25 degrees C) away from heat, moisture, and light. Do not store in the bathroom. Brief storage at temperatures between 59 and 86 degrees F (15 and 30 degrees C) is permitted. KEEP THIS MEDICINE out of the reach of children and away from pets. IF YOU MISS A DOSE OF THIS MEDICINE and you are taking it regularly, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. DO NOT take 2 doses at once.

**CAUTIONS:** DO NOT TAKE THIS MEDICINE if you have had an allergic reaction to it or if you are allergic to any ingredient in this product. DO NOT TAKE THIS MEDICINE if you have had a severe allergic reaction to codeine or another opioid (eg, morphine). DO NOT EXCEED THE RECOMMENDED DOSE or take this medicine for longer than prescribed without checking with your doctor. Exceeding the recommended dose or taking this medicine for longer than prescribed may be habit-forming or may increase the risk of seizures or breathing problems. If using this medicine for an extended period of time, DO NOT SUDDENLY STOP taking this medicine without your doctor's approval. Your dose may need to be slowly lowered to avoid side effects. When used for an extended period of time, this medicine may not work as well and may require different dosing. Talk with your doctor if this medicine stops working well. DO NOT TAKE MORE THAN PRESCRIBED. BEFORE YOU HAVE ANY MEDICAL OR DENTAL TREATMENTS, EMERGENCY CARE, OR SURGERY, tell the doctor or dentist that you are using this medicine. DO NOT DRINK ALCOHOL while you are using this medicine. CHECK WITH YOUR DOCTOR BEFORE you use medicines that may cause drowsiness (eg, sleep aids, muscle relaxers, narcotic pain medicines) while you are using this medicine; it may add to their effects. Ask your pharmacist if you

have questions about which medicines may cause drowsiness. This medicine may cause drowsiness or dizziness. DO NOT DRIVE, OPERATE MACHINERY, OR DO ANYTHING ELSE THAT COULD BE DANGEROUS until you know how you react to this medicine. Using this medicine alone, with other medicines, or with alcohol may lessen your ability to drive or to perform other potentially dangerous tasks. Alcohol, hot weather, exercise, or fever may increase dizziness. IF DIZZINESS OCCURS, sit up or stand slowly, especially in the morning. Sit or lie down at the first sign of this effect. THIS MEDICINE MAY INCREASE YOUR RISK OF SEIZURES. Your risk may be greater if you also have certain medical conditions, use certain medicines, or if you use a lot of alcohol. Talk to your doctor to see if you may have a greater risk of seizures while taking this medicine. SEROTONIN SYNDROME is a possibly fatal syndrome that can be caused by this medicine. Your risk may be greater if you take this medicine with certain other medicines (eg, "triptans," MAOIs, antidepressants). Symptoms may include agitation; confusion; hallucinations; coma; fever; fast or irregular heartbeat; tremor; excessive sweating; and nausea, vomiting, or diarrhea. Contact your doctor at once if you have any of these symptoms. BEFORE YOU BEGIN TAKING ANY NEW MEDICINES, either prescription or over-the-counter, check with your doctor or pharmacist. CAUTION IS ADVISED WHEN USING THIS MEDICINE IN THE ELDERLY because they may be more sensitive to the effects of this medicine. FOR WOMEN: This medicine may cause harm to the fetus. IF YOU THINK YOU MAY BE PREGNANT, discuss with your doctor the benefits and risks of using this medicine during pregnancy. THIS MEDICINE IS EXCRETED in breast milk. DO NOT BREAST-FEED while taking this medicine.

**POSSIBLE SIDE EFFECTS:** SIDE EFFECTS that may occur while taking this medicine include constipation; diarrhea; dizziness; drowsiness; dry mouth; headache; indigestion; mildly increased sweating; mild itching; nausea; trouble sleeping; vomiting; weakness. If they continue or are bothersome, check with your doctor. CONTACT YOUR DOCTOR IMMEDIATELY if you experience agitation; chest pain; confusion; difficult or painful urination; excessive sweating; fainting; fast or irregular heartbeat; fever; hallucinations; loss of coordination; mood or mental changes (eg, depression); red, blistered, swollen, or peeling skin; seizures; severe dizziness or lightheadedness; severe nausea, vomiting, or diarrhea; severe or persistent headache; slow or shallow breathing; suicidal thoughts or behaviors; tremor; or vision problems. AN ALLERGIC REACTION to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, or tongue; unusual hoarseness. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your health care provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** IF OVERDOSE IS SUSPECTED, contact your local poison control center or emergency room immediately. Symptoms may include bluish skin; cold, clammy skin; difficult, shallow, or slow breathing; drowsiness leading to unresponsiveness or coma; excessive sweating; limp muscles; pinpoint pupils; seizures; slow or irregular heartbeat.

**ADDITIONAL INFORMATION:** DO NOT SHARE THIS MEDICINE with others for whom it was not prescribed. DO NOT USE THIS MEDICINE for other health conditions. IF YOU WILL BE USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME, be sure to obtain necessary refills before your supply runs out. CHECK WITH YOUR PHARMACIST about how to dispose of unused medicine.

**KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.**

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

**RX # 0510864-09015**

DATE: 06/10/13

**TRAMADOL 50MG TABLETS**

QTY: 180 1 REFILL BEFORE 04/07/14

Refill NDC: 68382-0319-10

Retail Price: \$77.97 Your Insurance Saved You: \$67.97

**\$ 10.00**

W. CHRISTENSEN, MD  
MFG: ZYDUS  
SDB/ / ADO/ADO

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 131612711486024999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Customer  
Receipt

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

**RX # 0510864-09015**

DATE: 06/10/13

**TRAMADOL 50MG TABLETS**

QTY: 180 1 REFILL BEFORE 04/07/14

Refill NDC: 68382-0319-10

Retail Price: \$77.97 Your Insurance Saved You: \$67.97

**\$ 10.00**

W. CHRISTENSEN, MD  
MFG: ZYDUS  
SDB/ / ADO/ADO

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 131612711486024999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Duplicate  
Receipt

Pharmacy use only

MON 11:01AM  
Refill

TRAMADOL 50MG TABLETS  
68382-0319-10  
FAST RACK

QTY 180  
20 DRAM



WHITE  
FRONT: 319

SDB/ / ADO/ADO

Do not flush unused medications or pour down a sink or drain.

WIC# 216392



**INGREDIENT NAME:** HYDROCODONE (hye-droe-KO-done) and ACETAMINOPHEN (a-seat-a-MIN-oh-fen)

**COMMON USES:** This medicine is a narcotic and analgesic combination used to relieve moderate to moderately severe pain. It may also be used to treat other conditions as determined by your doctor.

**BEFORE USING THIS MEDICINE:** WARNING: THIS MEDICINE CONTAINS ACETAMINOPHEN. Severe and sometimes fatal liver problems, including the need for liver transplant, have been reported with the use of acetaminophen. Most cases of these liver problems occurred in patients taking excessive doses of acetaminophen (more than 4,000 mg per day). Also, patients who developed these liver problems were often using more than 1 medicine that contained acetaminophen. Discuss any questions or concerns with your doctor. Some medicines or medical conditions may interact with this medicine. **INFORM YOUR DOCTOR OR PHARMACIST** of all prescription and over-the-counter medicine that you are taking. **DO NOT TAKE THIS MEDICINE** if you are also taking sodium oxybate (XGB). **ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION** may be needed if you are taking anticoagulants (eg, warfarin), buprenorphine, isoniazid, linezolid, monoamine oxidase inhibitors (MAOIs) (eg, phenelzine), naltrexone, other narcotic pain medicines (eg, morphine), perampanel, sleep medicines (eg, zolpidem), tricyclic antidepressants (eg, amitriptyline), or medicines that may harm the liver (eg, methotrexate, ketoconazole, certain medicines for HIV infection). Ask your doctor if you are unsure if any of your medicines might harm the liver. **DO NOT START OR STOP** any medicine without doctor or pharmacist approval. Inform your doctor of any other medical conditions, including a history of liver problems (eg, hepatitis), kidney problems, lung or breathing problems (eg, asthma, chronic obstructive pulmonary disease / COPD), drug or alcohol abuse or dependence, heart problems (eg, cor pulmonale, slow or irregular heartbeat), low blood pressure, stomach or bowel problems (eg, blockage, inflammation, paralysis), gallbladder problems, prostate problems (eg, an enlarged prostate), difficulty urinating, seizures, thyroid problems, adrenal gland problems (eg, Addison disease), mental or mood problems, inflammation of the pancreas (pancreatitis), curvature of the spine (scoliosis), suicidal thoughts or actions, allergies, pregnancy, or breast-feeding. Tell your doctor if you have severe drowsiness, or have or recently have had any head injury, increased pressure in the brain, growths in the brain (eg, tumors), or infection of the brain or nervous system. Tell your doctor if you've recently had surgery (eg, stomach or bowel surgery) or are scheduled to have surgery. Tell your doctor if you are very poor or are very overweight or are dehydrated; have low blood volume, stomach pain, or constipation; drink alcohol; or are going through withdrawal from alcohol or other substances. **USE OF THIS MEDICINE IS NOT RECOMMENDED** if you have very slow or difficult breathing, or you are having an asthma attack. Contact your doctor or pharmacist if you have any questions or concerns about taking this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for using this medicine provided by your doctor. **TAKE THIS MEDICINE** by mouth with or without food. If stomach upset occurs, take with food to reduce stomach irritation. **IF YOU HAVE BEEN TAKING THIS MEDICINE REGULARLY** or for longer than a few weeks, do not suddenly stop taking it without checking with your doctor. Your doctor may need to gradually lower your dose. **SOME OF THESE PRODUCTS CONTAIN SULFITES.** Sulfites may cause an allergic reaction in some patients (eg, asthma patients). If you have ever had an allergic reaction to sulfites, ask your pharmacist if your product has sulfites in it. **STORE THIS MEDICINE** at room temperature, between 68 and 77 degrees F (20 and 25 degrees C), away from heat, moisture, and light. Do not store in the bathroom. **KEEP THIS MEDICINE** out of the reach of children and away from pets. **IF YOU MISS A DOSE OF THIS MEDICINE** and you are taking it regularly, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.

**CAUTIONS:** **DO NOT TAKE THIS MEDICINE** if you are allergic to any ingredient in this medicine. **BEFORE USING THIS MEDICINE**, check with your doctor if you have had an allergic reaction to other narcotic medicines (eg, codeine, morphine, oxycodone). **DO NOT EXCEED THE RECOMMENDED DOSE**, take more often than prescribed, or use for longer than prescribed without checking with your doctor. Exceeding the recommended dose or taking this medicine for longer than prescribed may be habit-forming. If using this medicine for an extended period of time, **DO NOT SUDDENLY STOP** taking this medicine without your doctor's approval. When using for an extended period, this medicine may not work as well and may require different dosing. Talk with your doctor if this medicine stops working well. **THIS MEDICINE MAY AFFECT CERTAIN LAB TEST RESULTS.** Make sure that all of your doctors and laboratory know you are taking this medicine. **BEFORE YOU HAVE ANY MEDICAL OR DENTAL TREATMENTS, emergency care, or surgery**, tell the doctor or dentist that you are using this medicine. **THIS MEDICINE MAY CAUSE DIZZINESS, lightheadedness, or fainting.** Alcohol, hot weather, exercise, and fever can increase these effects. To prevent them, sit up or stand slowly, especially in the morning. Also, sit or lie down at the first sign of dizziness, lightheadedness, or weakness. **THIS MEDICINE MAY ALSO CAUSE DROWSINESS OR BLURRED VISION.** Do not drive, operate machinery, or do anything else that could be dangerous until you know how you react to this medicine. Using this medicine alone, with other medicines, or with alcohol may lessen your ability to drive or to perform other potentially dangerous tasks. **DO NOT DRINK ALCOHOL** while you are using this medicine. **CHECK WITH YOUR DOCTOR** before you use medicines that may cause drowsiness (eg, sleep aids, muscle relaxers) while you are using this medicine; it may add to their effects. Ask your pharmacist if you have questions about which medicines may cause drowsiness. **THIS MEDICINE CONTAINS ACETAMINOPHEN.** Do not take additional acetaminophen for pain or fever without checking with your doctor or pharmacist. Ask your pharmacist if you have questions about which medicines contain acetaminophen. **THIS MEDICINE MAY HARM YOUR LIVER.** Your risk may be greater if you drink alcohol while you are using this medicine. Talk to your doctor before you take this medicine or other fever reducers if you drink alcohol. **CONTACT YOUR DOCTOR RIGHT AWAY** if you take more than 4,000 mg of acetaminophen per day, even if you feel well. **BEFORE YOU BEGIN TAKING ANY NEW MEDICINE**, either prescription or over-the-counter, check with your doctor or pharmacist. **CAUTION IS ADVISED WHEN USING THIS MEDICINE IN THE ELDERLY**, because they may be more sensitive to the effects of the medicine. **FOR YOU TO BECOME PREGNANT**, discuss with your doctor the benefits and risks of using this medicine during pregnancy. Long-term use of this medicine during pregnancy may cause dependence in the fetus or newborn. Discuss any questions or concerns with your doctor. **THIS MEDICINE IS EXCRETED** in breast milk. **DO NOT BREAST-FEED** while taking this medicine.

**POSSIBLE SIDE EFFECTS:** SIDE EFFECTS that may occur while taking this medicine include dizziness, drowsiness, lightheadedness, nausea, or vomiting. If

they continue or are bothersome, check with your doctor. **THIS MEDICINE MAY CAUSE CONSTIPATION.** Talk with your doctor or pharmacist about using a stool softener or laxative to prevent constipation. It is also important to maintain a diet adequate in fiber, drink plenty of water, and exercise to prevent constipation. If you become constipated while taking this medicine, talk with your doctor or pharmacist. **CONTACT YOUR DOCTOR IMMEDIATELY** if you experience a change in the amount of urine produced; change or loss in hearing; chest pain; confusion; fainting; fever, chills, or persistent sore throat; mental or mood changes (eg, agitation, anxiety, depression); red, swollen, blistered, or peeling skin; severe or persistent constipation; severe or persistent dizziness, drowsiness, headache, or lightheadedness; shortness of breath; slow, shallow, or difficult breathing; symptoms of liver problems (eg, yellowing of the skin or eyes, pale stools, dark urine, persistent loss of appetite, unusual or severe stomach pain); trouble urinating; unusual bruising or bleeding; unusual tiredness or weakness; or vision changes. **AN ALLERGIC REACTION** to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, throat, or tongue; or unusual hoarseness. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** IF OVERDOSE IS SUSPECTED, contact your local poison control center or emergency room immediately. Symptoms of overdose may include bluish skin or nails; cold and clammy skin; coma; decreased urination; difficult, shallow, or slow breathing; fainting; limp muscles; loss of consciousness; nausea or vomiting; pinpoint pupils; seizures; severe dizziness, drowsiness, or light-headedness; slow heartbeat; symptoms of liver problems (eg, yellowing of the skin or eyes, pale stools, dark urine, loss of appetite, unusual or severe stomach pain); unusual bruising or bleeding; or unusual sweating or tiredness.

**ADDITIONAL INFORMATION:** If your symptoms do not improve or if they become worse, contact your doctor. **DO NOT SHARE THIS MEDICINE** with others for whom it was not prescribed. **DO NOT USE THIS MEDICINE** for other health conditions. **CHECK WITH YOUR PHARMACIST** about how to dispose of unused medicine.

**KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.**

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

**RX # 0529365-09015**

DATE: 07/09/13

**HYDROCODONE /ACETAMINOPHEN 5-325 TB**

QTY: 120 1 REFILL BEFORE 11/20/13

Copy NDC: 00591-3202-01

Retail Price: \$87.09 Your Insurance Saved You: \$77.09

\$ 10.00

A. PARK, MD  
MFG: WATSON  
SDB/HYK/HYK/ADO/ADO

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 131906514767024999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Customer  
Receipt

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

**RX # 0529365-09015**

DATE: 07/09/13

**HYDROCODONE /ACETAMINOPHEN 5-325 TB**

QTY: 120 1 REFILL BEFORE 11/20/13

Copy NDC: 00591-3202-01

Retail Price: \$87.09 Your Insurance Saved You: \$77.09

\$ 10.00

A. PARK, MD  
MFG: WATSON  
SDB/HYK/HYK/ADO/ADO

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 131906514767024999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951


Duplicate  
Receipt

Pharmacy use only

TUE 9:35PM  
Copy

HYDROCODONE /ACETAMINOPHEN 5-325 TB  
00591-3202-01  
FAST RACK

QTY 120  
30 DRAM



WHITE AND MULTICOLOR  
FRONT: 3202  
BACK: WATSON

SDB/HYK/HYK/ADO/ADO

Call your doctor for medical advice about side effects.  
You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 216382



Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**INGREDIENT NAME:** TRAMADOL (TRAM-a-dol)

**COMMON USES:** This medicine is an analgesic used to treat moderate to moderately severe pain.

**BEFORE USING THIS MEDICINE:** Some medicines or medical conditions may interact with this medicine. **INFORM YOUR DOCTOR OR PHARMACIST** of all prescription and over-the-counter medicine that you are taking. **DO NOT TAKE THIS MEDICINE** if you are taking sodium oxybate (GHB), or another product that contains tramadol. **ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION** may be needed if you are taking alpha-2 receptor blockers (eg, yohimbine), anorexiant (eg, phentermine), anticoagulants (eg, warfarin), azole antifungals (eg, ketoconazole), butyrophenones (eg, haloperidol), carbamazepine, cyclobenzaprine, digoxin, furazolidone, linezolid, lithium, loxapine, macrolide antibiotics (eg, erythromycin), certain medicines for mental or mood disorders (eg, olanzapine), molindone, monoamine oxidase inhibitors (MAOIs) (eg, phenelzine, selegiline), nefazodone, opioid pain medicine (eg, codeine, hydrocodone), phenothiazines (eg, promethazine), quinine, rifampin, serotonin-norepinephrine reuptake inhibitors (SNRIs) (eg, duloxetine), selective serotonin reuptake inhibitors (SSRIs) (eg, fluoxetine, paroxetine), sleeping medicines (eg, zolpidem), St. John's wort, thioxanthenes (eg, thiothixene), tiagabine, tricyclic antidepressants (eg, amitriptyline), or "triptans" (eg, sumatriptan). **DO NOT START OR STOP** any medicine without doctor or pharmacist approval. Inform your doctor of any other medical conditions, including liver, kidney, or thyroid problems; blockage in the bladder; diabetes; heart problems (eg, cor pulmonale); high blood pressure; pancreas problems; prostate problems; metabolism problems; allergies; pregnancy; or breast-feeding. Tell your doctor if you have or recently have had any head injury, brain injury or tumor, increased pressure in the brain, or infection of the brain or nervous system. Tell your doctor if you have a history of lung or breathing problems (eg, asthma, chronic obstructive pulmonary disease / COPD); seizures (eg, epilepsy); recent stomach or bowel surgery; other stomach or bowel problems (eg, pain, inflammation, ulcers); alcohol or other substance abuse; mood or mental problems (eg, depression); or suicidal thoughts or actions. Tell your doctor if you drink alcohol or if you are going through withdrawal from alcohol or other substances. **USE OF THIS MEDICINE IS NOT RECOMMENDED** if you have severe or persistent diarrhea caused by antibiotic use; suicidal thoughts or actions; or a history of alcohol or other substance abuse or addiction. **USE OF THIS MEDICINE IS NOT RECOMMENDED** if you are intoxicated with alcohol, opioids or narcotics (eg, codeine, morphine), or sedatives or sleeping medicines (eg, temazepam, zolpidem). This medicine should not be used in **CHILDREN** younger than 16 years; safety and effectiveness in these children have not been confirmed. Contact your doctor or pharmacist if you have any questions or concerns about using this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for taking this medicine provided by your doctor. **TAKE THIS MEDICINE** by mouth with or without food. **STORE THIS MEDICINE** at 77 degrees F (25 degrees C) away from heat, moisture, and light. Do not store in the bathroom. Brief storage at temperatures between 59 and 86 degrees F (15 and 30 degrees C) is permitted. **KEEP THIS MEDICINE** out of the reach of children and away from pets. **IF YOU MISS A DOSE OF THIS MEDICINE** and you are taking it regularly, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. **DO NOT** take 2 doses at once.

**CAUTIONS:** **DO NOT TAKE THIS MEDICINE** if you have had an allergic reaction to it or if you are allergic to any ingredient in this product. **DO NOT TAKE THIS MEDICINE** if you have had a severe allergic reaction to codeine or another opioid (eg, morphine). **DO NOT EXCEED THE RECOMMENDED DOSE** or take this medicine for longer than prescribed without checking with your doctor. Exceeding the recommended dose or taking this medicine for longer than prescribed may be habit-forming or may increase the risk of seizures or breathing problems. If using this medicine for an extended period of time, **DO NOT SUDDENLY STOP** taking this medicine without your doctor's approval. Your dose may need to be slowly lowered to avoid side effects. When used for an extended period of time, this medicine may not work as well and may require different dosing. Talk with your doctor if this medicine stops working well. **DO NOT TAKE MORE THAN PRESCRIBED. BEFORE YOU HAVE ANY MEDICAL OR DENTAL TREATMENTS, EMERGENCY CARE, OR SURGERY**, tell the doctor or dentist that you are using this medicine. **DO NOT DRINK ALCOHOL** while you are using this medicine. **CHECK WITH YOUR DOCTOR** before you use medicines that may cause drowsiness (eg, sleep aids, muscle relaxers, narcotic pain medicines) while you are using this medicine; it may add to their effects. Ask your pharmacist if you

have questions about which medicines may cause drowsiness. This medicine may cause drowsiness or dizziness. **DO NOT DRIVE, OPERATE MACHINERY, OR DO ANYTHING ELSE THAT COULD BE DANGEROUS** until you know how you react to this medicine. Using this medicine alone, with other medicines, or with alcohol may lessen your ability to drive or to perform other potentially dangerous tasks. Alcohol, hot weather, exercise, or fever may increase dizziness. **IF DIZZINESS OCCURS**, sit up or stand slowly, especially in the morning. Sit or lie down at the first sign of this effect. **THIS MEDICINE MAY INCREASE YOUR RISK OF SEIZURES.** Your risk may be greater if you also have certain medical conditions, use certain medicines, or if you use a lot of alcohol. Talk to your doctor to see if you may have a greater risk of seizures while taking this medicine. **SEROTONIN SYNDROME** is a possibly fatal syndrome that can be caused by this medicine. Your risk may be greater if you take this medicine with certain other medicines (eg, "triptans," MAOIs, antidepressants). Symptoms may include agitation; confusion; hallucinations; coma; fever; fast or irregular heartbeat; tremor; excessive sweating; and nausea, vomiting, or diarrhea. Contact your doctor at once if you have any of these symptoms. **BEFORE YOU BEGIN TAKING ANY NEW MEDICINES**, either prescription or over-the-counter, check with your doctor or pharmacist. **CAUTION IS ADVISED WHEN USING THIS MEDICINE IN THE ELDERLY** because they may be more sensitive to the effects of this medicine. **FOR WOMEN:** This medicine may cause harm to the fetus. **IF YOU THINK YOU MAY BE PREGNANT**, discuss with your doctor the benefits and risks of using this medicine during pregnancy. **THIS MEDICINE IS EXCRETED** in breast milk. **DO NOT BREAST-FEED** while taking this medicine.

**POSSIBLE SIDE EFFECTS:** **SIDE EFFECTS** that may occur while taking this medicine include constipation; diarrhea; dizziness; drowsiness; dry mouth; headache; indigestion; mildly increased sweating; mild itching; nausea; trouble sleeping; vomiting; weakness. If they continue or are bothersome, check with your doctor. **CONTACT YOUR DOCTOR IMMEDIATELY** if you experience agitation; chest pain; confusion; difficult or painful urination; excessive sweating; fainting; fast or irregular heartbeat; fever; hallucinations; loss of coordination; mood or mental changes (eg, depression); red, blistered, swollen, or peeling skin; seizures; severe dizziness or lightheadedness; severe nausea, vomiting, or diarrhea; severe or persistent headache; slow or shallow breathing; suicidal thoughts or behaviors; tremor; or vision problems. **AN ALLERGIC REACTION** to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, or tongue; unusual hoarseness. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your health care provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** **IF OVERDOSE IS SUSPECTED**, contact your local poison control center or emergency room immediately. Symptoms may include bluish skin; cold, clammy skin; difficult, shallow, or slow breathing; drowsiness leading to unresponsiveness or coma; excessive sweating; limp muscles; pinpoint pupils; seizures; slow or irregular heartbeat.

**ADDITIONAL INFORMATION:** **DO NOT SHARE THIS MEDICINE** with others for whom it was not prescribed. **DO NOT USE THIS MEDICINE** for other health conditions. **IF YOU WILL BE USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME**, be sure to obtain necessary refills before your supply runs out. **CHECK WITH YOUR PHARMACIST** about how to dispose of unused medicine.

**KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.**

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

**RX # 0510864-09015**

DATE: 07/09/13

**TRAMADOL 50MG TABLETS**

QTY: 180 NO REFILLS - DR. AUTH REQUIRED

Refill NDC: 68382-0319-10

Retail Price: \$77.97 Your Insurance Saved You: \$67.97

\$ **10.00**

W. CHRISTENSEN, MD  
MFG: ZYDUS  
SDB/ / ADO/ADO

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 131906513334053999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Customer  
Receipt

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

**RX # 0510864-09015**

DATE: 07/09/13

**TRAMADOL 50MG TABLETS**

QTY: 180 NO REFILLS - DR. AUTH REQUIRED

Refill NDC: 68382-0319-10

Retail Price: \$77.97 Your Insurance Saved You: \$67.97

\$ **10.00**

W. CHRISTENSEN, MD  
MFG: ZYDUS  
SDB/ / ADO/ADO

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 131906513334053999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Duplicate  
Receipt

Pharmacy use only

TUE 9:35PM

Refill

TRAMADOL 50MG TABLETS

68382-0319-10

FAST RACK

QTY 180

20 DRAM



WHITE  
FRONT: 319

SDB/ / ADO/ADO

Do not flush unused medications or pour down a sink or drain.

WIC# 216382