

# YOUR PERSONAL PRESCRIPTION INFORMATION

## Your Walgreens Pharmacy Location

4702 N Jim Miller Rd  
Dallas, TX 75227-  
(214)388-4951

IF YOU HAVE QUESTIONS ABOUT YOUR PRESCRIPTION, PLEASE CONTACT YOUR WALGREENS PHARMACIST OR CALL 1-800-WALGREENS.

PATIENT ROBERT A PLOCK

BIRTH DATE 07/26/68

MEDICATION HYDROCODONE /ACETAMINOPHEN 10-325 T


QUANTITY 120

DIRECTIONS TAKE 1 TABLET BY MOUTH EVERY 4 HOURS AS NEEDED FOR PAIN. MAX 6 TABLETS A DAY

DOCTOR DR A. PARK

PATIENT ALLERGIES

DRUG DESCRIPTION



WHITE

FRONT: WATSON 853

INGREDIENT NAME: HYDROCODONE (hye-droe-KO-done) and ACETAMINOPHEN (a-seat-a-MIN-oh-fen)

COMMON USES: This medicine is a narcotic and analgesic combination used to relieve moderate to moderately severe pain. It may also be used to treat other conditions as determined by your doctor.

**BEFORE USING THIS MEDICINE: WARNING: THIS MEDICINE CONTAINS ACETAMINOPHEN.** Severe and sometimes fatal liver problems, including the need for liver transplant, have been reported with the use of acetaminophen. Most cases of these liver problems occurred in patients taking excessive doses of acetaminophen (more than 4,000 mg per day). Also, patients who developed these liver problems were often using more than 1 medicine that contained acetaminophen. Discuss any questions or concerns with your doctor. Some medicines or medical conditions may interact with this medicine. **INFORM YOUR DOCTOR OR PHARMACIST** of all prescription and over-the-counter medicine that you are taking. **DO NOT TAKE THIS MEDICINE** if you are also taking sodium oxybate (GHB). **ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION** may be needed if you are taking anticoagulants (eg, warfarin), buprenorphine, isoniazid, linezolid, monoamine oxidase inhibitors (MAOIs) (eg, phenelzine), naltrexone, other narcotic pain medicines (eg, morphine), perampanel, sleep medicines (eg, zolpidem), tricyclic antidepressants (eg, amitriptyline), or medicines that may harm the liver (eg, methotrexate, ketoconazole, certain medicines for HIV infection). Ask your doctor if you are unsure if any of your medicines might harm the liver. **DO NOT START OR STOP** any medicine without doctor or pharmacist approval. Inform your doctor of any other medical conditions, including a history of liver problems (eg, hepatitis), kidney problems, lung or breathing problems (eg, asthma, chronic obstructive pulmonary disease / COPD), drug or alcohol abuse or dependence, heart problems (eg, cor pulmonale, slow or irregular heartbeat), low blood pressure, stomach or bowel problems (eg, blockage, inflammation, paralysis), gallbladder problems, prostate problems (eg, an enlarged prostate), difficulty urinating, seizures, thyroid problems, adrenal gland problems (eg, Addison disease), mental or mood problems, inflammation of the pancreas (pancreatitis), curvature of the spine (scoliosis), suicidal thoughts or actions, allergies, pregnancy, or breast-feeding. Tell your doctor if you have severe drowsiness, or have or recently have had any head injury, increased pressure in the brain, growths in the brain (eg, tumors), or infection of the brain or nervous system. Tell your doctor if you have recently had surgery (eg, stomach or bowel surgery) or are scheduled to have surgery. Tell your doctor if you are in very poor health; are very overweight or are dehydrated; have low blood volume, stomach pain, or constipation; drink alcohol; or are going through withdrawal from alcohol or other substances. **USE OF THIS MEDICINE IS NOT RECOMMENDED** if you have very slow or difficult breathing, or you are having an asthma attack. Contact your doctor or pharmacist if you have any questions or concerns about taking this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for using this medicine provided by your doctor. **TAKE THIS MEDICINE** by mouth with or without food. If stomach upset occurs, take with food to reduce stomach irritation. **IF YOU HAVE BEEN TAKING THIS MEDICINE REGULARLY** or for longer than a few weeks, do not suddenly stop taking it without checking with your doctor. Your doctor may need to gradually lower your dose. **SOME OF THESE PRODUCTS CONTAIN SULFITES.** Sulfites may cause an allergic reaction in some patients (eg, asthma patients). If you have ever had an allergic reaction to sulfites, ask your pharmacist if your product has sulfites in it. **STORE THIS MEDICINE** at room temperature, between 68 and 77 degrees F (20 and 25 degrees C), away from heat, moisture, and light. **DO NOT store** in the bathroom. **KEEP THIS MEDICINE** out of the reach of children and away from pets. **IF YOU MISS A DOSE OF THIS MEDICINE** and you are taking it regularly, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.

**CAUTIONS: DO NOT TAKE THIS MEDICINE** if you are allergic to any ingredient in this medicine. **BEFORE USING THIS MEDICINE,** check with your doctor if you have had an allergic reaction to other narcotic medicines (eg, codeine, morphine, oxycodone). **DO NOT EXCEED THE RECOMMENDED DOSE,** take more often than prescribed, or use more than prescribed without checking with your doctor. Exceeding the recommended dose or taking this medicine for longer than prescribed may be habit-forming. If using this medicine for an extended period of time, **DO NOT SUDDENLY STOP** taking this medicine without your doctor's approval. When using for an extended period, this medicine may not work as well and may require different dosing. Talk with your doctor if this medicine stops working well. **THIS MEDICINE MAY AFFECT CERTAIN LAB TEST RESULTS.** Make sure that all of your doctors and laboratory personnel know you are taking this medicine. **BEFORE YOU HAVE ANY MEDICAL OR DENTAL TREATMENTS,** emergency care, or surgery, tell the doctor or dentist that you are using this medicine. **THIS MEDICINE MAY CAUSE DIZZINESS,** lightheadedness, or fainting. Alcohol, hot weather, exercise, and fever can increase these effects. To prevent them, sit up or stand slowly, especially in the morning. Also, sit or lie down at the first sign of dizziness, lightheadedness, or weakness. **THIS MEDICINE MAY ALSO CAUSE DROWSINESS OR BLURRED VISION.** Do not drive, operate machinery, or do anything else that could be dangerous until you know how you react to this medicine. Using this medicine alone, with other medicines, or with alcohol may lessen your ability to drive or to perform other potentially dangerous tasks. **DO NOT DRINK ALCOHOL** while you are using this medicine. **CHECK WITH YOUR DOCTOR** before you use medicines that may cause drowsiness (eg, sleep aids, muscle relaxers) while you are using this medicine; it may add to their effects. Ask your pharmacist if you have questions about which medicines may cause drowsiness. **THIS MEDICINE CONTAINS ACETAMINOPHEN.** Do not take additional acetaminophen for pain or fever without checking with your doctor or pharmacist. Ask your pharmacist if you have questions about which medicines contain acetaminophen. **THIS MEDICINE MAY HARM YOUR LIVER.** Your risk may be greater if you drink alcohol while you are using this medicine. Talk to your doctor before you take this medicine or other fever reducers if you drink alcohol. **CONTACT YOUR DOCTOR RIGHT AWAY** if you take more than 4,000 mg of acetaminophen per day, even if you feel well. **BEFORE YOU BEGIN TAKING ANY NEW MEDICINE,** either prescription or over-the-counter, check with your doctor or pharmacist. **CAUTION IS ADVISED WHEN USING THIS MEDICINE IN THE ELDERLY** because they may be more sensitive to the effects of the medicine. **FOR WOMEN: IF YOU BECOME PREGNANT,** discuss with your doctor the benefits and risks of using this medicine during pregnancy. Long-term use of this medicine during pregnancy may cause dependence in the fetus or newborn. Discuss any questions or concerns with your doctor. **THIS MEDICINE IS EXCRETED** in breast milk. **DO NOT BREAST-FEED** while taking this medicine.

**POSSIBLE SIDE EFFECTS:** SIDE EFFECTS that may occur while taking this medicine include dizziness, drowsiness, lightheadedness, nausea, or vomiting. If

they continue or are bothersome, check with your doctor. **THIS MEDICINE MAY CAUSE CONSTIPATION.** Talk with your doctor or pharmacist about using a stool softener or laxative to prevent constipation. It is also important to maintain a diet adequate in fiber, drink plenty of water, and exercise to prevent constipation. If you become constipated while taking this medicine, talk with your doctor or pharmacist. **A VERY BAD SKIN REACTION** (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever); red or irritated eyes; or sores in your mouth, throat, nose, or eyes. **CONTACT YOUR DOCTOR IMMEDIATELY** if you experience a change in the amount of urine produced; change or loss in hearing; chest pain; confusion; fainting; fever, chills, or persistent sore throat; mental or mood changes (eg, agitation, anxiety, depression); severe or persistent constipation; severe or persistent dizziness, drowsiness, headache, or lightheadedness; shortness of breath; slow, shallow, or difficult breathing; symptoms of liver problems (eg, yellowing of the skin or eyes, pale stools, dark urine, persistent loss of appetite, unusual or severe stomach pain); trouble urinating; unusual bruising or bleeding; unusual tiredness or weakness; or vision changes. **AN ALLERGIC REACTION** to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, throat, or tongue; or unusual hoarseness. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE: IF OVERDOSE IS SUSPECTED,** contact your local poison control center or emergency room immediately. Symptoms of overdose may include bluish skin or nails; cold and clammy skin; coma; decreased urination; difficult, shallow, or slow breathing; fainting; limp muscles; loss of consciousness; nausea or vomiting; pinpoint pupils; seizures; severe dizziness, drowsiness, or light-headedness; slow heartbeat; symptoms of liver problems (eg, yellowing of the skin or eyes, pale stools, dark urine, loss of appetite, unusual or severe stomach pain); unusual bruising or bleeding; or unusual sweating or tiredness.

**ADDITIONAL INFORMATION:** If your symptoms do not improve or if they become worse, contact your doctor. **DO NOT SHARE THIS MEDICINE** with others for whom it was not prescribed. **DO NOT USE THIS MEDICINE** for other health conditions. **CHECK WITH YOUR PHARMACIST** about how to dispose of unused medicine.

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

ROBERT A PLOCK

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

RX # 0544507-09015

DATE: 09/25/13

HYDROCODONE /ACETAMINOPHEN 10-325 T

QTY: 120 NO REFILLS - DR. AUTH REQUIRED

New

NDC: 00591-2612-05

Retail Price: \$91.99 Your Insurance Saved You: \$81.99

\$ 10.00

DR A. PARK

PLAN: W/PB

MFG: WATSON

GROUP# 01960917

SIM/DBP/DBP/DBP/DBP

CLAIM REF# 132683398818059999

ROBERT A PLOCK

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

RX # 0544507-09015

DATE: 09/25/13

HYDROCODONE /ACETAMINOPHEN 10-325 T

QTY: 120 NO REFILLS - DR. AUTH REQUIRED

New

NDC: 00591-2612-05

Retail Price: \$91.99 Your Insurance Saved You: \$81.99

\$ 10.00

DR A. PARK

PLAN: W/PB

MFG: WATSON

GROUP# 01960917

SIM/DBP/DBP/DBP/DBP

CLAIM REF# 132683398818059999

Walgreens

4702 N JIM MILLER RD DALLAS, TX 75227

PH: (214)388-4951

Customer Receipt

Pharmacy use only

WED 12:45PM

New

HYDROCODONE /ACETAMINOPHEN 10-325 T

00591-2612-05

FAST RACK

QTY 120

30 DRAM



WHITE

FRONT: WATSON 853

SIM/DBP/DBP/DBP/DBP

Do not flush unused medications or pour down a sink or drain.

WIC# 216382

Walgreens

4702 N JIM MILLER RD DALLAS, TX 75227


PH: (214)388-4951

Duplicate Receipt

YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location  
4702 N Jim Miller Rd  
Dallas, TX 75227-  
(214)388-4951

IF YOU HAVE QUESTIONS ABOUT YOUR PRESCRIPTION, PLEASE CONTACT YOUR WALGREENS PHARMACIST OR CALL 1-800-WALGREENS.

PATIENT	ROBERT A PLOCK	DOCTOR	DR A. PARK	DRUG DESCRIPTION
BIRTH DATE	07/26/68			
MEDICATION	TRAMADOL 50MG TABLETS			
QUANTITY	120	PATIENT ALLERGIES		
DIRECTIONS TAKE 1 TABLET BY MOUTH EVERY 6 HOURS AS NEEDED FOR PAIN				WHITE FRONT: 319

INGREDIENT NAME: TRAMADOL (TRAM-a-dol)  
COMMON USES: This medicine is an analgesic used to treat moderate to moderately severe pain.

BEFORE USING THIS MEDICINE: Some medicines or medical conditions may interact with this medicine. INFORM YOUR DOCTOR OR PHARMACIST of all prescription and over-the-counter medicine that you are taking. DO NOT TAKE THIS MEDICINE if you are taking sodium oxybate (GHB), or another product that contains tramadol. ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION may be needed if you are taking alpha-2 receptor blockers (eg, yohimbine), anorexiant (eg, phentermine), anticoagulants (eg, warfarin), azole antifungals (eg, ketoconazole), butyrophenones (eg, haloperidol), carbamazepine, cyclobenzaprine, digoxin, furazolidone, linezolid, lithium, loxapine, macrolide antibiotics (eg, erythromycin), certain medicines for mental or mood disorders (eg, olanzapine), molidone, monoamine oxidase inhibitors (MAOIs) (eg, phenelzine, selegiline), nefazodone, opioid pain medicine (eg, codeine, hydrocodone), phenothiazines (eg, promethazine), quinidine, rifampin, serotonin-norepinephrine reuptake inhibitors (SNRIs) (eg, duloxetine), selective serotonin reuptake inhibitors (SSRIs) (eg, fluoxetine, paroxetine), sleeping medicines (eg, zolpidem), St. John's wort, thioxanthenes (eg, thiothixene), tiagabine, tricyclic antidepressants (eg, amitriptyline), or "triptans" (eg, sumatriptan). DO NOT START OR STOP any medicine without doctor or pharmacist approval. Inform your doctor of any other medical conditions, including liver, kidney, or thyroid problems; blockage in the bladder; diabetes; heart problems (eg, cor pulmonale); high blood pressure; pancreas problems; prostate problems; metabolism problems; allergies; pregnancy; or breast-feeding. Tell your doctor if you have or recently have had any head injury, brain injury or tumor, increased pressure in the brain, or infection of the brain or nervous system. Tell your doctor if you have a history of lung or breathing problems (eg, asthma, chronic obstructive pulmonary disease / COPD); seizures (eg, epilepsy); recent stomach or bowel surgery; other stomach or bowel problems (eg, pain, inflammation, ulcers); alcohol or other substance abuse; mood or mental problems (eg, depression); or suicidal thoughts or actions. Tell your doctor if you drink alcohol or if you are going through withdrawal from alcohol or other substances. USE OF THIS MEDICINE IS NOT RECOMMENDED if you have severe or persistent diarrhea caused by antibiotic use; suicidal thoughts or actions; or a history of alcohol or other substance abuse or addiction. USE OF THIS MEDICINE IS NOT RECOMMENDED if you are intoxicated with alcohol, opioids or narcotics (eg, codeine, morphine), or sedatives or sleeping medicines (eg, temazepam, zolpidem). This medicine should not be used in CHILDREN younger than 16 years; safety and effectiveness in these children have not been confirmed. Contact your doctor or pharmacist if you have any questions or concerns about using this medicine.

HOW TO USE THIS MEDICINE: Follow the directions for taking this medicine provided by your doctor. TAKE THIS MEDICINE by mouth with or without food. STORE THIS MEDICINE at 77 degrees F (25 degrees C) away from heat, moisture, and light. Do not store in the bathroom. Brief storage at temperatures between 59 and 86 degrees F (15 and 30 degrees C) is permitted. KEEP THIS MEDICINE out of the reach of children and away from pets. IF YOU MISS A DOSE OF THIS MEDICINE and you are taking it regularly, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. DO NOT take 2 doses at once.

CAUTIONS: DO NOT TAKE THIS MEDICINE if you have had an allergic reaction to it or if you are allergic to any ingredient in this product. DO NOT TAKE THIS MEDICINE if you have had a severe allergic reaction to codeine or another opioid (eg, morphine). DO NOT EXCEED THE RECOMMENDED DOSE or take this medicine for longer than prescribed without checking with your doctor. Exceeding the recommended dose or taking this medicine for longer than prescribed may be habit-forming or may increase the risk of seizures or breathing problems. If using this medicine for an extended period of time, DO NOT SUDDENLY STOP taking this medicine without your doctor's approval. Your dose may need to be slowly lowered to avoid side effects. When used for an extended period of time, this medicine may not work as well and may require different dosing. Talk with your doctor if this medicine stops working well. DO NOT TAKE MORE THAN PRESCRIBED. BEFORE YOU HAVE ANY MEDICAL OR DENTAL TREATMENTS, EMERGENCY CARE, OR SURGERY, tell the doctor or dentist that you are using this medicine. DO NOT DRINK ALCOHOL while you are using this medicine. CHECK WITH YOUR DOCTOR BEFORE you use medicines that may cause drowsiness (eg, sleep aids, muscle relaxers, narcotic pain medicines) while you are using this medicine; it may add to their effects. Ask your pharmacist if you

have questions about which medicines may cause drowsiness. This medicine may cause drowsiness or dizziness. DO NOT DRIVE, OPERATE MACHINERY, OR DO ANYTHING ELSE THAT COULD BE DANGEROUS until you know how you react to this medicine. Using this medicine alone, with other medicines, or with alcohol may lessen your ability to drive or to perform other potentially dangerous tasks. Alcohol, hot weather, exercise, or fever may increase dizziness. IF DIZZINESS OCCURS, sit up or stand slowly, especially in the morning. Sit or lie down at the first sign of this effect. THIS MEDICINE MAY INCREASE YOUR RISK OF SEIZURES. Your risk may be greater if you also have certain medical conditions, use certain medicines, or if you use a lot of alcohol. Talk to your doctor to see if you may have a greater risk of seizures while taking this medicine. SEROTONIN SYNDROME is a possibly fatal syndrome that can be caused by this medicine. Your risk may be greater if you take this medicine with certain other medicines (eg, "triptans," MAOIs, antidepressants). Symptoms may include agitation; confusion; hallucinations; coma; fever; fast or irregular heartbeat; tremor; excessive sweating; and nausea, vomiting, or diarrhea. Contact your doctor at once if you have any of these symptoms. BEFORE YOU BEGIN TAKING ANY NEW MEDICINES, either prescription or over-the-counter, check with your doctor or pharmacist. CAUTION IS ADVISED WHEN USING THIS MEDICINE IN THE ELDERLY because they may be more sensitive to the effects of this medicine. FOR WOMEN: This medicine may cause harm to the fetus. IF YOU THINK YOU MAY BE PREGNANT, discuss with your doctor the benefits and risks of using this medicine during pregnancy. THIS MEDICINE IS EXCRETED in breast milk. DO NOT BREAST-FEED while taking this medicine.

POSSIBLE SIDE EFFECTS: SIDE EFFECTS that may occur while taking this medicine include constipation; diarrhea; dizziness; drowsiness; dry mouth; headache; indigestion; mildly increased sweating; mild itching; nausea; trouble sleeping; vomiting; weakness. If they continue or are bothersome, check with your doctor. CONTACT YOUR DOCTOR IMMEDIATELY if you experience agitation; chest pain; confusion; difficult or painful urination; excessive sweating; fainting; fast or irregular heartbeat; fever; hallucinations; loss of coordination; mood or mental changes (eg, depression); red, blistered, swollen, or peeling skin; seizures; severe dizziness or lightheadedness; severe nausea, vomiting, or diarrhea; severe or persistent headache; slow or shallow breathing; suicidal thoughts or behaviors; tremor; or vision problems. AN ALLERGIC REACTION to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, or tongue; unusual hoarseness. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your health care provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

OVERDOSE: IF OVERDOSE IS SUSPECTED, contact your local poison control center or emergency room immediately. Symptoms may include bluish skin; cold, clammy skin; difficult, shallow, or slow breathing; drowsiness leading to unresponsiveness or coma; excessive sweating; limp muscles; pinpoint pupils; seizures; slow or irregular heartbeat.

ADDITIONAL INFORMATION: DO NOT SHARE THIS MEDICINE with others for whom it was not prescribed. DO NOT USE THIS MEDICINE for other health conditions. IF YOU WILL BE USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME, be sure to obtain necessary refills before your supply runs out. CHECK WITH YOUR PHARMACIST about how to dispose of unused medicine.

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

ROBERT A PLOCK 6827 Latta Pkwy, Dallas, TX 75227 (214)799-7775	
RX # 0534730-09015	DATE: 08/06/13
TRAMADOL 50MG TABLETS	
QTY: 120	1 + REFILLS BEFORE 07/24/14
Copy	NDC: 68382-0319-10
Retail Price: \$53.98	Your Insurance Saved You: \$43.98
\$ 10.00	
DR A. PARK MFG: ZYDUS NJT/NJT/NJT/ADO/ADO	PLAN: WPB GROUP# 01960917 CLAIM REF# 132186453108009998

Walgreens 4702 N JIM MILLER RD DALLAS, TX 75227-  
PH: (214)388-4951 Customer Receipt

Pharmacy use only

TUE 9:24PM  
Copy  
TRAMADOL 50MG TABLETS  
68382-0319-10  
FAST RACK

ROBERT A PLOCK 6827 Latta Pkwy, Dallas, TX 75227 (214)799-7775	
RX # 0534730-09015	DATE: 08/06/13
TRAMADOL 50MG TABLETS	
QTY: 120	1 + REFILLS BEFORE 07/24/14
Copy	NDC: 68382-0319-10
Retail Price: \$53.98	Your Insurance Saved You: \$43.98
\$ 10.00	
DR A. PARK MFG: ZYDUS NJT/NJT/NJT/ADO/ADO	PLAN: WPB GROUP# 01960917 CLAIM REF# 132186453108009998

Walgreens 4702 N JIM MILLER RD DALLAS, TX 75227-  
PH: (214)388-4951 Duplicate Receipt

QTY 120  
20 DRAM  
WHITE  
FRONT: 319  
NJT/NJT/NJT/ADO/ADO

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 216382



Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

WIC# 216382

Do not flush unused medications or pour down a sink or drain.




# YOUR PERSONAL PRESCRIPTION INFORMATION

## Your Walgreens Pharmacy Location

4702 N Jim Miller Rd  
Dallas, TX 75227-  
(214)388-4951

IF YOU HAVE QUESTIONS ABOUT YOUR PRESCRIPTION, PLEASE CONTACT YOUR WALGREENS PHARMACIST OR CALL 1-800-WALGREENS.

<b>PATIENT</b>	ROBERT A PLOCK	<b>DOCTOR</b>	A. PARK, MD	<b>DRUG DESCRIPTION</b>
<b>BIRTH DATE</b>	07/26/68			
<b>MEDICATION</b>	HYDROCODONE /ACETAMINOPHEN 5-325 TB	<b>PATIENT ALLERGIES</b>		
<b>QUANTITY</b>	120			
<b>DIRECTIONS</b>	TAKE 1 TABLET BY MOUTH EVERY 6 HOURS AS NEEDED FOR PAIN . MAX 4 TABLETS PER DAY			



WHITE AND MULTICOLOR  
FRONT: 3202  
BACK: WATSON

**INGREDIENT NAME:** HYDROCODONE (hye-droe-KO-done) and ACETAMINOPHEN (a-seat-a-MIN-oh-ten)

**COMMON USES:** This medicine is a narcotic and analgesic combination used to relieve moderate to moderately severe pain. It may also be used to treat other conditions as determined by your doctor.

**BEFORE USING THIS MEDICINE:** WARNING: THIS MEDICINE CONTAINS ACETAMINOPHEN. Severe and sometimes fatal liver problems, including the need for liver transplant, have been reported with the use of acetaminophen. Most cases of these liver problems occurred in patients taking excessive doses of acetaminophen (more than 4,000 mg per day). Also, patients who developed these liver problems were often using more than 1 medicine that contained acetaminophen. Discuss any questions or concerns with your doctor. Some medicines or medical conditions may interact with this medicine. INFORM YOUR DOCTOR OR PHARMACIST of all prescription and over-the-counter medicine that you are taking. DO NOT TAKE THIS MEDICINE if you are also taking sodium oxybate (XOBY). ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION may be needed if you are taking anticoagulants (eg, warfarin), buprenorphine, isoniazid, linezolid, monoamine oxidase inhibitors (MAOIs) (eg, phenelzine), naltrexone, other narcotic pain medicines (eg, morphine), perampanel, sleep medicines (eg, zolpidem), tricyclic antidepressants (eg, amitriptyline), or medicines that may harm the liver (eg, methotrexate, ketoconazole, certain medicines for HIV infection). Ask your doctor if you are unsure if any of your medicines might harm the liver. DO NOT START OR STOP any medicine without doctor or pharmacist approval. Inform your doctor of any other medical conditions, including a history of liver problems (eg, hepatitis), kidney problems, lung or breathing problems (eg, asthma, chronic obstructive pulmonary disease / COPD), drug or alcohol abuse or dependence, heart problems (eg, cor pulmonale, slow or irregular heartbeat), low blood pressure, stomach or bowel problems (eg, blockage, inflammation, paralysis), gallbladder problems, prostate problems (eg, an enlarged prostate), difficulty urinating, seizures, thyroid problems, adrenal gland problems (eg, Addison disease), mental or mood problems, inflammation of the pancreas (pancreatitis), curvature of the spine (scoliosis), suicidal thoughts or actions, allergies, pregnancy, or breast-feeding. Tell your doctor if you have severe drowsiness, or have or recently have had any head injury, increased pressure in the brain, growths in the brain (eg, tumors), or infection of the brain or nervous system. Tell your doctor if you have recently had surgery (eg, stomach or lower surgery) or are scheduled to have surgery. Tell your doctor if you are in very poor health; are very overweight or are dehydrated; have low blood volume, stomach pain, or constipation; drink alcohol; or are going through withdrawal from alcohol or other substances. USE OF THIS MEDICINE IS NOT RECOMMENDED if you have very slow or difficult breathing, or you are having an asthma attack. Contact your doctor or pharmacist if you have any questions or concerns about taking this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for using this medicine provided by your doctor. TAKE THIS MEDICINE by mouth with or without food. If stomach upset occurs, take with food to reduce stomach irritation. IF YOU HAVE BEEN TAKING THIS MEDICINE REGULARLY or for longer than a few weeks, do not suddenly stop taking it without checking with your doctor. Your doctor may need to gradually lower your dose. SOME OF THESE PRODUCTS CONTAIN SULFITES. Sulfites may cause allergic reaction in some patients (eg, asthma patients). If you have ever had an allergic reaction to sulfites, ask your pharmacist if your product has sulfites in it. STORE THIS MEDICINE at room temperature, between 68 and 77 degrees F (20 and 25 degrees C), away from heat, moisture, and light. Do not store in the bathroom. KEEP THIS MEDICINE out of the reach of children and away from pets. IF YOU MISS A DOSE OF THIS MEDICINE and you are taking it regularly, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.

**CAUTIONS:** DO NOT TAKE THIS MEDICINE if you are allergic to any ingredient in this medicine. BEFORE USING THIS MEDICINE, check with your doctor if you have had an allergic reaction to other narcotic medicines (eg, codeine, morphine, oxycodone). DO NOT EXCEED THE RECOMMENDED DOSE, take more often than prescribed, or use for longer than prescribed without checking with your doctor. Exceeding the recommended dose or taking this medicine for longer than prescribed may be habit-forming. If using this medicine for an extended period of time, DO NOT SUDDENLY STOP taking this medicine without your doctor's approval. When using for an extended period, this medicine may not work as well and may require different dosing. Talk with your doctor if this medicine stops working well. THIS MEDICINE MAY AFFECT CERTAIN LAB TEST RESULTS. Make sure that all of your doctors and laboratory personnel know you are taking this medicine. BEFORE YOU HAVE ANY MEDICAL OR DENTAL TREATMENTS, emergency care, or surgery, tell the doctor or dentist that you are using this medicine. THIS MEDICINE MAY CAUSE DIZZINESS, lightheadedness, or fainting. Alcohol, hot weather, exercise, and fever can increase these effects. To prevent them, sit up or stand slowly, especially in the morning. Also, sit or lie down at the first sign of dizziness, lightheadedness, or weakness. THIS MEDICINE MAY ALSO CAUSE DROWSINESS OR BLURRED VISION. Do not drive, operate machinery, or do anything else that could be dangerous until you know how you react to this medicine. Using this medicine alone, with other medicines, or with alcohol may lessen your ability to drive or to perform other potentially dangerous tasks. DO NOT DRINK ALCOHOL while you are using this medicine. CHECK WITH YOUR DOCTOR before you use medicines that may cause drowsiness (eg, sleep aids, muscle relaxers) while you are using this medicine; it may add to their effects. Ask your pharmacist if you have questions about which medicines may cause drowsiness. THIS MEDICINE CONTAINS ACETAMINOPHEN. Do not take additional acetaminophen for pain or fever without checking with your doctor or pharmacist. Ask your pharmacist if you have questions about which medicines contain acetaminophen. THIS MEDICINE MAY HARM YOUR LIVER. Your risk may be greater if you drink alcohol while you are using this medicine. Talk to your doctor before taking this medicine or other fever reducers if you drink alcohol. CONTACT YOUR DOCTOR RIGHT AWAY if you take more than 4,000 mg of acetaminophen per day, even if you feel well. BEFORE YOU BEGIN TAKING ANY NEW MEDICINE, either prescription or over-the-counter, check with your doctor or pharmacist. CAUTION IS ADVISED WHEN USING THIS MEDICINE IN THE ELDERLY because they may be more sensitive to the effects of the medicine. FOR WOMEN: IF YOU BECOME PREGNANT, discuss with your doctor the benefits and risks of using this medicine during pregnancy. Long-term use of this medicine during pregnancy may cause dependence in the fetus or newborn. Discuss any questions or concerns with your doctor. THIS MEDICINE IS EXCRETED in breast milk. DO NOT BREAST-FEED while taking this medicine.

**POSSIBLE SIDE EFFECTS:** SIDE EFFECTS that may occur while taking this medicine include dizziness, drowsiness, lightheadedness, nausea, or vomiting. If

they continue or are bothersome, check with your doctor. THIS MEDICINE MAY CAUSE CONSTIPATION. Talk with your doctor or pharmacist about using a stool softener or laxative to prevent constipation. It is also important to maintain a diet adequate in fiber, drink plenty of water, and exercise to prevent constipation. If you become constipated while taking this medicine, talk with your doctor or pharmacist. CONTACT YOUR DOCTOR IMMEDIATELY if you experience a change in the amount of urine produced; change or loss in hearing; chest pain; confusion; fainting; fever, chills, or persistent sore throat; mental or mood changes (eg, agitation, anxiety, depression); red, swollen, blistered, or peeling skin; severe or persistent constipation; severe or persistent dizziness, drowsiness, headache, or lightheadedness; shortness of breath; slow, shallow, or difficult breathing; symptoms of liver problems (eg, yellowing of the skin or eyes, pale stools, dark urine, persistent loss of appetite, unusual or severe stomach pain); trouble urinating; unusual bruising or bleeding; unusual tiredness or weakness; or vision changes. AN ALLERGIC REACTION to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, throat, or tongue; or unusual hoarseness. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** IF OVERDOSE IS SUSPECTED, contact your local poison control center or emergency room immediately. Symptoms of overdose may include bluish skin or nails; cold and clammy skin; coma; decreased urination; difficult, shallow, or slow breathing; fainting; limp muscles; loss of consciousness; nausea or vomiting; pinpoint pupils; seizures; severe dizziness, drowsiness, or light-headedness; slow heartbeat; symptoms of liver problems (eg, yellowing of the skin or eyes, pale stools, dark urine, loss of appetite, unusual or severe stomach pain); unusual bruising or bleeding; or unusual sweating or tiredness.

**ADDITIONAL INFORMATION:** If your symptoms do not improve or if they become worse, contact your doctor. DO NOT SHARE THIS MEDICINE with others for whom it was not prescribed. DO NOT USE THIS MEDICINE for other health conditions. CHECK WITH YOUR PHARMACIST about how to dispose of unused medicine.

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

### ROBERT A PLOCK

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

**RX # 0529365-09015**

DATE: 08/06/13

**HYDROCODONE /ACETAMINOPHEN 5-325 TB**

QTY: 120 NO REFILLS - DR. AUTH REQUIRED

Refill NDC:00591-3202-01

Retail Price: \$87.09 Your Insurance Saved You: \$77.09

\$ 10.00

A. PARK, MD  
MFG:WATSON  
SDB/ADO/ADO/ /ADO

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 132186453718093999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Customer  
Receipt

### ROBERT A PLOCK

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

**RX # 0529365-09015**

DATE: 08/06/13

**HYDROCODONE /ACETAMINOPHEN 5-325 TB**

QTY: 120 NO REFILLS - DR. AUTH REQUIRED

Refill NDC:00591-3202-01

Retail Price: \$87.09 Your Insurance Saved You: \$77.09

\$ 10.00

A. PARK, MD  
MFG:WATSON  
SDB/ADO/ADO/ /ADO

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 132186453718093999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951


Duplicate  
Receipt

Pharmacy use only

TUE 9:24PM  
Refill

HYDROCODONE /ACETAMINOPHEN 5-325 TB  
00591-3202-01  
FAST RACK

QTY 120  
30 DRAM



WHITE AND MULTICOLOR  
FRONT: 3202  
BACK: WATSON

SDB/ADO/ADO/ /ADO

# YOUR PERSONAL PRESCRIPTION INFORMATION

## Your Walgreens Pharmacy Location

4702 N Jim Miller Rd  
Dallas, TX 75227-  
(214)388-4951

IF YOU HAVE QUESTIONS ABOUT YOUR PRESCRIPTION, PLEASE CONTACT YOUR WALGREENS PHARMACIST OR CALL 1-800-WALGREENS.

PATIENT ROBERT A PLOCK

BIRTH DATE 07/26/68

MEDICATION IBUPROFEN 800MG TABLETS


QUANTITY 90

DIRECTIONS TAKE 1 TABLET BY MOUTH EVERY 8 HOURS AS NEEDED FOR PAIN. NO MORE THAN 3 TABLETS PER DAY

DOCTOR DR A. PARK

DRUG DESCRIPTION

PATIENT ALLERGIES



WHITE

FRONT: IP 466

INGREDIENT NAME: IBUPROFEN (eye-byoo-PROE-fen)

**COMMON USES:** This medicine is a nonsteroidal anti-inflammatory drug (NSAID) used to treat mild to moderate pain, osteoarthritis, and rheumatoid arthritis. It may also be used to treat other conditions as determined by your doctor.

**BEFORE USING THIS MEDICINE:** WARNING: THE RISK OF SERIOUS AND SOMETIMES FATAL HEART PROBLEMS, HEART ATTACK, AND STROKE may be increased with the use of this medicine. This risk may be increased the longer you use this medicine. Risk may also be higher in patients who have heart problems or who are at risk for heart problems. THIS MEDICINE SHOULD NOT BE USED to treat pain before or after coronary artery heart bypass (CABG) surgery. THE RISK OF SERIOUS AND SOMETIMES FATAL STOMACH AND BOWEL PROBLEMS, including bleeding, ulcers, and holes in the stomach and bowel, is increased while using this medicine. These problems may occur at any time during therapy, with or without symptoms. Elderly patients are at higher risk for serious stomach problems. Ask your doctor or pharmacist for more information about this medicine and its side effects. Some medicines or medical conditions may interact with this medicine. INFORM YOUR DOCTOR OR PHARMACIST of all prescription and over-the-counter medicine that you are taking. DO NOT TAKE THIS MEDICINE if you are also taking heparin or warfarin. ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION may be needed if you are taking serotonin reuptake blocker medicines such as fluoxetine or citalopram, "blood thinners" such as warfarin, bisphosphonates such as alendronate or risedronate, cyclosporine, corticosteroids such as prednisone, high blood pressure medicines (including ACE inhibitors such as captopril, angiotensin II receptor antagonists such as losartan, and beta-blockers such as metoprolol), "water pills" (diuretics such as furosemide, hydrochlorothiazide, triamterene), lithium, methotrexate, or aspirin. DO NOT START OR STOP ANY MEDICINE without doctor or pharmacist approval. Inform your doctor of any other medical conditions including poorly controlled diabetes, dehydration, heart problems (such as heart failure or history of heart attack), swelling of the hands, feet, or ankles (edema), high blood pressure, history of stroke, blood clotting problems, stomach or bowel problems (such as bleeding or ulcers), history of tobacco use or alcohol use, kidney problems, liver problems, blood or bleeding problems (such as anemia), asthma, growths in the nose (nasal polyps), any allergies (especially history of angioedema with symptoms of lip, tongue, throat swelling), pregnancy, or breast-feeding. USE OF THIS MEDICINE IS NOT RECOMMENDED if you have a history of allergy to aspirin or other NSAIDs (e.g., naproxen, celecoxib). USE OF THIS MEDICINE IS NOT RECOMMENDED if you have history of severe kidney disease or if you are going to have or have recently had coronary artery heart bypass (CABG) surgery. Contact your doctor or pharmacist if you have any questions or concerns about taking this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for using this medicine provided by your doctor. This medicine may come with a medication guide. Read it carefully. Ask your doctor, nurse, or pharmacist any questions that you may have about this medicine. TAKE THIS MEDICINE with a full glass (8 oz./240 ml) of water. DO NOT lie down for 30 minutes after taking this medicine. The dosage is based on your medical condition and response to therapy. If repeat doses are needed, they are usually given 6 or 8 hours apart, or as directed by your doctor. When used in children, the dose is based on your child's weight. Read the product instructions to find the appropriate dose for your child's weight. Consult the pharmacist or doctor if you have questions or if you need help in choosing the appropriate dosage form. THIS MEDICINE MAY BE TAKEN WITH FOOD if it upsets your stomach. Taking it with food may not decrease the risk of stomach or bowel problems (such as bleeding or ulcers) that may occur while taking this medicine. Talk with your doctor or pharmacist if you experience persistent stomach upset. STORE THIS MEDICINE at room temperature, away from heat and light. Do not store in the bathroom. IF YOU MISS A DOSE OF THIS MEDICINE, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.

**CAUTIONS:** THIS MEDICINE INCREASES YOUR RISK OF SERIOUS STOMACH OR BOWEL PROBLEMS (such as ulcers and bleeding). This risk is increased if you are elderly or are in poor health, if you have a history of smoking or drinking alcohol, if you take corticosteroid medicines (such as prednisone) or "blood thinners" (such as warfarin), or if you take this medicine for a long period of time. THIS MEDICINE MAY ALSO INCREASE YOUR RISK for certain serious heart and blood vessel problems (such as heart attack and stroke). TAKE THIS MEDICINE EXACTLY AS PRESCRIBED BY YOUR DOCTOR, at the lowest possible dose for the

shortest time needed. Talk with your doctor or pharmacist for further information. DO NOT TAKE THIS MEDICINE IF YOU HAVE HAD A SEVERE ALLERGIC REACTION to aspirin or any medicine containing aspirin or to a nonsteroidal anti-inflammatory drug (such as Feldene, Motrin, Naprosyn, Clinoril). A severe reaction includes a severe rash, hives, breathing difficulties, or dizziness. If you have a question about whether you are allergic to this medicine or if a certain medicine is a nonsteroidal anti-inflammatory drug, contact your doctor or pharmacist. DO NOT EXCEED THE RECOMMENDED DOSE or take this medicine for longer than 10 days for pain or 3 days for fever, unless directed by your doctor. Laboratory and/or medical tests, including blood counts, liver function tests, and kidney function tests, may be performed to monitor your progress or to check for side effects, especially if you are taking this medicine for a long period of time. KEEP ALL DOCTOR AND LABORATORY APPOINTMENTS while you are taking this medicine. DO NOT DRIVE, OPERATE MACHINERY, OR DO ANYTHING ELSE THAT COULD BE DANGEROUS until you know how you react to this medicine. **ALCOHOL WARNING:** If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take this medicine or other pain relievers/fever reducers. BEFORE YOU BEGIN TAKING ANY NEW MEDICINE, either prescription or over-the-counter, check with your doctor or pharmacist. If you are also taking aspirin, as prescribed by your doctor for reasons such as heart attack or stroke prevention (usually these dosages are 81-325 mg per day), continue to take the aspirin and consult your doctor or pharmacist before using this medicine. CAUTION IS ADVISED WHEN USING THIS MEDICINE IN THE ELDERLY because they may be more sensitive to the effects of this medicine, especially the risk of stomach or bowel effects (such as bleeding or ulcers), or kidney effects. FOR WOMEN: USE OF THIS MEDICINE DURING PREGNANCY has resulted in fetal and newborn death. If you think you may be pregnant, contact your doctor immediately. THIS MEDICINE IS EXCRETED IN BREAST MILK. IF YOU ARE OR WILL BE BREAST-FEEDING while you are using this medicine, check with your doctor or pharmacist to discuss the risks to your baby.

**POSSIBLE SIDE EFFECTS:** SIDE EFFECTS, that may go away during treatment, include nausea, vomiting, diarrhea, gas, constipation, indigestion, dizziness, lightheadedness, drowsiness, or headache. If they continue or are bothersome, check with your doctor. CHECK WITH YOUR DOCTOR AS SOON AS POSSIBLE if you experience ringing in ears. CONTACT YOUR DOCTOR IMMEDIATELY if you experience rapid or pounding heartbeat; easy bruising or bleeding; very stiff neck; or mental/mood changes. CONTACT YOUR DOCTOR IMMEDIATELY if you experience sharp or crushing chest pain; sudden shortness of breath; sudden leg pain; sudden severe headache, vomiting, dizziness, or fainting; changes in vision; numbness of an arm or leg; slurred speech; one-sided weakness; sudden unexplained weight gain; change in amount of urine produced; severe or persistent stomach pain; vomit that looks like coffee grounds; black tarry stools; itching, reddened, swollen, blistered, painful, or peeling skin; yellowing of the skin or eyes; dark urine; right-sided tenderness; severe or persistent tiredness; fever, chills, or sore throat; severe or persistent nausea; swelling of hands, ankles, feet, face, lips, eyes, throat, or tongue; difficulty swallowing or breathing; or hoarseness. AN ALLERGIC REACTION TO THIS MEDICINE is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash, itching, swelling, severe dizziness, or trouble breathing. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms of overdose may include severe stomach pain, coffee ground-like vomit, unusually fast or slow heartbeat, trouble breathing, extreme drowsiness, loss of consciousness, and seizures.

**ADDITIONAL INFORMATION:** DO NOT SHARE THIS MEDICINE with others for whom it was not prescribed. DO NOT USE THIS MEDICINE for other health conditions. KEEP THIS MEDICINE out of the reach of children and pets. IF USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME, obtain refills before your supply runs out.

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

ROBERT A PLOCK

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

RX # 0534729-09015

DATE: 08/06/13

IBUPROFEN 800MG TABLETS

QTY: 90 NO REFILLS - DR. AUTH REQUIRED

Copy NDC: 53746-0466-05

Retail Price: \$22.39 Your Insurance Saved You: \$12.39

\$ 10.00

DR A. PARK  
MFG: AMNEAL  
NJT/NJT/NJT /ADO

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 132186452475012999

Walgreens

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Customer  
Receipt

ROBERT A PLOCK

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

RX # 0534729-09015

DATE: 08/06/13

IBUPROFEN 800MG TABLETS

QTY: 90 NO REFILLS - DR. AUTH REQUIRED

Copy NDC: 53746-0466-05

Retail Price: \$22.39 Your Insurance Saved You: \$12.39

\$ 10.00

DR A. PARK  
MFG: AMNEAL  
NJT/NJT/NJT /ADO

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 132186452475012999

Walgreens

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Duplicate  
Receipt

Pharmacy use only

TUE 9:24PM

Copy


IBUPROFEN 800MG TABLETS

53746-0466-05

FAST RACK

QTY 90

40 DRAM



WHITE

FRONT: IP 466

NJT/NJT/NJT /ADO

Med Guide

Do not flush unused medications or pour down a sink or drain.

WIC# 216382



# YOUR PERSONAL PRESCRIPTION INFORMATION

## Your Walgreens Pharmacy Location

4702 N Jim Miller Rd  
Dallas, TX 75227-  
(214)388-4951

IF YOU HAVE QUESTIONS ABOUT YOUR PRESCRIPTION, PLEASE CONTACT YOUR WALGREENS PHARMACIST OR CALL 1-800-WALGREENS.

**PATIENT** ROBERT A PLOCK  
**BIRTH DATE** 07/26/68  
**MEDICATION** TRAMADOL 50MG TABLETS  
**QUANTITY** 120

**DOCTOR** DR A. PARK

**DRUG DESCRIPTION**

**PATIENT  
ALLERGIES**



WHITE  
FRONT: 319

### INGREDIENT NAME: TRAMADOL (TRAM-a-dol)

**COMMON USES:** This medicine is an analgesic used to treat moderate to moderately severe pain.

**BEFORE USING THIS MEDICINE:** Some medicines or medical conditions may interact with this medicine. INFORM YOUR DOCTOR OR PHARMACIST of all prescription and over-the-counter medicine that you are taking. DO NOT TAKE THIS MEDICINE if you are taking sodium oxybate (GHB), or another product that contains tramadol. ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION may be needed if you are taking alpha-2 receptor blockers (eg, yohimbine), anorexiant (eg, phentermine), anticoagulants (eg, warfarin), azole antifungals (eg, ketoconazole), butyrophenones (eg, haloperidol), carbamazepine, cyclobenzaprine, digoxin, furazolidone, linezolid, lithium, loxapine, macrolide antibiotics (eg, erythromycin), certain medicines for mental or mood disorders (eg, olanzapine), molidone, monoamine oxidase inhibitors (MAOIs) (eg, phenelzine, selegiline), nefazodone, opioid pain medicine (eg, codeine, hydrocodone), phenothiazines (eg, promethazine), quinidine, rifampin, serotonin-norepinephrine reuptake inhibitors (SNRIs) (eg, duloxetine), selective serotonin reuptake inhibitors (SSRIs) (eg, fluoxetine, paroxetine), sleeping medicines (eg, zolpidem), St. John's wort, thioxanthenes (eg, thiothixene), tiagabine, tricyclic antidepressants (eg, amitriptyline), or "triptans" (eg, sumatriptan). DO NOT START OR STOP any medicine without doctor or pharmacist approval. Inform your doctor of any other medical conditions, including liver, kidney, or thyroid problems; blockage in the bladder; diabetes; heart problems (eg, cor pulmonale); high blood pressure; pancreas problems; prostate problems; metabolism problems; allergies; pregnancy; or breast-feeding. Tell your doctor if you have or recently have had any head injury, brain injury or tumor, increased pressure in the brain, or infection of the brain or nervous system. Tell your doctor if you have a history of lung or breathing problems (eg, asthma, chronic obstructive pulmonary disease / COPD); seizures (eg, epilepsy); recent stomach or bowel surgery; other stomach or bowel problems (eg, pain, inflammation, ulcers); alcohol or other substance abuse; mood or mental problems (eg, depression); or suicidal thoughts or actions. Tell your doctor if you drink alcohol or if you are going through withdrawal from alcohol or other substances. USE OF THIS MEDICINE IS NOT RECOMMENDED if you have severe or persistent diarrhea caused by antibiotic use; suicidal thoughts or actions; or a history of alcohol or other substance abuse or addiction. USE OF THIS MEDICINE IS NOT RECOMMENDED if you are intoxicated with alcohol, opioids or narcotics (eg, codeine, morphine), or sedatives or sleeping medicines (eg, temazepam, zolpidem). This medicine should not be used in CHILDREN younger than 16 years; safety and effectiveness in these children have not been confirmed. Contact your doctor or pharmacist if you have any questions or concerns about using this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for taking this medicine provided by your doctor. TAKE THIS MEDICINE by mouth with or without food. STORE THIS MEDICINE at 77 degrees F (25 degrees C) away from heat, moisture, and light. Do not store in the bathroom. Brief storage at temperatures between 59 and 86 degrees F (15 and 30 degrees C) is permitted. KEEP THIS MEDICINE out of the reach of children and away from pets. IF YOU MISS A DOSE OF THIS MEDICINE and you are taking it regularly, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. DO NOT take 2 doses at once.

**CAUTIONS:** DO NOT TAKE THIS MEDICINE if you have had an allergic reaction to it or if you are allergic to any ingredient in this product. DO NOT TAKE THIS MEDICINE if you have had a severe allergic reaction to codeine or another opioid (eg, morphine). DO NOT EXCEED THE RECOMMENDED DOSE or take this medicine for longer than prescribed without checking with your doctor. Exceeding the recommended dose or taking this medicine for longer than prescribed may be habit-forming or may increase the risk of seizures or breathing problems. If using this medicine for an extended period of time, DO NOT SUDDENLY STOP taking this medicine without your doctor's approval. Your dose may need to be slowly lowered to avoid side effects. When used for an extended period of time, this medicine may not work as well and may require different dosing. Talk with your doctor if this medicine stops working well. DO NOT TAKE MORE THAN PRESCRIBED. BEFORE YOU HAVE ANY MEDICAL OR DENTAL TREATMENTS, EMERGENCY CARE, OR SURGERY, tell the doctor or dentist that you are using this medicine. DO NOT DRINK ALCOHOL while you are using this medicine. CHECK WITH YOUR DOCTOR BEFORE you use medicines that may cause drowsiness (eg, sleep aids, muscle relaxers, narcotic pain medicines) while you are using this medicine; it may add to their effects. Ask your pharmacist if you

have questions about which medicines may cause drowsiness. This medicine may cause drowsiness or dizziness. DO NOT DRIVE, OPERATE MACHINERY, OR DO ANYTHING ELSE THAT COULD BE DANGEROUS until you know how you react to this medicine. Using this medicine alone, with other medicines, or with alcohol may lessen your ability to drive or to perform other potentially dangerous tasks. Alcohol, hot weather, exercise, or fever may increase dizziness. IF DIZZINESS OCCURS, sit up or stand slowly, especially in the morning. Sit or lie down at the first sign of this effect. THIS MEDICINE MAY INCREASE YOUR RISK OF SEIZURES. Your risk may be greater if you also have certain medical conditions, use certain medicines, or if you use a lot of alcohol. Talk to your doctor to see if you may have a greater risk of seizures while taking this medicine. SEROTONIN SYNDROME is a possibly fatal syndrome that can be caused by this medicine. Your risk may be greater if you take this medicine with certain other medicines (eg, "triptans," MAOIs, antidepressants). Symptoms may include agitation; confusion; hallucinations; coma; fever; fast or irregular heartbeat; tremor; excessive sweating; and nausea, vomiting, or diarrhea. Contact your doctor at once if you have any of these symptoms. BEFORE YOU BEGIN TAKING ANY NEW MEDICINES, either prescription or over-the-counter, check with your doctor or pharmacist. CAUTION IS ADVISED WHEN USING THIS MEDICINE IN THE ELDERLY because they may be more sensitive to the effects of this medicine. FOR WOMEN: This medicine may cause harm to the fetus. IF YOU THINK YOU MAY BE PREGNANT, discuss with your doctor the benefits and risks of using this medicine during pregnancy. THIS MEDICINE IS EXCRETED in breast milk. DO NOT BREAST-FEED while taking this medicine.

**POSSIBLE SIDE EFFECTS:** SIDE EFFECTS that may occur while taking this medicine include constipation; diarrhea; dizziness; drowsiness; dry mouth; headache; indigestion; mildly increased sweating; mild itching; nausea; trouble sleeping; vomiting; weakness. If they continue or are bothersome, check with your doctor. CONTACT YOUR DOCTOR IMMEDIATELY if you experience agitation; chest pain; confusion; difficult or painful urination; excessive sweating; fainting; fast or irregular heartbeat; fever; hallucinations; loss of coordination; mood or mental changes (eg, depression); red, blistered, swollen, or peeling skin; seizures; severe dizziness or lightheadedness; severe nausea, vomiting, or diarrhea; severe or persistent headache; slow or shallow breathing; suicidal thoughts or behaviors; tremor; or vision problems. AN ALLERGIC REACTION to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, or tongue; unusual hoarseness. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your health care provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** IF OVERDOSE IS SUSPECTED, contact your local poison control center or emergency room immediately. Symptoms may include bluish skin; cold, clammy skin; difficult, shallow, or slow breathing; drowsiness leading to unresponsiveness or coma; excessive sweating; limp muscles; pinpoint pupils; seizures; slow or irregular heartbeat.

**ADDITIONAL INFORMATION:** DO NOT SHARE THIS MEDICINE with others for whom it was not prescribed. DO NOT USE THIS MEDICINE for other health conditions. IF YOU WILL BE USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME, be sure to obtain necessary refills before your supply runs out. CHECK WITH YOUR PHARMACIST about how to dispose of unused medicine.

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

### ROBERT A PLOCK

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

**RX # 0534730-09015**

DATE: 09/04/13

### TRAMADOL 50MG TABLETS

QTY: 120 PARTIAL REFILL BEFORE 07/24/14

Refill NDC: 68382-0319-10

Retail Price: \$53.98 Your Insurance Saved You: \$43.98

\$ 10.00

DR A. PARK

PLAN: WPB

MFG: ZYDUS

GROUP# 01960917

SIM/PDR/PDR/ /PDR

CLAIM REF# 132473769246057998

### ROBERT A PLOCK

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

**RX # 0534730-09015**

DATE: 09/04/13

### TRAMADOL 50MG TABLETS

QTY: 120 PARTIAL REFILL BEFORE 07/24/14

Refill NDC: 68382-0319-10

Retail Price: \$53.98 Your Insurance Saved You: \$43.98

\$ 10.00

DR A. PARK

PLAN: WPB

MFG: ZYDUS

GROUP# 01960917

SIM/PDR/PDR/ /PDR

CLAIM REF# 132473769246057998

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Customer  
Receipt

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Duplicate  
Receipt

Pharmacy use only

WED 1:59PM  
Refill

TRAMADOL 50MG TABLETS  
68382-0319-10  
FAST RACK

QTY 120  
20 DRAM



WHITE  
FRONT: 319

SIM/PDR/PDR/ /PDR

Call your doctor for medical advice about side effects.  
You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 216382

# YOUR PERSONAL PRESCRIPTION INFORMATION

## Your Walgreens Pharmacy Location

4702 N Jim Miller Rd  
Dallas, TX 75227-  
(214)388-4951


IF YOU HAVE QUESTIONS ABOUT YOUR PRESCRIPTION, PLEASE CONTACT YOUR WALGREENS PHARMACIST OR CALL 1-800-WALGREENS.

**PATIENT** ROBERT A PLOCK  
**BIRTH DATE** 07/26/68  
**MEDICATION** CYCLOBENZAPRINE 10MG TABLETS  
**QUANTITY** 90  
**DIRECTIONS** TAKE 1 TABLET BY MOUTH EVERY 8 HOURS AS NEEDED FOR SPASMS

**DOCTOR** DR A. PARK

**DRUG DESCRIPTION**

**PATIENT ALLERGIES**



YELLOW  
FRONT: TL177

**INGREDIENT NAME:** CYCLOBENZAPRINE  
(sye-kloe-BEN-za-preen)

**COMMON USES:** This medicine is a muscle relaxant used to treat muscle spasms caused by painful muscle conditions. It should be used along with rest and physical therapy. It may also be used to treat other conditions as determined by your doctor.

**BEFORE USING THIS MEDICINE:** Some medicines or medical conditions may interact with this medicine. INFORM YOUR DOCTOR OR PHARMACIST of all prescription and over-the-counter medicine that you are taking. DO NOT TAKE THIS MEDICINE if you are taking or have taken a monoamine oxidase inhibitor (MAOI) (eg, phenelzine, rasagiline) within the last 14 days. ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION may be needed if you are taking bupropion, tramadol, selective serotonin reuptake inhibitors (SSRIs) (eg, fluoxetine), serotonin norepinephrine reuptake inhibitors (SNRIs) (eg, venlafaxine), anticholinergics (eg, methscopolamine, benztrapine), barbiturates (eg, phenobarbital), meperidine, mibefradil, phenothiazines (eg, chlorpromazine), guanethidine, guanfacine, tricyclic antidepressants (eg, amitriptyline), or verapamil. DO NOT START OR STOP any medicine without doctor or pharmacist approval. Inform your doctor if you have a history of liver problems, heart problems, cerebral palsy, brain or spinal cord disease, stroke, glaucoma, increased pressure in the eye, or trouble urinating. Inform your doctor of any other medical conditions including allergies, pregnancy, or breast-feeding. USE OF THIS MEDICINE IS NOT RECOMMENDED if you have an overactive thyroid, moderate to severe liver problems, certain heart problems (eg, irregular heartbeat, congestive heart failure, heart block, conduction problems), or if you have recently had a heart attack. Contact your doctor or pharmacist if you have any questions or concerns about using this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for using this medicine provided by your doctor. THIS MEDICINE MAY BE TAKEN WITH FOOD if it upsets your stomach. STORE THIS MEDICINE at room temperature between 68 and 77 degrees F (20 and 25 degrees C), away from heat, moisture, and light. KEEP THIS MEDICINE out of the reach of children and away from pets. IF YOU MISS A DOSE OF THIS MEDICINE, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. DO NOT take 2 doses at once.

**CAUTIONS:** DO NOT TAKE THIS MEDICINE if you have had an allergic reaction to it or are allergic to any ingredient in this product. DO NOT STOP USING THIS MEDICINE without first checking with your doctor. DO NOT EXCEED THE RECOMMENDED DOSE or take this medicine for longer than 2 to 3 weeks without checking with your doctor. If your symptoms (such as pain, tenderness, decreased range of motion) do not improve within 2 to 3 weeks or if they become worse, check with your doctor. THIS MEDICINE MAY CAUSE drowsiness, dizziness, or blurred vision. Do not drive, operate machinery, or do anything else that could be dangerous until you know how you react to this medicine. Talk with your doctor before you drink alcohol or use medicines that may cause drowsiness (such as sleep aids, muscle

relaxers) while you are using this medicine. This medicine may add to their effects. Ask your pharmacist if you have questions about which medicines may cause drowsiness. SEROTONIN SYNDROME is a possibly fatal syndrome that can be caused by this medicine. Your risk may be greater if you take this medicine with certain other medicines (eg, "triptans," MAOIs). Symptoms may include agitation; confusion; hallucinations; coma; fever; fast or irregular heartbeat; tremor; excessive sweating; and nausea, vomiting, or diarrhea. Contact your doctor at once if you have any of these symptoms. IF YOU EXPERIENCE DRY MOUTH, use sugarless candy or gum, or melt bits of ice in your mouth. If dry mouth continues for more than 2 weeks, contact your dentist or doctor DO NOT BECOME OVERHEATED IN HOT WEATHER or while you are being active; heatstroke may occur. BEFORE YOU BEGIN TAKING ANY NEW MEDICINE, either prescription or over-the-counter, check with your doctor or pharmacist. CAUTION IS ADVISED WHEN USING THIS MEDICINE IN THE ELDERLY because they may be more sensitive to the effects of the medicine especially confusion, hallucinations, and fast or irregular heartbeat. FOR WOMEN: IF YOU BECOME PREGNANT, discuss with your doctor the benefits and risks of using this medicine during pregnancy. IT IS UNKNOWN IF THIS MEDICINE IS EXCRETED in breast milk. IF YOU ARE OR WILL BE BREAST-FEEDING while you are using this medicine, check with your doctor or pharmacist to discuss the risks to your baby.

**POSSIBLE SIDE EFFECTS:** SIDE EFFECTS that may occur while taking this medicine include constipation, diarrhea, dizziness, drowsiness, dry mouth, fatigue, nausea, nervousness, and stomach pain or upset. If they continue or are bothersome, check with your doctor. CONTACT YOUR DOCTOR IMMEDIATELY if you experience chest pain, confusion, fainting, fast or irregular heartbeat, mental or mood changes, numbness of an arm or a leg, one-sided weakness, seizures, sudden severe stomach pain, severe dizziness or vomiting, speech or vision problems, trouble urinating, or yellowing of the skin or eyes. An allergic reaction to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; trouble breathing; tightness in the chest; swelling of the mouth, face, lips, or tongue; unusual hoarseness. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms of overdose may include agitation; chest pain; coma; confusion; fast or irregular heartbeat; loss of coordination; hallucinations; seizures; severe drowsiness, dizziness, nausea, or headache; slurred speech; tremor; or vomiting.

**ADDITIONAL INFORMATION:** DO NOT SHARE THIS MEDICINE with others for whom it was not prescribed. DO NOT USE THIS MEDICINE for other health conditions. CHECK WITH YOUR PHARMACIST about how to dispose of unused medicine.

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

**ROBERT A PLOCK**  
6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

**RX # 0532215-09015**

DATE: 09/04/13

**CYCLOBENZAPRINE 10MG TABLETS**

QTY: 90 1 REFILL BEFORE 07/24/14

Refill NDC: 59746-0177-10

Retail Price: \$46.99 Your Insurance Saved You: \$36.99

\$ 10.00

DR A. PARK  
MFG: CADISTA  
SIM/PDR/PDR/PDR

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 132473768806077999

**ROBERT A PLOCK**  
6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

**RX # 0532215-09015**

DATE: 09/04/13

**CYCLOBENZAPRINE 10MG TABLETS**

QTY: 90 1 REFILL BEFORE 07/24/14

Refill NDC: 59746-0177-10

Retail Price: \$46.99 Your Insurance Saved You: \$36.99

\$ 10.00

DR A. PARK  
MFG: CADISTA  
SIM/PDR/PDR/PDR

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 132473768806077999

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Customer  
Receipt

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951


Duplicate  
Receipt

Pharmacy use only

WED 1:58PM  
Refill

CYCLOBENZAPRINE 10MG TABLETS  
59746-0177-10  
FAST RACK

QTY 90  
10 DRAM



YELLOW  
FRONT: TL177

SIM/PDR/PDR/PDR

Do not flush unused medications or pour down a sink or drain.

WIC# 216382

Call your doctor for medical advice about side effects.  
You may report side effects to FDA at 1-800-FDA-1088.



# YOUR PERSONAL PRESCRIPTION INFORMATION

## Your Walgreens Pharmacy Location

4702 N Jim Miller Rd  
Dallas, TX 75227-  
(214)388-4951

IF YOU HAVE QUESTIONS ABOUT YOUR PRESCRIPTION, PLEASE CONTACT YOUR WALGREENS PHARMACIST OR CALL 1-800-WALGREENS.

PATIENT ROBERT A PLOCK

BIRTH DATE 07/26/68

MEDICATION CYCLOBENZAPRINE 10MG TABLETS


QUANTITY 90

DIRECTIONS TAKE 1 TABLET BY MOUTH EVERY 8 HOURS AS NEEDED FOR SPASMS

DOCTOR DR A. PARK

DRUG DESCRIPTION

PATIENT ALLERGIES



YELLOW

FRONT: TL177

INGREDIENT NAME: CYCLOBENZAPRINE  
(syne-kloe-BEN-za-preen)

**COMMON USES:** This medicine is a muscle relaxant used to treat muscle spasms caused by painful muscle conditions. It should be used along with rest and physical therapy. It may also be used to treat other conditions as determined by your doctor.

**BEFORE USING THIS MEDICINE:** Some medicines or medical conditions may interact with this medicine. INFORM YOUR DOCTOR OR PHARMACIST of all prescription and over-the-counter medicine that you are taking. DO NOT TAKE THIS MEDICINE if you are taking or have taken a monoamine oxidase inhibitor (MAOI) (eg, phenelzine, rasagiline) within the last 14 days. ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION may be needed if you are taking bupropion, tramadol, selective serotonin reuptake inhibitors (SSRIs) (eg, fluoxetine), serotonin norepinephrine reuptake inhibitors (SNRIs) (eg, venlafaxine), anticholinergics (eg, methscopolamine, benztropine), barbiturates (eg, phenobarbital), meperidine, mibefradil, phenothiazines (eg, chlorpromazine), guanethidine, guanfacine, tricyclic antidepressants (eg, amitriptyline), or verapamil. DO NOT START OR STOP any medicine without doctor or pharmacist approval. Inform your doctor if you have a history of liver problems, heart problems, cerebral palsy, brain or spinal cord disease, stroke, glaucoma, increased pressure in the eye, or trouble urinating. Inform your doctor of any other medical conditions including allergies, pregnancy, or breast-feeding. USE OF THIS MEDICINE IS NOT RECOMMENDED if you have an overactive thyroid, moderate to severe liver problems, certain heart problems (eg, irregular heartbeat, congestive heart failure, heart block, conduction problems), or if you have recently had a heart attack. Contact your doctor or pharmacist if you have any questions or concerns about using this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for using this medicine provided by your doctor. THIS MEDICINE MAY BE TAKEN WITH FOOD if it upsets your stomach. STORE THIS MEDICINE at room temperature between 68 and 77 degrees F (20 and 25 degrees C), away from heat, moisture, and light. KEEP THIS MEDICINE out of the reach of children and away from pets. IF YOU MISS A DOSE OF THIS MEDICINE, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. DO NOT take 2 doses at once.

**CAUTIONS:** DO NOT TAKE THIS MEDICINE if you have had an allergic reaction to it or are allergic to any ingredient in this product. DO NOT STOP USING THIS MEDICINE without first checking with your doctor. DO NOT EXCEED THE RECOMMENDED DOSE or take this medicine for longer than 2 to 3 weeks without checking with your doctor. If your symptoms (such as pain, tenderness, decreased range of motion) do not improve within 2 to 3 weeks or if they become worse, check with your doctor. THIS MEDICINE MAY CAUSE drowsiness, dizziness, or blurred vision. Do not drive, operate machinery, or do anything else that could be dangerous until you know how you react to this medicine. Talk with your doctor before you drink alcohol or use medicines that may cause drowsiness (such as sleep aids, muscle

relaxers) while you are using this medicine. This medicine may add to their effects. Ask your pharmacist if you have questions about which medicines may cause drowsiness. SEROTONIN SYNDROME is a possibly fatal syndrome that can be caused by this medicine. Your risk may be greater if you take this medicine with certain other medicines (eg, "triptans," MAOIs). Symptoms may include agitation; confusion; hallucinations; coma; fever; fast or irregular heartbeat; tremor; excessive sweating; and nausea, vomiting, or diarrhea. Contact your doctor at once if you have any of these symptoms. IF YOU EXPERIENCE DRY MOUTH, use sugarless candy or gum, or melt bits of ice in your mouth. If dry mouth continues for more than 2 weeks, contact your dentist or doctor DO NOT BECOME OVERHEATED IN HOT WEATHER or while you are being active; heatstroke may occur. BEFORE YOU BEGIN TAKING ANY NEW MEDICINE, either prescription or over-the-counter, check with your doctor or pharmacist. CAUTION IS ADVISED WHEN USING THIS MEDICINE IN THE ELDERLY because they may be more sensitive to the effects of the medicine especially confusion, hallucinations, and fast or irregular heartbeat. FOR WOMEN: IF YOU BECOME PREGNANT, discuss with your doctor the benefits and risks of using this medicine during pregnancy. IT IS UNKNOWN IF THIS MEDICINE IS EXCRETED in breast milk. IF YOU ARE OR WILL BE BREAST-FEEDING while you are using this medicine, check with your doctor or pharmacist to discuss the risks to your baby.

**POSSIBLE SIDE EFFECTS:** SIDE EFFECTS that may occur while taking this medicine include constipation, diarrhea, dizziness, drowsiness, dry mouth, fatigue, nausea, nervousness, and stomach pain or upset. If they continue or are bothersome, check with your doctor. CONTACT YOUR DOCTOR IMMEDIATELY if you experience chest pain, confusion, fainting, fast or irregular heartbeat, mental or mood changes, numbness of an arm or a leg, one-sided weakness, seizures, sudden severe stomach pain, severe dizziness or vomiting, speech or vision problems, trouble urinating, or yellowing of the skin or eyes. An allergic reaction to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; trouble breathing; tightness in the chest; swelling of the mouth, face, lips, or tongue; unusual hoarseness. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms of overdose may include agitation; chest pain; coma; confusion; fast or irregular heartbeat; loss of coordination; hallucinations; seizures; severe drowsiness, dizziness, nausea, or headache; slurred speech; tremor; or vomiting.

**ADDITIONAL INFORMATION:** DO NOT SHARE THIS MEDICINE with others for whom it was not prescribed. DO NOT USE THIS MEDICINE for other health conditions. CHECK WITH YOUR PHARMACIST about how to dispose of unused medicine.

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

ROBERT A PLOCK

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

RX # 0532215-09015

DATE: 07/24/13

CYCLOBENZAPRINE 10MG TABLETS

QTY: 90 2 REFILLS BEFORE 07/24/14

New

NDC: 59746-0177-10

Retail Price: \$46.99

Your Insurance Saved You: \$36.99

\$ 10.00

DR A. PARK  
MFG: CADISTA  
NJT/NJT/NJT / NJT

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 132055950731079999

Walgreens

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951

Customer  
Receipt

ROBERT A PLOCK

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

RX # 0532215-09015

DATE: 07/24/13

CYCLOBENZAPRINE 10MG TABLETS

QTY: 90 2 REFILLS BEFORE 07/24/14

New

NDC: 59746-0177-10

Retail Price: \$46.99

Your Insurance Saved You: \$36.99

\$ 10.00

DR A. PARK  
MFG: CADISTA  
NJT/NJT/NJT / NJT

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 132055950731079999

Walgreens

4702 N JIM MILLER RD DALLAS, TX 75227  
PH: (214)388-4951


Duplicate  
Receipt

Pharmacy use only

THU 8:00AM  
New

CYCLOBENZAPRINE 10MG TABLETS  
59746-0177-10  
FAST RACK

QTY 90  
10 DRAM



YELLOW  
FRONT: TL177

NJT/NJT/NJT / NJT

Do not flush unused medications or pour down a sink or drain.

WIC# 216382

# Walgreens

#09015 4702 N JIM MILLER RD  
DALLAS, TX 75227  
214-388-4951

860 4733 0091 09/25/2013 6:38 PM

FSA RX 0544508 10.00  
CLM RF# 132683403570029998  
FSA RX 0544507 10.00  
CLM RF# 132683398818059999

TOTAL 20.00  
VISA ACCT 2677 20.00  
CHANGE .00

TOTAL FSA ITEMS 0.00  
TOTAL RX ITEMS 20.00  
TOTAL FSA AND RX ITEMS 20.00

APPROVED FSA/HRA AMOUNT 0.00

THANK YOU FOR SHOPPING AT WALGREENS

YOU COULD HAVE EARNED AN ADDITIONAL 1000  
POINTS BY USING YOUR BALANCE REWARDS  
CARD TODAY. RESTRICTIONS APPLY, SEE  
PROGRAM RULES FOR DETAILS.

NOT A MEMBER? JOIN NOW AT ANY REGISTER,  
OR GO TO WALGREENS.COM/BALANCE.  
ENROLLING IS QUICK, EASY AND FREE!  
REDEEM AT THE REGISTER WITH AS FEW AS  
5000 POINTS.

RFN# 0901-5914-7335-1309-2503



# Walgreens

#09015 4702 N JIM MILLER RD  
DALLAS, TX 75227  
214-388-4951

860 8669 0091 07/27/2013 3:19 PM

FSA RX 0532215 10.00  
CLM RF# 132055950731079999

TOTAL 10.00  
VISA ACCT 7727 10.00  
CHANGE .00

TOTAL FSA ITEMS 0.00  
TOTAL RX ITEMS 10.00  
TOTAL FSA AND RX ITEMS 10.00

APPROVED FSA/HRA AMOUNT 0.00

THANK YOU FOR SHOPPING AT WALGREENS

YOU COULD HAVE EARNED AN ADDITIONAL 500  
POINTS BY USING YOUR BALANCE REWARDS  
CARD TODAY. RESTRICTIONS APPLY, SEE  
PROGRAM RULES FOR DETAILS.

NOT A MEMBER? JOIN NOW AT ANY REGISTER,  
OR GO TO WALGREENS.COM/BALANCE.  
ENROLLING IS QUICK, EASY AND FREE!  
REDEEM AT THE REGISTER WITH AS FEW AS  
5000 POINTS.

RFN# 0901-5918-6695-1307-2703



# Walgreens

#09015 4702 N JIM MILLER RD  
DALLAS, TX 75227  
214-388-4951

845 3012 0091 09/09/2013 8:14 AM

FSA RX 0532215 10.00  
CLM RF# 132473768806077999  
FSA RX 0534730 10.00  
CLM RF# 132473769246057998

REWARDS BONUS EARNED 1000

TOTAL 20.00  
VISA ACCT 7727 20.00  
CHANGE .00

TOTAL FSA ITEMS 0.00  
TOTAL RX ITEMS 20.00  
TOTAL FSA AND RX ITEMS 20.00

APPROVED FSA/HRA AMOUNT 0.00

THANK YOU FOR SHOPPING AT WALGREENS

REDEEM AT THE REGISTER WITH AS FEW AS  
5000 POINTS. RESTRICTIONS APPLY, SEE  
PROGRAM RULES FOR DETAILS. PLEASE GO  
TO WALGREENS.COM/BALANCE.

RFN# 0901-5913-0124-1309-0903



\*\*\*\*\*



POINT BALANCE 11500

BALANCE REWARDS ACCT # \*\*\*\*\*6625

OPENING BALANCE 10500  
EARNED THIS VISIT 1000  
CLOSING BALANCE 11500

POINTS TO NEXT AWARD 6500

\*\*\*\*\*

OTHER  
OTHER  
OTHER



# YOUR PERSONAL PRESCRIPTION INFORMATION

## Your Walgreens Pharmacy Location

4702 N Jim Miller Rd  
Dallas, TX 75227-  
(214)388-4951


IF YOU HAVE QUESTIONS ABOUT YOUR PRESCRIPTION, PLEASE CONTACT YOUR WALGREENS PHARMACIST OR CALL 1-800-WALGREENS.

**PATIENT** ROBERT A PLOCK  
**BIRTH DATE** 07/26/68  
**MEDICATION** CYCLOBENZAPRINE 10MG TABLETS  
**QUANTITY** 90  
**DIRECTIONS** TAKE 1 TABLET BY MOUTH EVERY 8 HOURS AS NEEDED FOR SPASMS

**DOCTOR** DR A. PARK

**DRUG DESCRIPTION**

**PATIENT ALLERGIES**



YELLOW  
FRONT: TL177

**INGREDIENT NAME:** CYCLOBENZAPRINE  
(sye-kloe-BEN-za-preen)

**COMMON USES:** This medicine is a muscle relaxant used to treat muscle spasms caused by painful muscle conditions. It should be used along with rest and physical therapy. It may also be used to treat other conditions as determined by your doctor.

**BEFORE USING THIS MEDICINE:** Some medicines or medical conditions may interact with this medicine. INFORM YOUR DOCTOR OR PHARMACIST of all prescription and over-the-counter medicine that you are taking. DO NOT TAKE THIS MEDICINE if you are taking or have taken a monoamine oxidase inhibitor (MAOI) (eg, phenelzine, rasagiline) within the last 14 days. ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION may be needed if you are taking bupropion, tramadol, selective serotonin reuptake inhibitors (SSRIs) (eg, fluoxetine), serotonin norepinephrine reuptake inhibitors (SNRIs) (eg, venlafaxine), anticholinergics (eg, methscopolamine, benztropine), barbiturates (eg, phenobarbital), meperidine, mibefradil, phenothiazines (eg, chlorpromazine), guanethidine, guanfacine, tricyclic antidepressants (eg, amitriptyline), or verapamil. DO NOT START OR STOP any medicine without doctor or pharmacist approval. Inform your doctor if you have a history of liver problems, heart problems, cerebral palsy, brain or spinal cord disease, stroke, glaucoma, increased pressure in the eye, or trouble urinating. Inform your doctor of any other medical conditions including allergies, pregnancy, or breast-feeding. USE OF THIS MEDICINE IS NOT RECOMMENDED if you have an overactive thyroid, moderate to severe liver problems, certain heart problems (eg, irregular heartbeat, congestive heart failure, heart block, conduction problems), or if you have recently had a heart attack. Contact your doctor or pharmacist if you have any questions or concerns about using this medicine.

**HOW TO USE THIS MEDICINE:** Follow the directions for using this medicine provided by your doctor. THIS MEDICINE MAY BE TAKEN WITH FOOD if it upsets your stomach. STORE THIS MEDICINE at room temperature between 68 and 77 degrees F (20 and 25 degrees C), away from heat, moisture, and light. KEEP THIS MEDICINE out of the reach of children and away from pets. IF YOU MISS A DOSE OF THIS MEDICINE, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. DO NOT take 2 doses at once.

**CAUTIONS:** DO NOT TAKE THIS MEDICINE if you have had an allergic reaction to it or are allergic to any ingredient in this product. DO NOT STOP USING THIS MEDICINE without first checking with your doctor. DO NOT EXCEED THE RECOMMENDED DOSE or take this medicine for longer than 2 to 3 weeks without checking with your doctor. If your symptoms (such as pain, tenderness, decreased range of motion) do not improve within 2 to 3 weeks or if they become worse, check with your doctor. THIS MEDICINE MAY CAUSE drowsiness, dizziness, or blurred vision. Do not drive, operate machinery, or do anything else that could be dangerous until you know how you react to this medicine. Talk with your doctor before you drink alcohol or use medicines that may cause drowsiness (such as sleep aids, muscle

relaxers) while you are using this medicine. This medicine may add to their effects. Ask your pharmacist if you have questions about which medicines may cause drowsiness. SEROTONIN SYNDROME is a possibly fatal syndrome that can be caused by this medicine. Your risk may be greater if you take this medicine with certain other medicines (eg, "triptans," MAOIs). Symptoms may include agitation; confusion; hallucinations; coma; fever; fast or irregular heartbeat; tremor; excessive sweating; and nausea, vomiting, or diarrhea. Contact your doctor at once if you have any of these symptoms. IF YOU EXPERIENCE DRY MOUTH, use sugarless candy or gum, or melt bits of ice in your mouth. If dry mouth continues for more than 2 weeks, contact your dentist or doctor DO NOT BECOME OVERHEATED IN HOT WEATHER or while you are being active; heatstroke may occur. BEFORE YOU BEGIN TAKING ANY NEW MEDICINE, either prescription or over-the-counter, check with your doctor or pharmacist. CAUTION IS ADVISED WHEN USING THIS MEDICINE IN THE ELDERLY because they may be more sensitive to the effects of the medicine especially confusion, hallucinations, and fast or irregular heartbeat. FOR WOMEN: IF YOU BECOME PREGNANT, discuss with your doctor the benefits and risks of using this medicine during pregnancy. IT IS UNKNOWN IF THIS MEDICINE IS EXCRETED in breast milk. IF YOU ARE OR WILL BE BREAST-FEEDING while you are using this medicine, check with your doctor or pharmacist to discuss the risks to your baby.

**POSSIBLE SIDE EFFECTS:** SIDE EFFECTS that may occur while taking this medicine include constipation, diarrhea, dizziness, drowsiness, dry mouth, fatigue, nausea, nervousness, and stomach pain or upset. If they continue or are bothersome, check with your doctor. CONTACT YOUR DOCTOR IMMEDIATELY if you experience chest pain, confusion, fainting, fast or irregular heartbeat, mental or mood changes, numbness of an arm or a leg, one-sided weakness, seizures, sudden severe stomach pain, severe dizziness or vomiting, speech or vision problems, trouble urinating, or yellowing of the skin or eyes. An allergic reaction to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash; hives; itching; trouble breathing; tightness in the chest; swelling of the mouth, face, lips, or tongue; unusual hoarseness. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**OVERDOSE:** If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms of overdose may include agitation; chest pain; coma; confusion; fast or irregular heartbeat; loss of coordination; hallucinations; seizures; severe drowsiness, dizziness, nausea, or headache; slurred speech; tremor; or vomiting.

**ADDITIONAL INFORMATION:** DO NOT SHARE THIS MEDICINE with others for whom it was not prescribed. DO NOT USE THIS MEDICINE for other health conditions. CHECK WITH YOUR PHARMACIST about how to dispose of unused medicine.

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

**RX # 0544508-09015**

DATE: 09/25/13

**CYCLOBENZAPRINE 10MG TABLETS**

QTY: 90 NO REFILLS - DR. AUTH REQUIRED

New NDC: 59746-0177-10

Retail Price: \$46.99 Your Insurance Saved You: \$36.99

\$ 10.00

DR A. PARK  
MFG: CADISTA  
SIM/DBP/DBP/DBP

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 132683403570029998

**ROBERT A PLOCK**

6827 Latta Pkwy, Dallas, TX 75227  
(214)799-7775

**RX # 0544508-09015**

DATE: 09/25/13

**CYCLOBENZAPRINE 10MG TABLETS**

QTY: 90 NO REFILLS - DR. AUTH REQUIRED

New NDC: 59746-0177-10

Retail Price: \$46.99 Your Insurance Saved You: \$36.99

\$ 10.00

DR A. PARK  
MFG: CADISTA  
SIM/DBP/DBP/DBP

PLAN: WPB  
GROUP# 01960917  
CLAIM REF# 132683403570029998

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227-  
PH: (214)388-4951

Customer  
Receipt

Pharmacy use only

**Walgreens**

4702 N JIM MILLER RD DALLAS, TX 75227-  
PH: (214)388-4951

Duplicate  
Receipt

WED 12:45PM  
New

CYCLOBENZAPRINE 10MG TABLETS  
59746-0177-10  
FAST RACK

QTY 90  
10 DRAM



YELLOW  
FRONT: TL177

SIM/DBP/DBP/DBP

Call your doctor for medical advice about side effects.  
You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 216382



Everyday FREE Shipping at \$25 or more. [See details >](#) 

**Walgreens**

 Balance Rewards

Savings & Deals

Weekly Ad

Store Locator

Healthcare Clinic

[Home](#) > [Account Home](#) > [Order Status & History](#)

## In-Store Order Details

**Robert Plock**

[Account Home](#)

[Pharmacy & Health](#)

**Orders & Activity**

[Balance Rewards](#)

[Personal Settings](#)


[< Back to orders](#)

### Prescription Order Details

Review your order details below.

**Rx#** 545313-9015

**Order Date:** 09/28/2013

Rx Description	Insurance	Delivery type	
<b>Status</b> 			<b>Price</b>
<b>OXYCODONE/ACETAMINOPHEN 10-325MG TB</b> Qty:120 Prescription Number: 0545313-09015  PARK Prescription for Robert Plock	WPB	Pickup  Pick up time: N/A  Pickup Location: 4702 N Jim Miller Rd Dallas, TX 75227 214-388-4951	
Picked up			\$10.00
Filled Date on 09/28/2013			
<b>Payment Details:</b>		<b>Order Total:</b>	<b>\$10.00</b>

[< Back to orders](#)